

LIFTING HARMONIC MORPHISMS I: METRIZED COMPLEXES AND BERKOVICH SKELETA

OMID AMINI, MATTHEW BAKER, ERWAN BRUGALLÉ, AND JOSEPH RABINOFF

ABSTRACT. Let K be an algebraically closed, complete non-Archimedean field. The purpose of this paper is to carefully study the extent to which finite morphisms of algebraic K -curves are controlled by certain combinatorial objects, called *skeleta*. A skeleton is a metric graph embedded in the Berkovich analytification of X . A skeleton has the natural structure of a metrized complex of curves. We prove that a finite morphism of K -curves gives rise to a finite harmonic morphism of a suitable choice of skeleta. We use this to give analytic proofs of stronger “skeletonized” versions of some foundational results of Liu-Lorenzini, Coleman, and Liu on simultaneous semistable reduction of curves. We then consider the inverse problem of lifting finite harmonic morphisms of metrized complexes to morphisms of curves over K . We prove that every tamely ramified finite harmonic morphism of Λ -metrized complexes of k -curves lifts to a finite morphism of K -curves. If in addition the ramification points are marked, we obtain a complete classification of all such lifts along with their automorphisms. This generalizes and provides new analytic proofs of earlier results of Saïdi and Wewers. As an application, we discuss the relationship between harmonic morphisms of metric graphs and induced maps between component groups of Néron models, providing a negative answer to a question of Ribet motivated by number theory.

This article is the first in a series of two. The second article contains several applications of our lifting results to questions about lifting morphisms of tropical curves.

Throughout this paper, unless explicitly stated otherwise, K denotes a complete algebraically closed non-Archimedean field with nontrivial valuation $\text{val} : K \rightarrow \mathbf{R} \cup \{\infty\}$. Its valuation ring is denoted R , its maximal ideal is \mathfrak{m}_R , and the residue field is $k = R/\mathfrak{m}_R$. We denote the value group of K by $\Lambda = \text{val}(K^\times) \subset \mathbf{R}$.

1. INTRODUCTION

This article is the first in a series of two. The second, entitled *Lifting harmonic morphisms II: tropical curves and metrized complexes*, will be cited as [ABBR14]; references of the form “Theorem II.1.1” will refer to Theorem 1.1 in [ABBR14].

1.1. The purpose of this paper is to carefully study the extent to which finite morphisms of algebraic K -curves are controlled by certain combinatorial objects, called *skeleta*. Let X be a smooth, proper, connected K -curve. Roughly speaking, a *triangulation* $(X, V \cup D)$ of X (with respect to a finite set of punctures $D \subset X(K)$) is a finite set V of points in the Berkovich analytification X^{an} of X whose removal partitions X^{an} into open balls and finitely many open annuli (with the punctures belonging to distinct open balls). Triangulations of (X, D) are naturally in one-to-one correspondence with semistable models \mathfrak{X} of (X, D) : see Section 5. A triangulation $(X, V \cup D)$ gives rise to a skeleton $\Sigma(X, V \cup D)$ of X . The skeleton of a triangulated punctured curve is the dual graph of the special fiber \mathfrak{X}_k of the corresponding semistable model, equipped with a canonical metric, along with completed infinite rays in the directions of the punctures. There are many skeleta in X^{an} , although if $X \setminus D$ is hyperbolic (i.e., has negative Euler characteristic $\chi(X, D) = 2 - 2g(X) - \#D$), then there exists a unique minimal skeleton. A skeleton of (X, D) is by definition a subset $\Sigma \subset X^{\text{an}}$ of the form $\Sigma(X, V \cup D)$ for some triangulation of X .

1.2. Skeletal simultaneous semistable reduction. First we will prove that finite morphisms of curves induce morphisms of skeleta:

The authors thank Amaury Thuillier and Antoine Ducros for their help with some of the descent arguments in (5.2). We are grateful to Andrew Obus for a number of useful comments based on a careful reading of the first arXiv version of this manuscript. M.B. was partially supported by NSF grant DMS-1201473. E.B. was partially supported by the ANR-09-BLAN-0039-01.

Theorem A. *Let $\varphi : X' \rightarrow X$ be a finite morphism of smooth, proper, connected K -curves and let $D \subset X(K)$ be a finite set. There exists a skeleton $\Sigma \subset X$ such that $\Sigma' := \varphi^{-1}(\Sigma)$ is a skeleton of X' . For any such Σ the map $\varphi : \Sigma' \rightarrow \Sigma$ is a finite harmonic morphism of metric graphs.*

See Corollaries 4.18 and 4.28. Harmonic morphisms of graphs are defined in Section 2. Due to the close relationship between semistable models and skeleta, Theorem A should be interpreted as a very strong simultaneous semistable reduction theorem. In fact, we will show how to formally derive from Theorem A the simultaneous semistable reduction theorems of Liu–Lorenzini [LL99], Coleman [Col03], and Liu [Liu06]. Our version of these results hold over any non-Archimedean field K_0 , not assumed to be discretely valued. As an example, the following weak form of Liu’s theorem is a consequence of Theorem A:

Corollary. *Let X, X' be smooth, proper, geometrically connected curves over a non-Archimedean field K_0 and let $\varphi : X' \rightarrow X$ be a finite morphism. Then there exists a finite, separable extension K_1 of K_0 and semistable models $\mathfrak{X}, \mathfrak{X}'$ of the curves X_{K_1}, X'_{K_1} , respectively, such that φ_{K_1} extends to a **finite** morphism $\mathfrak{X}' \rightarrow \mathfrak{X}$.*

We discuss simultaneous semistable reduction theorems in Section 5. We wish to emphasize that Theorem A does not follow from any classical simultaneous semistable reduction theorem, in that there exist finite morphisms $\varphi : \mathfrak{X}' \rightarrow \mathfrak{X}$ of semistable models such that the inverse image of the corresponding skeleton of X is not a skeleton of X' . (See, however, [CKK] where a skeletal version of Liu’s theorem is derived from Liu’s method of proof in the discretely valued case.) Our proof of Theorem A is entirely analytic, resting on an analysis of morphisms between open annuli and open balls; in particular it makes almost no reference to semistable reduction theory, and is therefore quite different from Liu and Liu–Lorenzini’s approach.

A skeletal simultaneous semistable reduction theorem is important precisely when one wants to obtain a well-behaved morphism of graphs from a finite morphism of curves. This is crucial for obtaining the spectral lower bound on the gonality in [CKK], for instance.

1.3. Lifting harmonic morphisms. Our second goal is in a sense inverse to the first: we wish to lift finite harmonic morphisms of metric graphs to finite morphisms of curves. More precisely, let (X, D) be a punctured K -curve as above, let Σ be a skeleton, and let $\bar{\varphi} : \Sigma' \rightarrow \Sigma$ be a finite harmonic morphism of metric graphs. It is natural to ask whether there exists a curve X' and a finite morphism $\varphi : X' \rightarrow X$ such that $\varphi^{-1}(\Sigma)$ is a skeleton of X' and $\varphi^{-1}(\Sigma) \cong \Sigma'$ as metric graphs over Σ . In general the answer is “no”: there are subtle “Hurwitz obstructions” to such a lift, as described below. One solution is to enrich Σ, Σ' with the extra structure of *metrized complexes of curves*. A metrized complex of curves is basically a metric graph Γ and for every (finite) vertex $p \in \Gamma$, the data of a smooth, proper k -curve C_p , along with an identification of the edges adjacent to p with distinct k -points of C_p . There is a notion of a finite harmonic morphism of metrized complexes of curves, which consists of a finite harmonic morphism $\varphi : \Gamma' \rightarrow \Gamma$ of underlying metric graphs, and for every vertex $p' \in \Gamma'$ with $p = \varphi(p')$, a finite morphism $\varphi_{p'} : C_{p'} \rightarrow C_p$, satisfying various compatibility conditions.

Now let $\Sigma = \Sigma(X, V \cup D)$ for a triangulated punctured curve $(X, V \cup D)$. We will show that Σ is naturally a metrized complex of curves, and that finite harmonic morphisms $\Sigma' \rightarrow \Sigma$ of metrized complexes do lift to finite morphisms of curves, under a mild tameness hypothesis. Moreover we have very precise control over the set of such lifts entirely in terms of the morphism of metrized complexes. The main lifting theorem, stated somewhat imprecisely, is as follows:

Theorem B. (Lifting theorem) *Let $(X, V \cup D)$ be a triangulated punctured K -curve, let Σ be its skeleton, and let $\Sigma' \rightarrow \Sigma$ be a tame covering of metrized complexes of curves. Then there exists a curve X' and a finite morphism $\varphi : X' \rightarrow X$, branched only over D , such that $\varphi^{-1}(\Sigma)$ is a skeleton of $(X', \varphi^{-1}(D))$ and $\varphi^{-1}(\Sigma) \cong \Sigma'$ as metrized complexes of curves over Σ . There are finitely many such lifts X' up to X -isomorphism. There are explicit descriptions of the set of lifts and the automorphism group of each lift (as a cover of X) in terms of the morphism $\Sigma' \rightarrow \Sigma$.*

See Theorems 7.4, 7.7, and 7.10 for the precise statements. Tame coverings are defined in Definition 2.21. The hardest part of the proof of Theorem B is a local lifting result, Theorem 6.18, in which the target curve X is replaced by a neighborhood of a vertex $p \in \Sigma$. Theorem 6.18 eventually reduces to classical results about the tamely ramified étale fundamental group.

It is worth mentioning that given a metrized complex of curves \mathcal{C} with edge lengths contained in Λ , it is not hard to construct a triangulated punctured K -curve $(X, V \cup D)$ such that $\Sigma(X, V \cup D) \cong \mathcal{C}$. See Theorem 3.24.

1.4. Our lifting results extend several theorems in the literature, as well as making them more precise. Saïdi [Sai97, Théorème 3.7] proves a version of the lifting theorem for semistable formal curves (without punctures) over a complete discrete valuation ring. His methods also make use of the tamely ramified étale fundamental group and analytic gluing arguments. Wewers [Wew99] works more generally with marked curves over a complete Noetherian local ring using deformation-theoretic arguments, proving that every tamely ramified admissible cover of a marked semistable curve over the residue field of a complete Noetherian local ring lifts. Wewers classifies the possible lifts (fixing, as we do, a lift of the target) in terms of certain non-canonical “deformation data” depending on compatible choices of formal coordinates at the nodes on both the source and target curves.

Our results have several advantages compared to these previous approaches. For one, we are able to work over non-Noetherian rank-1 valuation rings, and descend the lifting theorems *a posteriori* to an essentially arbitrary non-Archimedean field: see (7.11). The real novelty, however, is the systematic use of metrized complexes of curves in the formulation of the lifting theorems, in particular the classification of the set of lifts X' of $\Sigma' \rightarrow \Sigma$. For example, in our situation, Wewers’ non-canonical deformation data are replaced by certain *canonical* gluing data, which means that the automorphism group of the morphism of metrized complexes acts naturally on the set of gluing data (but not on the set of deformation data). This is what allows us to classify lifts up to isomorphism as covers of the target curve, as well as determine the automorphism group of such a lift. Saïdi’s lifting theorem is more similar to ours, with generators of inertia groups over nodal points playing the role of gluing data. It is not clear to the authors whether his methods can be extended to give a classification of lifts.

The question of lifting (and classification of all possible liftings) in the wildly ramified case is more subtle and cannot be guaranteed in general. See Remark 7.6.

1.5. Homomorphisms of component groups. We will discuss in Section 8 the relationship between harmonic morphisms of metric graphs and component groups of Néron models, connecting our lifting results to natural questions in arithmetic algebraic geometry. The idea is as follows. Let X be a smooth, proper, geometrically connected curve over a discretely valued field K_0 , and suppose that X admits a semistable model \mathfrak{X} over the valuation ring R_0 of K_0 . Let J be the Jacobian of X and let Φ_X be the group of connected components of the special fiber of the Néron model of J .

Given a metric graph Γ with integer edge lengths, there is a naturally associated abelian group $\text{Jac}_{\text{reg}}(\Gamma)$, called its *regularized Jacobian*. The association $\Gamma \mapsto \text{Jac}_{\text{reg}}(\Gamma)$ satisfies Picard and Albanese functoriality: that is, given a finite harmonic morphism $\Gamma' \rightarrow \Gamma$ there are natural push-forward and pull-back homomorphisms $\text{Jac}_{\text{reg}}(\Gamma') \rightarrow \text{Jac}_{\text{reg}}(\Gamma)$ and $\text{Jac}_{\text{reg}}(\Gamma) \rightarrow \text{Jac}_{\text{reg}}(\Gamma')$.

A series of theorems of Raynaud, translated into our language, imply that $\Phi_X \cong \text{Jac}_{\text{reg}}(\Sigma_{\mathfrak{X}})$ canonically, where $\Sigma_{\mathfrak{X}}$ is the skeleton induced by the semistable model \mathfrak{X} . We observe that this isomorphism respects both Picard and Albanese functoriality.

For modular curves such as $X_0(N)$ with N prime, the minimal regular model has a special fiber consisting of two projective lines intersecting transversely. Maps from such curves to elliptic curves, and their induced maps on component groups of Néron models, play an important role in arithmetic geometry; for example, Ribet’s work establishing that the Shimura-Taniyama conjecture implies Fermat’s Last Theorem was motivated by the failure of such induced maps on component groups to be surjective in general. (The cokernel of such maps controls congruences between modular forms.) Ribet noticed that if one considers instead the case where the target curve has genus at least 2, then in

all examples he knew of, the induced map on component groups was surjective, and in personal communication with the second author, he asked whether this was a general property of curves whose reduction looks like that of $X_0(N)$. As a concrete application of Theorem B, the next proposition provides a negative answer to this question.

Proposition C. *There exists a finite morphism $f : X' \rightarrow X$ of semistable curves over a discretely valued field K_0 with $g(X) \geq 2$ such that:*

- *the special fiber of the minimal regular model of X' consists of two projective lines intersecting transversely;*
- *the induced map $f_* : \Phi_{X'} \rightarrow \Phi_X$ on component groups of Néron models of Jacobians is not surjective.*

In order to prove Proposition C, we proceed as follows. As component groups can be calculated in terms of metric graphs, one first finds a finite harmonic morphism of metric graphs $\varphi : \Gamma' \rightarrow \Gamma$ such that Γ' has two vertices and at least three edges, all of which have length one, and such that $\varphi_* : \text{Jac}_{\text{reg}}(\Gamma') \rightarrow \text{Jac}_{\text{reg}}(\Gamma)$ is not surjective. One then enriches φ to a morphism of metrized complexes of curves, with rational residue curves, and uses Theorem 3.24 and Theorem B (along with a descent argument) to lift this to a finite morphism of curves $X' \rightarrow X$. Such a morphism satisfies the conditions of Proposition C.

1.6. Applications to tropical lifting theorems. Tropical geometry has many applications to algebraic geometry, in particular enumerative algebraic geometry. The basic strategy is to associate a combinatorial “tropical” object to an algebraic object, and prove that counting the former is somehow equivalent to counting the latter. An important ingredient in any such argument is a tropical lifting theorem, which is a precise description of the algebraic lifts of a given tropical object.

Many of the intended applications of our lifting results are contained in the second paper [ABBR14], in which we prove a number of tropical lifting theorems. The skeleton of a curve, viewed as a metric graph, is a tropical object associated to the curve; our lifting results can therefore be interpreted as tropical lifting theorems for morphisms of metric graphs, in the following sense. Recall that $\Lambda \subset \mathbf{R}$ is the value group of K . A Λ -metric graph is a metric graph Γ whose edge lengths are contained in Λ . We say that a finite harmonic morphism $\bar{\varphi} : \Gamma' \rightarrow \Gamma$ of Λ -metric graphs is *liftable* provided that there exists a finite morphism of K -curves $\varphi : X' \rightarrow X$, a set of punctures $D \subset X(K)$, and a skeleton Σ of (X, D) such that $\Sigma \cong \Gamma$ as metric graphs, $\varphi^{-1}(\Sigma)$ is a skeleton of $(X', \varphi^{-1}(D))$, and $\Sigma' \cong \Gamma'$ as metric graphs over $\Sigma \cong \Gamma$. Theorem B (along with Theorem 3.24) shows that the obstruction to lifting a morphism of metric graphs $\bar{\varphi} : \Gamma' \rightarrow \Gamma$ to a morphism of curves is the same as the obstruction to enriching φ with the structure of a morphism of metrized complexes of curves. The following more precise statement is an immediate consequence of Proposition 7.15 and Theorem 3.24:

Corollary D. *Assume $\text{char}(k) = 0$. Let $\bar{\varphi} : \Gamma' \rightarrow \Gamma$ be a finite harmonic morphism of Λ -metric graphs. Suppose that φ can be enriched to a morphism of metrized complexes of curves. Then φ is liftable.*

We remark that we actually take advantage of the stronger form of Theorem B, in particular the calculation of the automorphism group of a lift, when lifting group actions on metrized complexes in [ABBR14].

The problem of enriching a morphism of metric graphs to a morphism of metrized complexes is essentially equivalent to the problem of assigning k -curves $C_p, C_{p'}$ to vertices $p \in \Gamma, p' \in \Gamma'$ and finding morphisms of k -curves $C_{p'} \rightarrow C_p$ with prescribed ramification profiles. The latter is a question about whether certain Hurwitz numbers are nonzero. See Proposition II.3.3. Allowing arbitrary choices of residue curves $C_p, C_{p'}$, this can always be done:

Theorem. *Suppose that $\text{char}(k) = 0$. Any finite harmonic morphism $\varphi : \Gamma' \rightarrow \Gamma$ of Λ -metric graphs is liftable to a morphism of curves.*

See Theorem II.3.11. Imposing requirements on the lifted curves X, X' give rise to variants of this tropical lifting problem. The most basic such requirement is to specify the genera of the curves. An

augmented metric graph is a metric graph Γ along with an integer $g(p) \in \mathbf{Z}_{\geq 0}$ for every finite vertex p of Γ , called the *genus*. The *genus* of an augmented graph Γ is defined to be

$$g(\Gamma) = h_1(\Gamma) + \sum_p g(p),$$

where $h_1(\Gamma)$ is the first Betti number. Any metrized complex of curves has an underlying augmented metric graph, where $g(p) := g(C_p)$, the genus of the residue curve. If Σ is a skeleton of a curve X , viewed as an augmented metric graph, then we have $g(\Sigma) = g(X)$.

A finite harmonic morphism of augmented metric graphs $\Gamma' \rightarrow \Gamma$ is *liftable* if the morphism of underlying metric graphs is liftable to a finite morphism of curves $X' \rightarrow X$ such that the isomorphism of Γ (resp. Γ') with a skeleton of X (resp. X') respects the augmented metric graph structure. In this case, $g(X) = g(\Gamma)$ and $g(X') = g(\Gamma')$. By our lifting theorem, $\Gamma' \rightarrow \Gamma$ is liftable to a morphism of curves if and only if it can be enhanced to a morphism of metrized complexes such that $g(C_p) = g(p)$ for every finite vertex. Now the obstruction to lifting is nontrivial: there exist finite harmonic morphisms of metric graphs which cannot be enhanced to a morphism of metrized complexes. This corresponds to the emptiness of certain Hurwitz spaces over k . See Section II.5 for the following striking example of this phenomenon:

Theorem. *There exists an augmented metric graph Γ admitting a finite morphism of degree $d > 0$ to an augmented metric graph T of genus zero, but which is not isomorphic to a skeleton of a d -gonal curve X .*

In [ABBR14] we also give applications of the lifting theorem to the following kinds of tropical lifting problems, among others:

- When T is an augmented metric graph of genus zero, we study a variant of the lifting problem in which the genus of the source curve is prescribed, but the degree of the morphism is not. We use this to prove that linear equivalence of divisors on a tropical curve C coincides with the equivalence relation generated by declaring that the fibers of every finite harmonic morphism from C to the tropical projective line are equivalent.
- We study liftability of metrized complexes equipped with a finite group action, and as an application classify all (augmented) metric graphs arising as the skeleton of a hyperelliptic curve. As mentioned above, this study actually takes advantage of the more precise form of Theorem B, not just the existence of a lift.

1.7. Organization of the paper. In Section 2 we recall some basic definitions of graphs with additional structures, including metrized complexes of curves, and define harmonic morphisms between them. We provide somewhat more detail than is strictly necessary for the exposition in this paper, but will prove necessary in [ABBR14].

In Section 3 we review the structure theory of analytic curves, and define the skeleton of a curve. We show that the skeleton is naturally a metrized complex of curves, and we prove that any metrized complex arises in this way. In Section 4 we develop “relative” versions of the results of Section 3. We prove that a finite morphism of curves induces a finite morphism between suitable choices of skeleta, and that the map on skeleta is a finite harmonic morphism of metrized complexes of curves. In Section 5 we show how the previous two sections can be used to derive various simultaneous semistable reduction theorems.

In Section 6 we prove a local lifting theorem, which essentially says that a tame covering of metrized complexes of curves can be lifted in a unique way to a tame covering of analytic K -curves in a neighborhood of a vertex. This is the technical heart of the lifting theorem. In Section 7 we globalize the considerations of the previous section, giving the classification of lifts of a tame covering of metrized complexes of curves to a tame covering of triangulated punctured K -curves.

Section 8 contains an example application, in which we construct a morphism of curves over a discretely valued field such that the covariantly associated morphism of component groups of Jacobians has a prescribed behavior. We use this to answer a question of Ribet.

1.8. Related work. Another framework for the foundations of tropical geometry has been proposed by Kontsevich-Soibelman [KS01, KS06] and Mikhalkin [Mik06, Mik05], in which tropical objects are associated to real one-parameter families of complex varieties. We refer to [KS01, KS06] for some conjectures relating this framework to Berkovich spaces. In this setting, the notion of metrized complex of curves is similar to the notion of phase-tropical curves, and Proposition 7.15 is a consequence of Riemann’s Existence Theorem. We refer the interested reader to the forthcoming paper [Mik] for more details (see also [BBM11] where this statement is implicit, as well as Corollary II.3.8).

Harmonic morphisms of finite graphs were introduced by Urakawa in [Ura00] and further explored in [BN09]. Harmonic morphisms of metric graphs have been introduced independently by several different people. Except for the integrality condition on the slopes, they appear already in Anand’s paper [Ana00]. The definition we use is the same as the one given in [Mik06, Cha12].

Finite harmonic morphisms of metrized complexes can be regarded as a metrized version of the notion of admissible cover due to Harris and Mumford [HM82] (where, in addition, arbitrary ramification above smooth points is allowed). Recall that for two semistable curves Y' and Y over a field k , a finite surjective morphism $\varphi : Y' \rightarrow Y$ is an *admissible cover* if $\varphi^{-1}(Y^{\text{sing}}) = Y'^{\text{sing}}$ and for each singular point y' of Y' , the ramification indices at y' along the two branches intersecting at y' coincide. (In addition, one usually imposes that all the other ramifications of φ are simple). An admissible cover naturally gives rise to a finite harmonic morphism of metrized complexes: denoting by \mathcal{C} the regularization of Y (the metrized complex associated to Y in which each edge has length one), define \mathcal{C}' as the metrized complex obtained from Y' by letting the length of the edge associated to the double point y' be $1/r_{y'}$ (where $r_{y'}$ is the ramification index of φ at y' along either of the two branches). The morphism φ of semistable curves naturally extends to a finite harmonic morphism $\varphi : \mathcal{C}' \rightarrow \mathcal{C}$: on each edge e' of \mathcal{C}' corresponding to a double point y' of Y' , the restriction of φ to e' is linear with slope (or “expansion factor”) $r_{y'}$. Conversely, a finite harmonic morphism of metrized complexes of curves gives rise to an admissible cover of semistable curves (without the supplementary condition on simple ramification) by forgetting the metrics on both sides, remembering only the expansion factor along each edge.

A related but somewhat different Berkovich-theoretic point of view on simultaneous semistable reduction for curves can be found in Welliaveetil’s recent preprint [Wel13]; harmonic morphisms of metrized complexes of curves play an implicit role in his paper.

2. METRIC GRAPHS AND METRIZED COMPLEXES OF CURVES

In this section we recall several definitions of graphs with some additional structures and morphisms between them. A number of these definitions are now standard in tropical geometry; we refer for example to [BN07], [Mik06], [BBM11], and [AB12]. We also provide a number of examples. Some of the definitions in this section are not necessary in the generality or the form in which they are presented for the purposes of this paper, but will be useful in [ABBR14].

Throughout this section, we fix Λ a non-trivial subgroup of $(\mathbf{R}, +)$.

2.1. Metric graphs. Given $r \in \mathbf{Z}_{\geq 1}$, we define $S_r \subset \mathbf{C}$ to be a “star with r branches”, i.e., a topological space homeomorphic to the convex hull in \mathbf{R}^2 of $(0, 0)$ and any r points, no two of which lie on a common line through the origin. We also define $S_0 = \{0\}$.

A *finite topological graph* Γ is the topological realization of a finite graph. That is to say, Γ is a compact 1-dimensional topological space such that for any point $p \in \Gamma$, there exists a neighborhood U_p of p in Γ homeomorphic to some S_r ; moreover there are only finitely many points p with U_p homeomorphic to S_r with $r \neq 2$.

The unique integer r such that U_p is homeomorphic to S_r is called the *valence* of p and denoted $\text{val}(p)$. A point of valence different from 2 is called an *essential vertex* of Γ . The set of *tangent directions* at p is $T_p(\Gamma) = \varinjlim_{U_p} \pi_0(U_p \setminus \{p\})$, where the limit is taken over all neighborhoods of p isomorphic to a star with r branches. The set $T_p(\Gamma)$ has $\text{val}(p)$ elements.

Definition 2.2. A *metric graph* is a finite connected topological graph Γ equipped with a complete inner metric on $\Gamma \setminus V_\infty(\Gamma)$, where $V_\infty(\Gamma) \subsetneq \Gamma$ is some (finite) set of 1-valent vertices of Γ called *infinite vertices* of Γ . (An *inner metric* is a metric for which the distance between two points x and y is the minimum of the lengths of all paths between x and y .)

Let Γ be a metric graph containing a finite essential vertex. We say that Γ is a Λ -*metric graph* if the distance between any two finite essential vertices of Γ lies in Λ . A Λ -*point* of Γ is a point of Γ whose distance to any finite essential vertex of Γ lies in Λ .

Let Γ be a metric graph such that all essential vertices are infinite, so Γ is homeomorphic to a circle or a completed line. We say that Γ is a Λ -*metric graph* if it is a completed line or a circle whose circumference is in Γ . Choose a point $v \in \Gamma$ of valency equal to 2. A Λ -*point* of Γ is a point of Γ whose distance to v lies in Λ .

One can equip Γ with a degenerate metric in which the infinite vertices are at infinite distance from any other point of Γ . When no confusion is possible about the subgroup Λ , we will sometimes write simply *metric graph* instead of Λ -*metric graph*.

Definition 2.3. Let Γ be a Λ -metric graph. A *vertex set* $V(\Gamma)$ is a finite subset of the Λ -points of Γ containing all essential vertices. An element of a fixed vertex set $V(\Gamma)$ is called a *vertex* of Γ , and the closure of a connected component of $\Gamma \setminus V(\Gamma)$ is called an *edge* of Γ . An edge which is homeomorphic to a circle is called a *loop edge*. An edge adjacent to an infinite vertex is called an *infinite edge*. We denote by $V_f(\Gamma)$ the set of finite vertices of Γ , and by $E_f(\Gamma)$ the set of finite edges of Γ .

Fix a vertex set $V(\Gamma)$. We denote by $E(\Gamma)$ the set of edges of Γ . Since Γ is a metric graph, we can associate to each edge e of Γ its length $\ell(e) \in \Lambda \cup \{+\infty\}$. Since the metric on $\Gamma \setminus V_\infty(\Gamma)$ is complete, an edge e is infinite if and only if $\ell(e) = +\infty$. The notion of vertices and edges of Γ depends, of course, on the choice of a vertex set; we will fix such a choice without comment whenever there is no danger of confusion.

Definition 2.4. Fix vertex sets $V(\Gamma')$ and $V(\Gamma)$ for two Λ -metric graphs Γ' and Γ , respectively, and let $\varphi : \Gamma' \rightarrow \Gamma$ be a continuous map.

- The map φ is called a $(V(\Gamma'), V(\Gamma))$ -*morphism of Λ -metric graphs* if we have $\varphi(V(\Gamma')) \subset V(\Gamma)$, $\varphi^{-1}(E(\Gamma)) \subset E(\Gamma')$, and the restriction of φ to any edge e' of Γ' is a dilation by some factor $d_{e'}(\varphi) \in \mathbf{Z}_{\geq 0}$.
- The map φ is called a *morphism of Λ -metric graphs* if there exists a vertex set $V(\Gamma')$ of Γ' and a vertex set $V(\Gamma)$ of Γ such that φ is a $(V(\Gamma'), V(\Gamma))$ -morphism of Λ -metric graphs.
- The map φ is said to be *finite* if $d_{e'}(\varphi) > 0$ for any edge $e' \in E(\Gamma')$.

An edge e' of Γ' is mapped to a vertex of Γ if and only if $d_{e'}(\varphi) = 0$. Such an edge is said to be *contracted* by φ . A morphism $\varphi : \Gamma' \rightarrow \Gamma$ is finite if and only if there are no contracted edges, which holds if and only if $\varphi^{-1}(p)$ is a finite set for any point $p \in \Gamma$. If φ is finite, then $p' \in V_f(\Gamma')$ if and only if $\varphi(p') \in V_f(\Gamma)$.

The integer $d_{e'}(\varphi) \in \mathbf{Z}_{\geq 0}$ in Definition 2.4 is called the *degree* of φ along e' (it is also sometimes called the *weight* of e' or *expansion factor* along e' in the literature). Since $\ell(\varphi(e')) = d_{e'}(\varphi) \cdot \ell(e')$, it follows in particular that if $d_{e'}(\varphi) \geq 1$ then e' is infinite if and only if $\varphi(e')$ is infinite. Let $p' \in V(\Gamma')$, let $v' \in T_{p'}(\Gamma')$, and let $e' \in E(\Gamma')$ be the edge in the direction of v' . We define the *directional derivative* of φ in the direction v' to be $d_{v'}(\varphi) := d_{e'}(\varphi)$. If we set $p = \varphi(p')$, then φ induces a map

$$d\varphi(p') : \{v' \in T_{p'}(\Gamma') : d_{v'}(\varphi) \neq 0\} \longrightarrow T_p(\Gamma)$$

in the obvious way.

Example 2.5. Figure 1 depicts four examples of morphisms of metric graphs $\varphi : \Gamma' \rightarrow \Gamma$. We use the following conventions in our pictures: black dots represent vertices of Γ' and Γ ; we label an edge with its degree if and only if the degree is different from 0 and 1; we do not specify the lengths of edges of Γ' and Γ .

The morphisms depicted in Figure 1(a), (b), and (c) are finite, while the one depicted in Figure 1(d) is not.




FIGURE 1

Definition 2.6. Let $\varphi : \Gamma' \rightarrow \Gamma$ be a morphism of Λ -metric graphs, let $p' \in \Gamma'$, and let $p = \varphi(p')$. The morphism φ is *harmonic at p'* provided that, for every tangent direction $v \in T_p(\Gamma)$, the number

$$d_{p'}(\varphi) := \sum_{\substack{v' \in T_{p'}(\Gamma') \\ v' \mapsto v}} d_{v'}(\varphi)$$

is independent of v . The number $d_{p'}(\varphi)$ is called the *degree of φ at p'* .

We say that φ is *harmonic* if it is *surjective* and harmonic at all $p' \in \Gamma'$; in this case the number $\deg(\varphi) = \sum_{p' \mapsto p} d_{p'}(\varphi)$ is independent of $p \in \Gamma$, and is called the *degree of φ* .

Remark 2.7. In the above situation, if Γ consists of a single vertex v and $\varphi : \Gamma' \rightarrow \Gamma$ is a morphism of metric graphs, then φ is by definition harmonic. In this case the quantity $d_{p'}(\varphi)$ is not defined, so we include the choice of a positive integer $d_{p'}(\varphi)$ for each $p' \in V_f(\Gamma')$ as part of the data of the morphism φ . Note that if φ is finite and $\Gamma = \{p\}$, then $\Gamma' = \{p'\}$.

If both Γ' and Γ have at least one edge, then a non-constant morphism which is harmonic at all $p' \in V(\Gamma')$ is automatically surjective.

Example 2.8. The morphism depicted in Figure 1(a) is not harmonic, while the ones depicted in Figure 1(b), (c), and (d) are (for suitable choices of lengths).

2.9. Harmonic morphisms and harmonic functions. Given a metric graph Γ and a non-empty open set U in Γ , a function $f : U \rightarrow \mathbf{R}$ is said to be *harmonic on U* if f is piecewise affine with integer slopes and for all $x \in U$, the sum of the slopes of f along all outgoing tangent directions at x is equal to 0.

Since we will not use it elsewhere in the paper, we omit the proof of the following result (which is very similar to the proof of [BN09, Proposition 2.6]):

Proposition 2.10. *A morphism $\varphi : \Gamma' \rightarrow \Gamma$ of metric graphs is harmonic if and only if for every open set $U \subseteq \Gamma$ and every harmonic function $f : U \rightarrow \mathbf{R}$, the pullback function $\varphi^* f : \varphi^{-1}(U) \rightarrow \mathbf{R}$ is also harmonic.*

Equivalently, a morphism of metric graphs is harmonic if and only if germs of harmonic functions pull back to germs of harmonic functions.

2.11. For a metric graph Γ , we let $\text{Div}(\Gamma)$ denote the free abelian group on Γ . Given a harmonic morphism $\varphi : \Gamma' \rightarrow \Gamma$ of metric graphs, there are natural pull-back and push-forward homomorphisms $\varphi^* : \text{Div}(\Gamma) \rightarrow \text{Div}(\Gamma')$ and $\varphi_* : \text{Div}(\Gamma') \rightarrow \text{Div}(\Gamma)$ defined by

$$\varphi^*(p) = \sum_{p' \mapsto p} d_{p'}(\varphi) (p') \quad \text{and} \quad \varphi_*(p') = (\varphi(p'))$$

and extending linearly. It is clear that for $D \in \text{Div}(\Gamma)$ we have $\deg(\varphi^*(D)) = \deg(\varphi) \cdot \deg(D)$ and $\deg(\varphi_*(D)) = \deg(D)$.

2.12. Augmented metric graphs. Here we consider Λ -metric graphs together with the data of a non-negative integer at each point. For a finite point p , this integer should be thought as the genus of a “virtual algebraic curve” lying above p .

Definition 2.13. An *augmented Λ -metric graph* is a Λ -metric graph Γ along with a function $g : \Gamma \rightarrow \mathbf{Z}_{\geq 0}$, called the *genus function*, such that $g(p) = 0$ for all points of Γ except possibly for finitely many (finite) Λ -points $p \in \Gamma$. The *essential vertices* of Γ are the points p for which $\text{val}(p) \neq 2$ or $g(p) > 0$. A *vertex set* of Γ is a vertex set $V(\Gamma)$ of the underlying metric graph which contains all essential vertices of Γ as an augmented graph.

An augmented metric graph is said to be *totally degenerate* if the genus function is identically zero.

The *genus* of Γ is defined to be

$$g(\Gamma) = h_1(\Gamma) + \sum_{p \in \Gamma} g(p),$$

where $h_1(\Gamma) = \dim_{\mathbf{Q}} H_1(\Gamma, \mathbf{Q})$ is the first Betti number of Γ . The *canonical divisor* on Γ is

$$(2.13.1) \quad K_{\Gamma} = \sum_{p \in \Gamma} (\text{val}(p) + 2g(p) - 2)(p).$$

A *harmonic morphism of augmented Λ -metric graphs* $\varphi : \Gamma' \rightarrow \Gamma$ is a map which is a harmonic morphism between the underlying metric graphs of Γ' and Γ .

Note that both summations in the above definition are in fact over essential vertices of Γ . The degree of the canonical divisor of an augmented Λ -metric graph Γ is $\deg(K_{\Gamma}) = 2g(\Gamma) - 2$. An augmented metric graph of genus 0 will also be called a *rational augmented metric graph*.

2.14. Let $\varphi : \Gamma' \rightarrow \Gamma$ be a harmonic morphism of augmented Λ -metric graphs. The *ramification divisor* of φ is the divisor $R = \sum R_{p'}(p')$, where

$$(2.14.1) \quad R_{p'} = d_{p'}(\varphi) \cdot (2 - 2g(\varphi(p'))) - (2 - 2g(p')) - \sum_{v' \in T_{p'}(\Gamma')} (d_{v'}(\varphi) - 1).$$

Note that if $p' \in V_{\infty}(\Gamma')$, then $R_{p'} \geq 0$ if $d_{p'}(\varphi) > 0$, and $R_{p'} = -1$ if $d_{p'}(\varphi) = 0$. The definition of $R_{p'}$ in (2.14.1) is rigged so that we have the following graph-theoretic analogue of the *Riemann–Hurwitz formula*:

$$(2.14.2) \quad K_{\Gamma'} = \varphi^*(K_{\Gamma}) + R.$$

In particular, the sum $R = \sum R_{p'}(p')$ is in fact finite.

Definition 2.15. Let $\varphi : \Gamma' \rightarrow \Gamma$ be a finite harmonic morphism of augmented Λ -metric graphs. We say that φ is *étale* at a point $p' \in \Gamma'$ provided that $R_{p'} = 0$. We say that φ is *generically étale* if R is supported on the set of infinite vertices of Γ' and that φ is *étale* if $R = 0$.

2.16. Metrized complexes of curves. Recall that k is an algebraically closed field. A *metrized complex of curves* is, roughly speaking, an augmented metric graph Γ together with a chosen vertex set $V(\Gamma)$ and a marked algebraic k -curve of genus $g(p)$ for each finite vertex $p \in V(\Gamma)$. More precisely:

Definition 2.17. A *Λ -metrized complex of k -curves* consists of the following data:

- An augmented Λ -metric graph Γ equipped with the choice of a distinguished vertex set $V(\Gamma)$.
- For every finite vertex p of Γ , a smooth, proper, connected k -curve C_p of genus $g(p)$.
- An injective function $\text{red}_p : T_p(\Gamma) \rightarrow C_p(k)$, called the *reduction map*. We call the image of red_p the set of *marked points* on C_p .

Example 2.18. Figure 2 depicts a particular metrized complex of curves over \mathbf{C} . At each finite vertex p there is an associated compact Riemann surface C_p , together with a finite set of marked points (in red).

Definition 2.19. A *harmonic morphism of metrized complexes of curves* consists of a harmonic $(V(\Gamma'), V(\Gamma))$ -morphism $\varphi : \Gamma' \rightarrow \Gamma$ of augmented metric graphs, and for every finite vertex p' of Γ'




FIGURE 2

with $d_{p'}(\varphi) > 0$ a finite morphism of curves $\varphi_{p'} : C'_{p'} \rightarrow C_{\varphi(p')}$, satisfying the following compatibility conditions:

- (1) For every finite vertex $p' \in V(\Gamma')$ and every tangent direction $v' \in T_{p'}(\Gamma')$ with $d_{v'}(\varphi) > 0$, we have $\varphi_{p'}(\text{red}_{p'}(v')) = \text{red}_{\varphi(p')}(d\varphi_{p'}(v'))$, and the ramification degree of $\varphi_{p'}$ at $\text{red}_{p'}(v')$ is equal to $d_{v'}(\varphi)$.
- (2) For every finite vertex $p' \in V(\Gamma')$ with $d_{p'}(\varphi) > 0$, every tangent direction $v \in T_{\varphi(p')}(\Gamma)$, and every point $x' \in \varphi_{p'}^{-1}(\text{red}_{\varphi(p')}(v)) \subset C'_{p'}(k)$, there exists $v' \in T_{p'}(\Gamma')$ such that $\text{red}_{p'}(v') = x'$.
- (3) For every finite vertex $p' \in V(\Gamma')$ with $d_{p'}(\varphi) > 0$ we have $d_{p'}(\varphi) = \deg(\varphi_{p'})$.

A harmonic morphism of metrized complexes of curves is called *finite* if the underlying harmonic morphism of (augmented) metric graphs is finite. Metrized complexes of curves with harmonic morphisms between them form a category, with finite harmonic morphisms giving rise to a subcategory.

If Γ has at least one edge, then (3) follows from (1) and (2), since the sum of the ramification degrees of a finite morphism of curves along any fiber is equal to the degree of the morphism.

Let $\varphi : C' \rightarrow C$ be a finite morphism of smooth, proper, connected k -curves. Recall that φ is *tamely ramified* if either $\text{char}(k) = 0$ or the ramification degree of φ at every closed point of C' is prime to the characteristic of k .

Remark 2.20.

- (1) Let $\varphi : C' \rightarrow C$ be a finite morphism. Let D be the (finite) set of branch points, let $D' = \varphi^{-1}(D) \subset C'(k)$, and let $U = C \setminus D$ and $U' = C' \setminus D'$. Then φ is tamely ramified if and only if $\varphi|_{U'} : U' \rightarrow U$ is a tamely ramified cover of U over C relative to D in the sense of [SGA1, XIII.2.1.3]: see the discussion in §2.0 of *loc. cit.*
- (2) A tamely ramified morphism is separable (i.e. generically étale).
- (3) The Riemann–Hurwitz formula applies to a tamely ramified morphism.

Definition 2.21. Let $\varphi : C' \rightarrow C$ be a finite harmonic morphism of metrized complexes of curves. We say that φ is a *tame harmonic morphism* if either $\text{char}(k) = 0$, or $\text{char}(k) = p > 0$ and $\varphi_{p'}$ is tamely ramified for all finite vertices $p' \in V(\Gamma')$. We call φ a *tame covering* if in addition $\varphi : \Gamma' \rightarrow \Gamma$ is a generically étale finite morphism of augmented metric graphs.

Let $\varphi : C' \rightarrow C$ be a tame harmonic morphism. Then each $\varphi_{p'} : C'_{p'} \rightarrow C_{\varphi(p')}$ is a separable morphism. Note that if $\text{char}(k) = p$ then $d_{e'}(\varphi)$ is prime to p for all edges e' of Γ' . We have the following kind of converse statement:

Proposition 2.22. Let $\varphi : C' \rightarrow C$ be a finite harmonic morphism of metrized complexes of curves. Suppose that φ is generically étale and that $\varphi_{p'} : C'_{p'} \rightarrow C_{\varphi(p')}$ is a separable morphism of curves for all finite vertices p' of Γ' . Let p' be a finite vertex of Γ' , and suppose that $\varphi_{p'}$ is ramified at a point $x' \in C'_{p'}(k)$. Then there exists $v' \in T_{p'}(\Gamma')$ such that $\text{red}_{p'}(v') = x'$. In particular, if $\text{char}(k) = p > 0$ and $d_{e'}(\varphi)$ is prime to p for all edges e' of C' , then $\varphi_{p'}$ is tamely ramified and φ is a tame covering.

Proof. Let $p = \varphi(p')$. Since φ is generically étale we have

$$0 = R_{p'} = d_{p'}(\varphi)(2 - 2g(p)) - (2 - 2g(p')) - \sum_{v' \in T_{p'}(\Gamma')} (d_{v'}(\varphi) - 1).$$

Since $g(p) = g(C_p)$, $g(p') = g(C'_{p'})$, and $d_{p'}(\varphi)$ is the degree of $\varphi_{p'} : C'_{p'} \rightarrow C_p$, this gives

$$\sum_{v' \in T_{p'}(\Gamma')} (d_{v'}(\varphi) - 1) = \deg(\varphi_{p'}) (2 - 2g(C_p)) - (2 - 2g(C'_{p'})).$$

The Riemann–Hurwitz formula as applied to $\varphi_{p'}$ says that the right-hand side of the above equation is equal to the degree of the ramification divisor of $\varphi_{p'}$ (this holds whenever $\varphi_{p'}$ is separable), so the proposition follows because for all $v' \in T_{p'}(\Gamma')$, the ramification degree of $\varphi_{p'}$ at $\text{red}_{p'}(v')$ is $d_{v'}(\varphi)$. ■

Remark 2.23. Loosely speaking, Proposition 2.22 says that all ramification of all morphisms $\varphi_{p'}$ is “visible” in the edge degrees of the underlying morphism of metric graphs. It follows from Proposition 2.22 that if $\Gamma' = \{p'\}$, then $\varphi_{p'}$ is étale.

3. ANALYTIC CURVES AND THEIR SKELETA

In this section we recall the definition and basic properties of the skeleton of the analytification of an algebraic curve as outlined in [Tem] and elaborated in [BPR11, §5]. These foundational results are well-known to experts, and appear in various guises in the literature. See for example Berkovich [Ber90, Ber99], Thuillier [Thu05], and Ducros [Duc08]. In this section we also show that the skeleton is naturally a metrized complex of curves, and prove that any metrized complex arises as the skeleton of a curve. Finally, we introduce the important notion of a triangulated punctured curve, which is essentially a punctured curve along with the data of a skeleton with a distinguished set of vertices.

Recall that K is an algebraically closed field which is complete with respect to a nontrivial non-Archimedean valuation $\text{val} : K \rightarrow \mathbf{R} \cup \{\infty\}$. Its valuation ring is R , its maximal ideal is \mathfrak{m}_R , its residue field is $k = R/\mathfrak{m}_R$, and its value group is $\Lambda = \text{val}(K^\times) \subset \mathbf{R}$.

By an *analytic curve* we mean a strictly K -analytic space X of pure dimension 1. For our purposes the most important example of an analytic curve is the analytification of a smooth, connected, projective algebraic K -curve. If X is an analytic curve, then we define $\mathbf{H}(X) = X \setminus X(K)$ as in [BPR11, 3.10].

3.1. Analytic building blocks. We identify the set underlying the analytification of the affine line with the set of multiplicative seminorms $\|\cdot\| : K[t] \rightarrow \mathbf{R}_{\geq 0}$ which extend the absolute value on K . We let $\text{val} : \mathbf{A}^{1,\text{an}} \rightarrow \mathbf{R} \cup \{\infty\}$ denote the valuation map

$$\text{val}(x) = -\log |x| = -\log \|t\|_x.$$

For $a \in K^\times$ the *standard closed ball of radius $|a|$* is the affinoid domain

$$\mathbf{B}(a) = \text{val}^{-1}([\text{val}(a), \infty]) \subset \mathbf{A}^{1,\text{an}}$$

and the *standard open ball of radius $|a|$* is the open analytic domain

$$\mathbf{B}(a)_+ = \text{val}^{-1}((\text{val}(a), \infty]) \subset \mathbf{A}^{1,\text{an}}.$$

Note that scaling by a gives isomorphisms $\mathbf{B}(1) \xrightarrow{\sim} \mathbf{B}(a)$ and $\mathbf{B}(1)_+ \xrightarrow{\sim} \mathbf{B}(a)_+$. We call $\mathbf{B}(1)$ (resp. $\mathbf{B}(1)_+$) the *standard closed* (resp. *open*) *unit ball*. For $a \in K^\times$ with $\text{val}(a) \geq 0$ the *standard closed annulus of modulus $\text{val}(a)$* is the affinoid domain

$$\mathbf{S}(a) = \text{val}^{-1}([0, \text{val}(a)]) \subset \mathbf{A}^{1,\text{an}}$$

and if $\text{val}(a) > 0$ the *standard open annulus of modulus $\text{val}(a)$* is the open analytic domain

$$\mathbf{S}(a)_+ = \text{val}^{-1}((0, \text{val}(a))) \subset \mathbf{A}^{1,\text{an}}.$$

The *standard punctured open ball* is the open analytic domain

$$\mathbf{S}(0)_+ = \text{val}^{-1}((0, \infty)) = \mathbf{B}(1)_+ \setminus \{0\} \subset \mathbf{A}^{1,\text{an}}.$$

By a *closed unit ball* (resp. *open unit ball*, resp. *closed annulus*, resp. *open annulus*, resp. *punctured open ball*) we will mean a K -analytic space which is isomorphic to the standard closed unit ball

(resp. the standard open unit ball, resp. the standard closed annulus of modulus $\text{val}(a)$ for some $a \in R \setminus \{0\}$, resp. the standard open annulus of modulus $\text{val}(a)$ for some $a \in \mathfrak{m}_R \setminus \{0\}$, resp. the standard punctured open ball). A *generalized open annulus* is a K -analytic space which is either an open annulus or a punctured open ball. It is a standard fact that the modulus of a closed (resp. open) annulus is an isomorphism invariant; see for instance [BPR11, Corollary 5.6]. We define the modulus of a punctured open ball to be ∞ .

Let $a \in \mathfrak{m}_R$ (so $a = 0$ is allowed). There is a natural section $\sigma : (0, \text{val}(a)) \rightarrow \mathbf{S}(a)_+$ of the valuation map $\text{val} : \mathbf{S}(a)_+ \rightarrow (0, \text{val}(a))$ sending r to the maximal point of the affinoid domain $\text{val}^{-1}(r)$ if $r \in \Lambda$, resp. the only point of $\text{val}^{-1}(r)$ if $r \notin \Lambda$. The *skeleton* of $\mathbf{S}(a)_+$ is defined to be the image of σ and is denoted $\Sigma(\mathbf{S}(a)_+) = \sigma((0, \text{val}(a)))$. We will always identify the skeleton of the standard generalized open annulus $\mathbf{S}(a)_+$ with the interval or ray $(0, \text{val}(a))$. It follows from [BPR11, Proposition 5.5] that the skeleton of a standard open annulus or standard punctured open ball is an isomorphism invariant, so if A is a generalized open annulus, then we can define the *skeleton* $\Sigma(A)$ of A to be the image of the skeleton of a standard open annulus or standard punctured open ball $\mathbf{S}(a)_+$ under any isomorphism $\mathbf{S}(a)_+ \xrightarrow{\sim} A$.

Let T be a metric space and let $x, y \in T$. A *geodesic segment from x to y* is the image of a locally isometric embedding $[a, b] \hookrightarrow T$ with $a \mapsto x$ and $b \mapsto y$. We often identify a geodesic segment with its image in T . Recall that an **R-tree** is a metric space T with the following properties:

- (i) For all $x, y \in T$ there is a unique geodesic segment $[x, y]$ from x to y .
- (ii) For all $x, y, z \in T$, if $[x, y] \cap [y, z] = \{y\}$, then $[x, z] = [x, y] \cup [y, z]$.

See [BR10, Appendix B]. It is proved in §1.4 of loc. cit. that $\mathbf{H}(\mathbf{B}(1))$ is naturally an **R-tree**. Since any path-connected subspace of an **R-tree** is an **R-tree** as well, if X is a standard open annulus or standard (punctured) open ball then $\mathbf{H}(X)$ is an **R-tree**. It is proved in [BPR11, Corollary 5.61] that the metric structure on $\mathbf{H}(X)$ is an isomorphism invariant, so the same statement applies to open balls and generalized open annuli. For $a \in \mathfrak{m}_R$ the section $\sigma : (0, \text{val}(a)) \rightarrow \mathbf{S}(a)_+$ is an isometry.

It also follows from the results in [BR10, §1.4] that for any $x \in \mathbf{H}(\mathbf{B}(1))$ and any type-1 point $y \in \mathbf{B}(1)$, there is a unique continuous injection $\alpha : [0, \infty) \hookrightarrow \mathbf{B}(1)$ with $\alpha(0) = x$ and $\alpha(\infty) = y$, $\alpha([0, \infty)) \subset \mathbf{H}(\mathbf{B}(1))$, and such that α is an isometry when restricted to $[0, \infty)$. We let $[x, y]$ denote the image of α and we call $[x, y]$ the *geodesic segment from x to y* . Similarly, if x and y are both type-1 points then there is a unique continuous injection $\alpha : [-\infty, \infty] \rightarrow \mathbf{B}(1)$ with $\alpha(-\infty) = x$ and $\alpha(\infty) = y$, $\alpha(\mathbf{R}) \subset \mathbf{H}(\mathbf{B}(1))$, and such that $\alpha|_{\mathbf{R}}$ is an isometry; the image of α is called the *geodesic segment from x to y* and is denoted $[x, y]$. Restricting to a suitable analytic subdomain of $\mathbf{B}(1)$ allows us to define geodesic segments between any two points of an open ball or generalized open annulus.

3.2. Open balls. The closure of $\mathbf{B}(1)_+$ in $\mathbf{B}(1)$ consists of $\mathbf{B}(1)_+$ and a single type-2 point x , called the *end* of $\mathbf{B}(1)_+$. The end is the Shilov boundary point of $\mathbf{B}(1)$: see for instance the proof of [BPR11, Lemma 5.16]. The valuation map $\text{val} : \mathbf{B}(1)_+ \rightarrow (0, \infty]$ extends to a continuous map $\text{val} : \mathbf{B}(1)_+ \cup \{x\} \rightarrow [0, \infty]$. The set $\mathbf{B}(1)_+ \cup \{x\}$ is path-connected and compact, being a closed subspace of the compact space $\mathbf{B}(1)$. (In fact $\mathbf{B}(1)_+ \cup \{x\}$ is the one-point compactification of $\mathbf{B}(1)_+$.) For any $y \in \mathbf{B}(1)_+$ the geodesic segment $[x, y] \subset \mathbf{B}(1)$ is contained in $\mathbf{B}(1)_+ \cup \{x\}$.

Lemma 3.3. *Let $X = \text{Spec}(A)$ be an irreducible affine K -curve and let $\varphi : \mathbf{B}(1)_+ \rightarrow X^{\text{an}}$ be a morphism with finite fibers. Then φ extends in a unique way to a continuous map $\mathbf{B}(1)_+ \cup \{x\} \rightarrow X^{\text{an}}$, and the image of x is a type-2 point of X^{an} .*

Proof. Let $f \in A$ be nonzero and define $F : (0, \infty) \rightarrow \mathbf{R}$ by $F(r) = -\log |f \circ \varphi(\sigma(r))|$, where $\sigma : (0, \infty) \rightarrow \mathbf{S}(0)_+ \subset \mathbf{B}(1)_+$ is the canonical section of val . By [BPR11, Proposition 5.10], F is a piecewise affine function which is differentiable away from finitely many points $r \in \Lambda \cap (0, \infty)$, and for any point $r \in (0, \infty)$ at which F is differentiable, the derivative of F is equal to the number of zeros y of $f \circ \varphi$ with $\text{val}(y) > r$. It follows that $F(\Lambda \cap (0, \infty)) \subset \Lambda$. Since $f \circ \varphi$ has finitely many zeros, the limit $\lim_{r \rightarrow 0} F(r)$ exists and is contained in Λ . We define $\|f\|_0 = \exp(-\lim_{r \rightarrow 0} F(r))$. It is easy to see that $f \mapsto \|f\|_0$ is a multiplicative seminorm on A extending the absolute value on K , so $\|\cdot\|_0$ is

a point in X^{an} . Define $\varphi(x) = \|\cdot\|_0$. One shows as in the proof of [BPR11, Lemma 5.16] that φ is continuous on $\mathbf{B}(1)_+ \cup \{x\}$.

It remains to show that $\varphi(x) = \|\cdot\|_0$ is a type-2 point of X^{an} . Let $f \in A$ be a non-constant function that has a zero on $\varphi(\mathbf{B}(1)_+)$. Since $-\log \|f\|_0 \in \Lambda$, there exists $\alpha \in K^\times$ such that $\|\alpha f\|_0 = 1$; replacing f by αf , we may and do assume that $\|f\|_0 = 1$. Since $\varphi \circ f$ has a zero, by the above we have that $F = -\log |f \circ \varphi \circ \sigma|$ is monotonically increasing, so $F(r) > 0$ for all $r \in (0, \infty)$. For any $r \in (0, \infty)$, the maximal point of $\text{val}^{-1}(r) \subset \mathbf{B}(1)_+$ is equal to $\sigma(r)$, so for any $y \in \mathbf{B}(1)_+$ such that $\text{val}(y) = r$ we have $F(r) \leq -\log |f \circ \varphi(y)|$. It follows that $|f \circ \varphi(y)| < 1$ for all $y \in \mathbf{B}(1)_+$, so $\varphi(f(\mathbf{B}(1)_+)) \subset \mathbf{B}(1)_+$. Since $\mathbf{B}(1)_+$ is dense in $\mathbf{B}(1)_+ \cup \{x\}$, the image of x under $\varphi \circ f$ is contained in the closure $\mathbf{B}(1)_+ \cup \{x\}$ of $\mathbf{B}(1)_+$ in $\mathbf{B}(1)$. Since $1 = \|f\|_0 = |f \circ \varphi(x)|$, the point $f \circ \varphi(x) \notin \mathbf{B}(1)_+$, so $f \circ \varphi(x) = x$, which is a type-2 point of $\mathbf{A}^{1,\text{an}}$. Therefore $\varphi(x)$ is a type-2 point of X^{an} . \blacksquare

Applying Lemma 3.3 to a morphism $\varphi : \mathbf{B}(1)_+ \rightarrow \mathbf{B}(1)_+ \subset \mathbf{A}^{1,\text{an}}$, we see that any automorphism of $\mathbf{B}(1)_+$ extends (uniquely) to a homeomorphism $\mathbf{B}(1)_+ \cup \{x\} \rightarrow \mathbf{B}(1)_+ \cup \{x\}$, so it makes sense to speak of the *end* of any open ball. If B is an open ball with end x , we let \overline{B} denote $B \cup \{x\}$.

Let B be an open ball with end x . We define a partial ordering on \overline{B} by declaring $y \leq z$ if $y \in [x, z]$; again see [BR10, §1.4]. Equivalently, for $y, z \in B$ we have $y \leq z$ if and only if $|f(y)| \geq |f(z)|$ for all analytic functions f on B . The following lemma is proved as in [BPR11, Proposition 5.27].

Lemma 3.4. *Let B be an open ball and let $y \in B$ be a type-2 point. Then $B \setminus \{y\}$ is a disjoint union of the open annulus $A = \{z \in B : z \not\geq y\}$ with infinitely many open balls, and $B_y = \{z \in B : z \geq y\}$ is an affinoid subdomain of B isomorphic to the closed ball $\mathbf{B}(1)$.*

3.5. Open annuli. Let $a \in \mathfrak{m}_R \setminus \{0\}$. The closure of $\mathbf{S}(a)_+$ in $\mathbf{B}(1)$ consists of $\mathbf{S}(a)_+$ and the two type-2 points $x = \sigma(0)$ and $y = \sigma(\text{val}(a))$, called the *ends* of $\mathbf{S}(a)_+$: again see the proof of [BPR11, Lemma 5.16]. The valuation map $\text{val} : \mathbf{S}(a)_+ \rightarrow (0, \text{val}(a))$ extends to a continuous map $\text{val} : \mathbf{S}(a)_+ \cup \{x, y\} \rightarrow [0, \text{val}(a)]$, and for any $z \in \mathbf{S}(a)_+$ the geodesic segments $[x, z]$ and $[y, z]$ are contained in $\mathbf{S}(a)_+ \cup \{x, y\}$. The following lemma is proved in the same way as the first part of Lemma 3.3.

Lemma 3.6. *Let $X = \text{Spec}(A)$ be an irreducible affine K -curve, let $a \in \mathfrak{m}_R \setminus \{0\}$, and let $\varphi : \mathbf{S}(a)_+ \rightarrow X^{\text{an}}$ be a morphism with finite fibers. Let $x = \sigma(0)$ and $y = \sigma(\text{val}(a))$ be the ends of $\mathbf{S}(a)_+$. Then φ extends in a unique way to a continuous map $\mathbf{S}(a)_+ \cup \{x, y\} \rightarrow X^{\text{an}}$.*

It follows from Lemma 3.6 that any automorphism of $\mathbf{S}(a)_+$ extends to a homeomorphism $\mathbf{S}(a)_+ \cup \{x, y\} \xrightarrow{\sim} \mathbf{S}(a)_+ \cup \{x, y\}$, so it makes sense to speak of the *ends* of any open annulus. If A is an open annulus with ends x, y , then we let \overline{A} denote the compact space $A \cup \{x, y\}$.

The closure of the punctured open ball $\mathbf{S}(0)_+$ in $\mathbf{B}(1)$ is equal to $\mathbf{S}(0)_+ \cup \{0, x\}$, where x is the end of $\mathbf{B}(1)_+$. We define x to be the *end* of $\mathbf{S}(0)_+$ and 0 to be the *puncture*. As above, the end and puncture of $\mathbf{S}(0)_+$ are isomorphism invariants, so it makes sense to speak of the *end* and the *puncture* of any punctured open ball. If A is a punctured open ball with end x and puncture y , then we let \overline{A} denote the compact space $A \cup \{x, y\}$.

Let A be a generalized open annulus. Each connected component of $A \setminus \Sigma(A)$ is an open ball [BPR11, Lemma 5.12], and if B is such a connected component with end x , then the inclusion $B \hookrightarrow A$ extends to an inclusion $\overline{B} \hookrightarrow A$ with x mapping into $\Sigma(A)$; the image of \overline{B} is the closure of B in A . We define the retraction to the skeleton $\tau : A \rightarrow \Sigma(A)$ by fixing $\Sigma(A)$ and sending each connected component of $A \setminus \Sigma(A)$ to its end. If $A = \mathbf{S}(a)_+$, then the retraction τ coincides with $\sigma \circ \text{val} : \mathbf{S}(a)_+ \rightarrow \Sigma(\mathbf{S}(a)_+)$; in particular, if $(r, s) \subset (0, \text{val}(a)) = \mathbf{S}(a)_+$, then $\tau^{-1}((r, s)) = \text{val}^{-1}((r, s))$.

3.7. The skeleton of a curve. Let X be a smooth, connected, proper algebraic K -curve and let $D \subset X(K)$ be a finite set of closed points. The set $\mathbf{H}(X^{\text{an}})$ is naturally a metric space [BPR11, Corollary 5.62], although the metric topology on $\mathbf{H}(X^{\text{an}})$ is much finer than the topology induced by the topology on the K -analytic space X^{an} . The metric on $\mathbf{H}(X^{\text{an}})$ is locally modeled on an \mathbf{R} -tree [BPR11, Proposition 5.63].

Definition 3.8. A *semistable vertex set* of X is a finite set V of type-2 points of X^{an} such that $X^{\text{an}} \setminus V$ is a disjoint union of open balls and finitely many open annuli. A *semistable vertex set of (X, D)* is a semistable vertex set of X such that the points of D are contained in distinct open ball connected components of $X^{\text{an}} \setminus V$.

It is a consequence of the semistable reduction theorem of Deligne-Mumford that there exist semistable vertex sets of (X, D) : see [BPR11, Theorem 5.49]. In the sequel it will be convenient to consider a curve along with a choice of semistable vertex set, so we give such an object a name.

Definition 3.9. A *triangulated punctured curve* $(X, V \cup D)$ is a smooth, connected, proper algebraic K -curve X equipped with a finite set $D \subset X(K)$ of *punctures* and a semistable vertex set V of (X, D) .

Remark 3.10. This terminology is loosely based on that used in [Duc08] as well as the forthcoming book of Ducros on analytic spaces and analytic curves. Strictly speaking, what we have defined should be called a *semistably triangulated punctured curve*, but as these are the only triangulations that we consider, we will not need the added precision.

3.11. Let $(X, V \cup D)$ be a triangulated punctured curve, so

$$(3.11.1) \quad X^{\text{an}} \setminus (V \cup D) = A_1 \cup \cdots \cup A_n \cup \bigcup_{\alpha} B_{\alpha},$$

where each A_i is a generalized open annulus and $\{B_{\alpha}\}$ is an infinite collection of open balls. The *skeleton* of $(X, V \cup D)$ is the subset

$$\Sigma(X, V \cup D) = V \cup D \cup \Sigma(A_1) \cup \cdots \cup \Sigma(A_n).$$

(This set is denoted $\widehat{\Sigma}(X, V \cup D)$ in [BPR11].) For each i and each α the inclusions $\mathbf{H}(A_i) \hookrightarrow \mathbf{H}(X^{\text{an}})$ and $\mathbf{H}(B_{\alpha}) \hookrightarrow \mathbf{H}(X^{\text{an}})$ are local isometries [BPR11, Proposition 5.60]. For each open ball B_{α} the map $B_{\alpha} \hookrightarrow X^{\text{an}}$ extends to an inclusion $\overline{B}_{\alpha} \hookrightarrow X^{\text{an}}$ sending the end of B_{α} to a point $x_{\alpha} \in V$. We say that B_{α} is *adjacent* to x_{α} . For each open annulus A_i the map $A_i \hookrightarrow X^{\text{an}}$ extends to a continuous map $\overline{A}_i \hookrightarrow X^{\text{an}}$ sending the ends of A_i to points $x_i, y_i \in V$. We say that A_i is *adjacent* to x_i and y_i . The length $\ell(e_i) \in \Lambda$ of the geodesic segment $e_i = \Sigma(A_i) \cup \{x_i, y_i\}$ is then the modulus of A_i . We say that V is *strongly semistable* if $x_i \neq y_i$ for each open annulus A_i . For each generalized open annulus A_i the map $A_i \hookrightarrow X^{\text{an}}$ extends to a continuous map $\overline{A}_i \hookrightarrow X^{\text{an}}$ sending the end of A_i to a point $x_i \in V$ and sending the puncture to a point $y_i \in D$. We say that A_i is *adjacent* to x_i and y_i . We define the length of $e_i = \Sigma(A_i) \cup \{x_i, y_i\}$ to be $\ell(e_i) = \infty$.

The skeleton $\Sigma = \Sigma(X, V \cup D)$ naturally has the structure of a Λ -metric graph with distinguished finite vertex set $V_f(\Sigma) = V$, infinite vertex set $V_{\infty}(\Sigma) = D$, and edges $\{e_1, \dots, e_n\}$ as above. Note that the Λ -points of Σ are exactly the type-2 points of X^{an} contained in Σ . For $x \in V$ the residue field $\widetilde{\mathcal{H}}(x)$ of the completed residue field $\mathcal{H}(x)$ at the type-2 point x is a finitely generated field extension of k of transcendence degree 1; we let C_x be the smooth k -curve with function field $\widetilde{\mathcal{H}}(x)$. For $x \in V$ we let $g(x)$ be the genus of C_x , and for $x \in \Sigma \setminus V$ we set $g(x) = 0$. These extra data give Σ the structure of an augmented Λ -metric graph. (In (3.22) we will see that Σ is in fact naturally a metrized complex of k -curves.) By the genus formula [BPR11, (5.45.1)], the genus of X is equal to the genus of the augmented Λ -metric graph Σ .

The open analytic domain $X^{\text{an}} \setminus \Sigma$ is isomorphic to an infinite disjoint union of open balls [BPR11, Lemma 5.18(3)]. If B is a connected component of $X^{\text{an}} \setminus \Sigma$, then the inclusion $B \hookrightarrow X^{\text{an}}$ extends to a map $\overline{B} \hookrightarrow X^{\text{an}}$ sending the end of B to a point of Σ . We define the *retraction* $\tau = \tau_{\Sigma} : X^{\text{an}} \rightarrow \Sigma$ by fixing Σ and sending a point $x \in X^{\text{an}}$ not in Σ to the end of the connected component of x in $X^{\text{an}} \setminus \Sigma$. This is a continuous map. If $x \in B_{\alpha}$ is in an open ball connected component of $X^{\text{an}} \setminus (V \cup D)$, then $\tau(x) \in V$ is the end of B_{α} , and if $x \in A_i$, then $\tau(x)$ coincides with the image of x under the retraction map $\tau : A_i \rightarrow \Sigma(A_i)$.

Here we collect some additional facts about skeleta from [BPR11, §5].

Proposition 3.12. *Let $(X, V \cup D)$ be a triangulated punctured curve with skeleton $\Sigma = \Sigma(X, V \cup D)$.*

- (1) The skeleton Σ is the set of points in X^{an} that do not admit an open neighborhood isomorphic to $\mathbf{B}(1)_+$ and disjoint from $V \cup D$.
- (2) Let V_1 be a semistable vertex set of (X, D) such that $V_1 \supset V$. Then $\Sigma(X, V_1 \cup D) \supset \Sigma(X, V \cup D)$ and $\tau_{\Sigma(X, V \cup D)} = \tau_{\Sigma(X, V_1 \cup D)} \circ \tau_{\Sigma(X, V_1 \cup D)}$.
- (3) Let $W \subset X^{\text{an}}$ be a finite set of type-2 points. Then there exists a semistable vertex set of (X, D) containing $V \cup W$.
- (4) Let $W \subset \Sigma$ be a finite set of type-2 points. Then $V \cup W$ is a semistable vertex set of (X, D) and $\Sigma(X, V \cup D) = \Sigma(X, V \cup W \cup D)$.
- (5) Let $x, y \in \Sigma \cap \mathbf{H}(X^{\text{an}})$. Then any geodesic segment from x to y in $\mathbf{H}(X^{\text{an}})$ is contained in Σ .
- (6) Let $x, y \in X^{\text{an}}$ be points of type-2 or 3 and let $[x, y]$ be a geodesic segment from x to y in $\mathbf{H}(X^{\text{an}})$. Then there exists a semistable vertex set V_1 of (X, D) such that $V_1 \supset V$ and $[x, y] \subset \Sigma(X, V_1 \cup D)$.

Definition 3.13. A skeleton of (X, D) is a subset of X^{an} of the form $\Sigma = \Sigma(X, V \cup D)$ for some semistable vertex set V of (X, D) . Such a semistable vertex set V is called a *vertex set for Σ* . A skeleton of X is a skeleton of (X, \emptyset) .

The augmented Λ -metric graph structure of a skeleton Σ of (X, D) does not depend on the choice of vertex set for Σ .

3.14. Modifying the skeleton. Let X be a smooth, proper, connected K -curve and let $D \subset X(K)$ be a finite set. Let Σ be a skeleton of (X, D) , let $y \in X^{\text{an}} \setminus \Sigma$, let B be the connected component of $X^{\text{an}} \setminus \Sigma$ containing y , and let $x = \tau(y) \in \Sigma$ be the end of B . If $y \in \mathbf{H}(X^{\text{an}})$, then the geodesic segment $[x, y] \subset \overline{B}$ is the unique geodesic segment in $\mathbf{H}(X^{\text{an}})$ connecting x and y . If $y \in X(K)$, then we define $[x, y]$ to be the geodesic segment $[x, y] \subset \overline{B}$ as in (3.1). The following strengthening of Proposition 3.12(3) will be important in the sequel.

Lemma 3.15. Let V be a semistable vertex set of (X, D) , let $\Sigma = \Sigma(X, V \cup D)$, let $W \subset X^{\text{an}}$ be a finite set of type-2 points, and let $E \subset X(K)$ be a finite set of type-1 points.

- (1) There exists a minimal semistable vertex set V_1 of $(X, D \cup E)$ which contains $V \cup W$, in the sense that any other such semistable vertex set contains V_1 .
- (2) Let B be a connected component of $X^{\text{an}} \setminus \Sigma$ with end x and let y_1, \dots, y_n be the points of $(W \cup E) \cap B$. Then

$$\Sigma(X, V_1 \cup D \cup E) \cap \overline{B} = [x, y_1] \cup \dots \cup [x, y_n]$$

if $n > 0$, and $\Sigma(X, V_1 \cup D \cup E) \cap \overline{B} = \{x\}$ otherwise. Therefore

$$\Sigma(X, V_1 \cup D \cup E) = \Sigma \cup \bigcup_{y \in W \cup E} [\tau_{\Sigma}(y), y].$$

- (3) The skeleton $\Sigma_1 = \Sigma(X, V_1 \cup D \cup E)$ is minimal in the sense that any other skeleton containing Σ and $W \cup E$ must contain Σ_1 .

Proof. To prove the first part we may assume that $W = \{y\}$ consists of a single type-2 point not contained in Σ and $E = \emptyset$, or $E = \{y\}$ is a single type-1 point not contained in D and $W = \emptyset$. In the first case one sees using Lemma 3.4 that $V_1 = V \cup \{y, \tau(y)\}$ is the minimal semistable vertex set of (X, D) containing V and y , and in the second case $V_1 = V \cup \{\tau(y)\}$ is the minimal semistable vertex set of $(X, D \cup \{y\})$ containing V . For W and E arbitrary, let V_1 be the minimal semistable vertex set of $(X, D \cup E)$ containing $V \cup W$ and let $\Sigma_1 = \Sigma(X, V_1 \cup D)$. If $E = \emptyset$, then it is clear from Proposition 3.12(5) that $[x, y_i] \subset \Sigma_1$ for each i ; the other inclusion is proved by induction on n , adding one point at a time as above. The case $E \neq \emptyset$ is similar and is left to the reader. The final assertion follows easily from the first two and Proposition 3.12. \blacksquare

Remark 3.16. The above lemma shows in particular that the metric graph $\Sigma(X, V_1 \cup D \cup E)$ is obtained from $\Sigma(X, V \cup D)$ by a sequence of tropical modifications and their inverses (see Definition II.2.12 and Example II.2.16). It is easy to see that any tropical modification of $\Sigma(X, V \cup D)$ is

of the form $\Sigma(X, V_1 \cup D \cup E)$ for a semistable vertex set V_1 which contains V and a finite subset $E \subset X(K)$.

3.17. The minimal skeleton. The Euler characteristic of $X \setminus D$ is defined to be $\chi(X \setminus D) = 2 - 2g(X) - \#D$, where $g(X)$ is the genus of X . We say that (X, D) is *stable* if $\chi(X \setminus D) < 0$. A semistable vertex set V of (X, D) is *minimal* if there is no semistable vertex set of (X, D) properly contained in V , and V is *stable* if there is no $x \in V$ of genus zero (resp. one) and valency less than 3 (resp. one) in $\Sigma(X, V \cup D)$. It is clear that minimal semistable vertex sets exist. A skeleton Σ of (X, D) is *minimal* if there is no skeleton of (X, D) properly contained in Σ .

The following consequence of the stable reduction theorem can be found in [BPR11, Theorem 5.49].

Proposition 3.18. *Let V be a minimal semistable vertex set of (X, D) and let $\Sigma = \Sigma(X, V \cup D)$.*

- (1) *If $\chi(X \setminus D) \leq 0$, then Σ is the unique minimal skeleton of (X, D) ; moreover, Σ is equal to the set of points of $X^{\text{an}} \setminus D$ that do not admit an open neighborhood isomorphic to $\mathbf{B}(1)_+$ and disjoint from D .*
- (2) *If $\chi(X \setminus D) < 0$, then V is the unique minimal semistable vertex set of (X, D) , V is stable, and*

$$V = \{x \in \Sigma : x \text{ has valency } \geq 3 \text{ or genus } \geq 1\}.$$

Remark 3.19. We have $\chi(X \setminus D) > 0$ if and only if $X \cong \mathbf{P}^1$ and D contains at most one point; in this case, any type-2 point of X^{an} serves as a minimal semistable vertex set of (X, D) , and there does not exist a unique minimal skeleton. If $\chi(X \setminus D) = 0$, then either $X \cong \mathbf{P}^1$ and D consists of two points, or X is an elliptic curve and D is empty. In the first case, the minimal skeleton Σ of (X, D) is the extended line connecting the points of D and any type-2 point on Σ is a minimal semistable vertex set. If X is an elliptic curve and X^{an} contains a type-2 point x of genus 1, then $\{x\}$ is both the unique minimal semistable vertex set and the minimal skeleton. Otherwise X is a Tate curve, Σ is a circle, and any type-2 point of Σ is a minimal semistable vertex set. See [BPR11, Remark 5.51].

3.20. Tangent directions and the slope formula. As above we let X be a smooth, proper, connected algebraic K -curve. A continuous function $F : X^{\text{an}} \rightarrow \mathbf{R} \cup \{\pm\infty\}$ is called *piecewise affine* provided that $F(\mathbf{H}(X^{\text{an}})) \subset \mathbf{R}$ and $F \circ \alpha : [a, b] \rightarrow \mathbf{R}$ is a piecewise affine function for every geodesic segment $\alpha : [a, b] \hookrightarrow \mathbf{H}(X^{\text{an}})$. To any point $x \in X^{\text{an}}$ is associated a set T_x of *tangent directions* at x , defined as the set of germs of geodesic segments in X^{an} beginning at x . If $F : X^{\text{an}} \rightarrow \mathbf{R} \cup \{\pm\infty\}$ is a piecewise affine function and $v \in T_x$ we denote by $d_v F(x)$ the outgoing slope of F in the direction v . We say that F is *harmonic* at a point $x \in X^{\text{an}}$ provided that there are only finitely many $v \in T_x$ with $d_v F(x) \neq 0$, and $\sum_{v \in T_x} d_v F(x) = 0$. See [BPR11, 5.65].

Let $x \in X^{\text{an}}$ be a type-2 point, let $\mathcal{H}(x)$ be the completed residue field of X^{an} at x , and let $\widetilde{\mathcal{H}}(x)$ be its residue field, as in (3.11). Then $\widetilde{\mathcal{H}}(x)$ is a finitely generated field extension of k of transcendence degree 1, and there is a canonical bijection between the set T_x of tangent directions to X^{an} at x and the set $\text{DV}(\widetilde{\mathcal{H}}(x)/k)$ of discrete valuations on $\widetilde{\mathcal{H}}(x)$ which are trivial on k . For $v \in T_x$ we let $\text{ord}_v : \widetilde{\mathcal{H}}(x) \rightarrow \mathbf{Z}$ denote the corresponding valuation. Let f be a nonzero analytic function on X^{an} defined on a neighborhood of x , let $c \in K^\times$ be such that $|c| = |f(x)|$, and let $\widetilde{f}_x \in \widetilde{\mathcal{H}}(x)$ denote the residue of $c^{-1}f$. Then \widetilde{f}_x is defined up to multiplication by k^\times , so for any $v \in T_x$ the integer $\text{ord}_v(\widetilde{f}_x)$ is well-defined.

The following theorem is called the slope formula in [BPR11, Theorem 5.69]:

Theorem 3.21. *Let $f \in K(X)^\times$ be a nonzero rational function on X and let $F = -\log|f| : X^{\text{an}} \rightarrow \mathbf{R} \cup \{\pm\infty\}$. Let $D \subset X(K)$ contain the zeros and poles of f and let Σ be a skeleton of (X, D) . Then:*

- (1) *$F = F \circ \tau$, where $\tau : X^{\text{an}} \rightarrow \Sigma$ is the retraction.*
- (2) *F is piecewise affine with integer slopes, and F is affine on each edge of Σ (with respect to a choice of vertex set V).*
- (3) *If x is a type-2 point of X^{an} and $v \in T_x$, then $d_v F(x) = \text{ord}_v(\widetilde{f}_x)$.*
- (4) *F is harmonic at all points of $X^{\text{an}} \setminus D$.*

(5) Let $x \in X(K)$ and let v be the unique tangent direction at x . Then $d_v F(x) = \text{ord}_x(f)$.

3.22. The skeleton as a metrized complex of curves. Let $(X, V \cup D)$ be a triangulated punctured curve with skeleton $\Sigma = \Sigma(X, V \cup D)$. Recall that Σ is an augmented Λ -metric graph with infinite vertices D . We enrich Σ with the structure of a Λ -metrized complex of k -curves as follows. For $x \in V$ let C_x be the smooth, proper, connected k -curve with function field $\widetilde{\mathcal{H}}(x)$ as in (3.11). By definition C_x has genus $g(x)$. We have natural bijections

$$(3.22.1) \quad T_x \cong \text{DV}(\widetilde{\mathcal{H}}(x)/k) \cong C_x(k)$$

where the first bijection $v \mapsto \text{ord}_v$ is defined in (3.20) and the second associates to a closed point $\xi \in C_x(k)$ the discrete valuation ord_ξ on the function field $\widetilde{\mathcal{H}}(x)$ of C_x . Let $v \in T_x$ be a tangent direction and define $\text{red}_x(v)$ to be the point of C_x corresponding to the discrete valuation $\text{ord}_v \in \text{DV}(\widetilde{\mathcal{H}}(x)/k)$. These data make Σ into a Λ -metrized complex of k -curves.

3.23. Lifting metrized complexes of curves. We now prove that every metrized complex of curves over k arises as the skeleton of a smooth, proper, connected K -curve. This fact appears in the literature in various contexts (over discretely-valued fields): see for instance [Bak08, Appendix B] and [Sai96, Lemme 6.3]. For this reason we only sketch a proof using our methods.

Theorem 3.24. *Let \mathcal{C} be a Λ -metrized complex of k -curves. There exists a triangulated punctured curve $(X, V \cup D)$ such that the skeleton $\Sigma(X, V \cup D)$ is isomorphic to \mathcal{C} .*

Proof. Let C be a smooth, proper, connected k -curve. By elementary deformation theory, there is a smooth, proper admissible formal R -scheme \mathfrak{C} with special fiber C . By GAGA the analytic generic fiber \mathfrak{C}_K is the analytification of a smooth, proper, connected K -curve \mathcal{C} . There is a reduction map $\text{red} : \mathfrak{C}_K \rightarrow C$ from the analytic generic fiber of \mathfrak{C} to (the set underlying) C , under which the inverse image of the generic point of C is a single distinguished point x . The set $\{x\}$ is a semistable vertex set of \mathcal{C} , with associated skeleton also equal to $\{x\}$. Moreover, there is a canonical identification of $\widetilde{\mathcal{H}}(x)$ with the field of rational functions on \overline{C} by [Ber90, Proposition 2.4.4]. See (5.6) for a more detailed discussion of the relationship between semistable vertex sets and admissible formal models.

Let Γ be the metric graph underlying \mathcal{C} . Let V be the vertices of Γ and let E be its edges. By adding valence-2 vertices we may assume that Γ has no loop edges. Assume for the moment that Γ has no infinite edges. For a vertex $x \in V$ let C_x denote the smooth, proper, connected k -curve associated to x , and choose an admissible formal curve \mathfrak{C}_x with special fiber isomorphic to C_x as above. For clarity we let $\text{red}_{\mathfrak{C}_x}$ denote the reduction map $(\mathfrak{C}_x)_K \rightarrow C_x$. By [BL93, Proposition 2.2], for every $\bar{x} \in C_x(k)$ the formal fiber $\text{red}_{\mathfrak{C}_x}^{-1}(\bar{x})$ is isomorphic to $\mathbf{B}(1)_+$. Let e be an edge of Γ with endpoints x, y , and let $\bar{x}_e \in C_x(k), \bar{y}_e \in C_y(k)$ be the reductions of the tangent vectors in the direction of e at x, y , respectively. Remove closed balls from $\text{red}_{\mathfrak{C}_x}^{-1}(\bar{x}_e)$ and $\text{red}_{\mathfrak{C}_y}^{-1}(\bar{y}_e)$ whose radii are such that the remaining open annuli have modulus equal to $\ell(e)$. We form a new analytic curve X^{an} by gluing these annuli together using some isomorphism of annuli for each edge e . The resulting curve is proper, hence is the analytification of an algebraic curve X . By construction the image of the set of distinguished points of the curves $(\mathfrak{C}_x)_K$ in X^{an} is a semistable vertex set, and the resulting skeleton (considered as a metric graph) is isomorphic to Γ .

If Γ does have infinite edges, then we apply the above procedure to the finite part of Γ , then puncture the curve X in the formal fibers over the smooth points of the residue curves which correspond to the directions of the infinite tails of Γ . \blacksquare

As an immediate application, we obtain the following property of the ‘‘abstract tropicalization map’’ from the moduli space of stable marked curves to the moduli space of stable abstract tropical curves:

Corollary 3.25. *If g and n are nonnegative integers with $2 - 2g - n < 0$, the natural map $\text{trop} : M_{g,n} \rightarrow M_{g,n}^{\text{trop}}$ (see [BPR11, Remark 5.52]) is surjective.*

See [ACP12] for an interpretation of the above map as a contraction from $M_{g,n}^{\text{an}}$ onto its skeleton (which also implies surjectivity).

4. MORPHISMS BETWEEN CURVES AND THEIR SKELETA

This section is a relative version of the previous one, in that we propose to study the behavior of semistable vertex sets and skeleta under finite morphisms of curves. We introduce finite morphisms of triangulated punctured curves and we prove that any finite morphism of punctured curves can be enriched to a finite morphism of triangulated punctured curves. This powerful result can be used to prove the simultaneous semistable reduction theorems of Coleman, Liu–Lorenzini, and Liu, which we do in Section 5. It is interesting to note that we use only analytic methods on analytic K -curves, making (almost) no explicit reference to semistable models; hence our proofs of the results of Liu–Lorenzini and Liu are very different from theirs. Using a relative version of the slope formula, we also show that a finite morphism of triangulated punctured curves induces (by restricting to skeleta) a finite harmonic morphism of metrized complexes of curves, which will be crucial in the sequel.

4.1. Morphisms between open balls and generalized open annuli. The main results of this section rest on a careful study of the behavior of certain morphisms between open balls and generalized open annuli. Some of the lemmas and intermediate results appear in some form in the literature — see in particular [Ber90] and [Ber93] — although they are hard to find in the form we need them. As the proofs are not difficult, for the convenience of the reader we include complete arguments.

Lemma 4.2. *Let $a, a' \in \mathfrak{m}_R$ and let $\varphi : \mathbf{S}(a')_+ \rightarrow \mathbf{S}(a)_+$ be a morphism.*

- (1) *If $\text{val} \circ \varphi$ is constant on $\Sigma(\mathbf{S}(a')_+)$, then $\varphi(\mathbf{S}(a')_+)$ is contained in $\mathbf{S}(a)_+ \setminus \Sigma(\mathbf{S}(a)_+)$.*
- (2) *If $\text{val} \circ \varphi$ is not constant on $\Sigma(\mathbf{S}(a')_+)$, then $\varphi(\Sigma(\mathbf{S}(a')_+)) \subset \Sigma(\mathbf{S}(a))$ and the restriction of φ to $\Sigma(\mathbf{S}(a')_+)$ has the form $x \mapsto m \cdot x + b$ for some nonzero integer m and some $b \in \Lambda$.*
- (3) *If φ extends to a continuous map $\bar{\varphi} : \overline{\mathbf{S}(a')_+} \rightarrow \overline{\mathbf{S}(a)_+}$ taking an end of $\mathbf{S}(a')_+$ to an end of $\mathbf{S}(a)_+$, then $\text{val} \circ \varphi$ is not constant on $\Sigma(\mathbf{S}(a')_+)$.*

Proof. Part (2) is exactly [BPR11, Proposition 5.5], so suppose that $\text{val} \circ \varphi$ is constant on $\Sigma(\mathbf{S}(a))$. The map $\varphi : \mathbf{S}(a')_+ \rightarrow \mathbf{S}(a)_+ \subset \mathbf{G}_m^{\text{an}}$ is given by a unit f on $\mathbf{S}(a')_+$. By [BPR11, Proposition 5.2], we can write $f = \alpha(1 + g)$, where $\alpha \in K^\times$ and $|g(x')| < 1$ for all $x' \in \mathbf{S}(a')_+$. Hence $|\varphi(x') - \alpha| < |\alpha|$ for all $x' \in \mathbf{S}(a')_+$; here $|\varphi(x') - \alpha|$ should be interpreted as the absolute value of the function $f - \alpha$ in $\mathcal{H}(x')$, or equivalently as the absolute value of the function $t - \alpha$ in $\mathcal{H}(\varphi(x))$, where t is a parameter on \mathbf{G}_m . But $|t(x) - \alpha| \geq |\alpha|$ for every $x \in \Sigma(\mathbf{S}(a))$, so $\varphi(\mathbf{S}(a')_+) \cap \Sigma(\mathbf{S}(a)) = \emptyset$.

Let x' be an end of $\mathbf{S}(a')_+$, let x be an end of $\mathbf{S}(a)_+$, and suppose that $\bar{\varphi}(x') = x$. Since x' is a limit point of $\Sigma(\mathbf{S}(a')_+)$, there exists a sequence of points $x'_1, x'_2, \dots \in \Sigma(\mathbf{S}(a')_+)$ converging to x' . Then $\varphi(x'_1), \varphi(x'_2), \dots$ converge to x , so $\text{val}(\varphi(x'_1)), \text{val}(\varphi(x'_2)), \dots$ converges to $\text{val}(x)$. Since each $\text{val}(\varphi(x'_i)) \in (0, \text{val}(a))$ but $\text{val}(x) \notin (0, \text{val}(a))$, $\text{val} \circ \varphi$ is not constant on $\Sigma(\mathbf{S}(a')_+)$. \blacksquare

Lemma 4.3. *Let B' be an open ball, let A be a generalized open annulus, and let $\varphi : B' \rightarrow A$ be a morphism. Then $\varphi(B') \cap \Sigma(A) = \emptyset$, and φ extends to a continuous map $\bar{B}' \rightarrow A$.*

Proof. The assertions of the Lemma are clear if φ is constant, so assume that φ is non-constant. We identify A with $\mathbf{S}(a)_+$ for $a \in \mathfrak{m}_R$ and B' with $\mathbf{B}(1)_+$. Any unit on B' has constant absolute value, so $\text{val} \circ \varphi$ is constant on B' . Since $\mathbf{B}(1)_+ \setminus \{0\}$ is a generalized open annulus, and since the type-1 point 0 maps to a type-1 point of A , it follows from Lemma 4.2 that $\varphi(B') \cap \Sigma(A) = \emptyset$. Let B be the connected component of $A \setminus \Sigma(A)$ containing $\varphi(B')$. Then the morphism $B' \rightarrow B \hookrightarrow A$ extends to a continuous map $\bar{B}' \rightarrow \bar{B} \hookrightarrow A$. \blacksquare

Definition 4.4. Let T be a metric space and let $m \in \mathbf{R}_{>0}$. A continuous injection $\varphi : [a, b] \hookrightarrow T$ is an *embedding with expansion factor m* provided that $r \mapsto \varphi(r/m) : [ma, mb] \hookrightarrow T$ is a geodesic segment. A continuous injection $\varphi : [a, b] \hookrightarrow T$ is *piecewise affine* provided that there exist $a = a_0 < a_1 < \dots < a_r = b$ and $m_1, \dots, m_r \in \mathbf{R}_{>0}$ such that $\varphi|_{[a_{i-1}, a_i]}$ is an embedding with expansion factor m_i for each $i = 1, \dots, r$.

Lemma 4.5. *Let B, B' be open balls, let x' be the end of B , and let $\varphi : B' \rightarrow B$ be a morphism with finite fibers. Let $y' \in B'$, let $y = \varphi(y')$, and let $x = \varphi(x')$. Then the restriction of φ to the geodesic*

segment $[x', y']$ is injective, and $\varphi([x', y'])$ is equal to the geodesic segment $[x, y]$. If in addition there exists $N > 0$ such that all fibers of φ have fewer than N elements, then $\varphi|_{[x', y']}$ is piecewise affine.

Proof. First suppose that y' does not have type 4. Choose isomorphisms $B' \cong \mathbf{B}(1)_+$ such that $0 > y'$ and $B \cong \mathbf{B}(1)_+$ such that $\varphi(0) = 0$. Since $y' \in [x', 0]$ we may replace $[x', y']$ by the larger geodesic segment $[x', 0]$ to assume that $y' = 0$. Define $F : \Sigma(\mathbf{S}(0)_+) \rightarrow \mathbf{R}$ by $F(x) = \text{val} \circ \varphi(x) = -\log |\varphi(x)|$. By [BPR11, Proposition 5.10], F is a piecewise affine function, and for any point $z' \in \Sigma(\mathbf{S}(0)_+)$ at which F is differentiable, the derivative of F is equal to the number of zeros q' of φ with $\text{val}(q') > \text{val}(z')$. It follows that F is monotonically increasing, so φ is injective on $[x', 0]$. Let

$$Z = \{\tau(q') : q' \in \mathbf{S}(0)_+, \varphi(q') = 0\} \subset \Sigma(\mathbf{S}(0)_+).$$

If $C \subset \Sigma(\mathbf{S}(0)_+) \setminus Z$ is a connected component, then $A' = \tau^{-1}(C)$ is a generalized open annulus mapping to $\mathbf{S}(0)_+$. By the above, $\text{val} \circ \varphi$ is not constant on $\Sigma(A') = \Sigma(\mathbf{S}(0)_+) \cap A'$, so by Lemma 4.2, $\varphi(\Sigma(A')) \subset \Sigma(\mathbf{S}(0)_+)$. Since $\Sigma(\mathbf{S}(0)_+) \setminus Z$ is dense in $\Sigma(\mathbf{S}(0)_+)$, we have that $[x', 0] = \Sigma(\mathbf{S}(0)_+) \cup \{0\}$ maps bijectively onto $[x, 0]$. It is clear in this case that the restriction of φ to the closure of any connected component of $\Sigma(\mathbf{S}(0)_+) \setminus Z$ is an embedding with integer expansion factor, so $\varphi|_{[x', 0]}$ is piecewise affine. Note that $\varphi|_{[x', 0]}$ changes expansion factor at most $\#\varphi^{-1}(0)$ times.

Now suppose that y' has type 4. Suppose that $z', w' \in [x', y']$ are two distinct points such that $\varphi(z') = \varphi(w')$. Assume without loss of generality that $z' < w'$. Since w' and z' have the same image under φ , they both have the same type, so $w' \neq y'$ because y' is the only type-4 point in $[x', y']$. Applying the above to the geodesic $[x', w']$ gives a contradiction. Therefore, φ is injective on $[x', y']$, so $\varphi([x', y']) = [x, y]$ since B is uniquely path-connected ([BR10, Corollary 1.14]).

Assume now that all fibers of φ have size bounded by N . The above argument proves that for all $z' < y'$, the restriction of φ to $[x', z']$ is piecewise affine, and $\varphi|_{[x', z']}$ changes expansion factor at most N times. Therefore there exists $z'_0 < y'$ and $m \in \mathbf{Z}_{>0}$ such that for all $z' \in [z'_0, y']$, the restriction of φ to $[z'_0, z']$ is an embedding with expansion factor m . It follows that $\varphi|_{[z'_0, y']}$ is an embedding with expansion factor m , so $\varphi|_{[x', y']}$ is piecewise affine. \blacksquare

Lemma 4.6. *Let B, B' be open balls and let $\varphi : B' \rightarrow B$ be a morphism. Suppose that φ is open, separated, and has finite fibers, so it extends canonically to a continuous map $\varphi : \overline{B'} \rightarrow \overline{B}$ by Lemma 3.3. Let x' be the end of B' and let $x = \varphi(x') \in \overline{B}$. Then $B_1 = \varphi(B')$ is an open ball connected component of $\overline{B} \setminus \{x\}$ and $\varphi : B' \rightarrow B_1$ is finite and order-preserving. In particular, if x is the end of B , then $\varphi : B' \rightarrow B$ is finite and order-preserving.*

Proof. Let $y' \in B'$. By Lemma 4.5, the restriction of φ to the geodesic $[x', y']$ is injective, so $\varphi(y') \neq x$. Therefore $x \notin B_1$. Let C be the connected component of $B \setminus \{x\}$ containing B_1 . Since x is an end of C and x' maps to x , Lemma 4.3 implies that C is not an open annulus, so it is an open ball. By hypothesis, B_1 is an open subset of C ; since $B' \cup \{x'\}$ is compact, $\varphi(B' \cup \{x'\}) = B_1 \cup \{x\}$ is closed in \overline{B} , and therefore $B_1 = (B_1 \cup \{x\}) \cap C$ is closed in C . Since C is connected, we have $B_1 = C$.

Since φ has finite fibers, $\varphi : B' \rightarrow B_1$ is finite if and only if it is proper by [Ber90, Corollary 3.3.8]. If $D \subset B_1$ is compact, then $\varphi^{-1}(D)$ is compact since $\varphi^{-1}(D)$ is closed as a subset of the compact space $\overline{B'}$. One has $\text{Int}(B'/B_1) = B'$ by [Ber90, Proposition 3.1.3(i)] since B' is a boundaryless K -analytic space. Therefore φ is proper.

The fact that $\varphi : B' \rightarrow B_1$ is order-preserving follows immediately from Lemma 4.5. \blacksquare

4.7. Let $\varphi : B' \rightarrow B$ be a finite morphism of open balls. Since an open ball is a smooth curve, φ is flat in the sense that if $\mathcal{M}(\mathcal{A}) \subset B$ is an affinoid domain and $\varphi^{-1}(\mathcal{M}(\mathcal{A})) = \mathcal{M}(\mathcal{A}')$, then \mathcal{A}' is a (finite) flat \mathcal{A} -algebra. For $x \in \mathcal{M}(\mathcal{A})$ the fiber over x is

$$\varphi^{-1}(x) = \mathcal{M}(\mathcal{A}' \widehat{\otimes}_{\mathcal{A}} \mathcal{H}(x)) = \mathcal{M}(\mathcal{A}' \otimes_{\mathcal{A}} \mathcal{H}(x)),$$

where the second equality holds because \mathcal{A}' is a finite \mathcal{A} -algebra. It follows that for $x \in B$ the quantity $\sum_{x' \mapsto x} \dim_{\mathcal{H}(x)} \mathcal{O}_{\varphi^{-1}(x), x'}$ is independent of x ; we call this number the *degree* of φ .

Proposition 4.8. *Let B, B' be open balls and let $\varphi : B' \rightarrow B$ be a finite morphism. Let x' be the end of B' , let $x = \varphi(x')$ be the end of B , let $y \in B$, and let $y'_1, \dots, y'_n \in B'$ be the inverse images of y in B' .*

Then

$$\varphi^{-1}([x, y]) = [x', y'_1] \cup \cdots \cup [x', y'_n].$$

Proof. Let $T' = [x', y'_1] \cup \cdots \cup [x', y'_n]$. The inclusion $T' \subset \varphi^{-1}([x, y])$ follows from Lemma 4.5. First we claim that for $z \in [x, y]$ near enough to x , there is only one preimage of z in B' . Shrinking $[x, y]$ if necessary, we may assume that y has type 2. Choose a type-1 point $w \in B$ such that $w > y$, so $(x, y) \subset (x, w) = \Sigma(B \setminus \{w\})$. Let $w' \in B'$ be a preimage of w and choose $z' \in \Sigma(B' \setminus \{w'\})$ such that $z' \leq \tau(q')$ for all $q' \in \varphi^{-1}(w)$, where $\tau : B' \setminus \{w'\} \rightarrow \Sigma(B' \setminus \{w'\})$ is the retraction. Let A' be the open annulus $\tau^{-1}((x', z')) \subset B'$. Then $\varphi(A') \subset B \setminus \{w\}$, and φ takes the end x' of A' to the end x of $B \setminus \{w\}$, so by Lemma 4.2(2,3), $\varphi(\Sigma(A')) \subset \Sigma(B \setminus \{w\})$ and the map $\Sigma(A') \rightarrow \Sigma(B \setminus \{w\})$ is injective. If $u' \in A' \setminus \Sigma(A')$, then u' is contained in an open ball in A' , so $\varphi(u') \notin \Sigma(B \setminus \{w\})$ by Lemma 4.3. Therefore every point of $\Sigma(B \setminus \{w\})$ has at most one preimage in A' . Let $u' \in B' \setminus A'$, so $u' \geq z'$. Then $\varphi(u') \geq \varphi(z')$, so every point $u \in (x, y]$ with $u < \varphi(z')$ has exactly one preimage in B' (note that $\varphi(z') \in \Sigma(B \setminus \{w\})$ by Lemma 4.5).

Let d be the degree of the finite morphism φ , so for every $z \in B$ we have

$$d = \sum_{z' \in \varphi^{-1}(z)} \dim_{\mathcal{H}(z)} \mathcal{O}_{\varphi^{-1}(z), z'}.$$

For $y \in B$ as in the statement of the Proposition, define a function $\delta : (x, y] \rightarrow \mathbf{Z}$ by

$$\delta(z) = \sum_{\substack{z' \in \varphi^{-1}(z) \\ z' \in T'}} \dim_{\mathcal{H}(z)} \mathcal{O}_{\varphi^{-1}(z), z'}.$$

Clearly $\delta(z) \leq d$ for all $z \in (x, y]$, and $\delta(z) = d$ if and only if $\varphi^{-1}(z) \subset T'$. By definition of T' we have $\delta(y) = d$, and by the above, $\delta(z) = d$ for $z \in (x, y]$ close enough to x (any geodesic segment $[x', y'_i]$ surjects onto $[x, y]$, hence contains the unique preimage of z). Therefore it is enough to show that $\delta(z_1) \geq \delta(z_2)$ if $z_1 \leq z_2$.

If $z \in B$ is a point of type 2 or 3, then $B_z := \{w \in B : w \geq z\}$ is a (not necessarily strict) affinoid subdomain of B . If $w' \in B'$ is such that $\varphi(w') \geq z$, then $\varphi([x', w']) = [x, \varphi(w')]$ is a geodesic containing z , so there exists $z' \in [x', w']$ mapping to z . It follows that

$$\varphi^{-1}(B_z) = \coprod_{z' \mapsto z} \{w' \in B' : w' \geq z'\} = \coprod_{z' \mapsto z} B_{z'}$$

is a disjoint union of affinoid domains. Hence each map $B_{z'} \rightarrow B_z$ is finite, and its degree is equal to $\dim_{\mathcal{H}(z)} \mathcal{O}_{\varphi^{-1}(z), z'}$. Let $z_1, z_2 \in (x, y]$ and assume that $z_1 < z_2$, so z_1 has type 2 or 3. For $z'_1 \in \varphi^{-1}(z_1)$ we have

$$\dim_{\mathcal{H}(z_1)} \mathcal{O}_{\varphi^{-1}(z_1), z'_1} = \sum_{\substack{z'_2 \mapsto z_2 \\ z'_2 \geq z'_1}} \dim_{\mathcal{H}(z_2)} \mathcal{O}_{\varphi^{-1}(z_2), z'_2}.$$

Summing over all $z'_1 \in \varphi^{-1}(z_1) \cap T'$, we obtain

$$\begin{aligned} \delta(z_1) &= \sum_{\substack{z'_1 \mapsto z_1 \\ z'_1 \in T'}} \dim_{\mathcal{H}(z_1)} \mathcal{O}_{\varphi^{-1}(z_1), z'_1} = \sum_{\substack{z'_1 \mapsto z_1 \\ z'_1 \in T'}} \sum_{\substack{z'_2 \mapsto z_2 \\ z'_2 \geq z'_1}} \dim_{\mathcal{H}(z_2)} \mathcal{O}_{\varphi^{-1}(z_2), z'_2} \\ &\geq \sum_{\substack{z'_1 \mapsto z_1 \\ z'_1 \in T'}} \sum_{\substack{z'_2 \mapsto z_2 \\ z'_2 \geq z'_1 \\ z'_2 \in T'}} \dim_{\mathcal{H}(z_2)} \mathcal{O}_{\varphi^{-1}(z_2), z'_2} = \delta(z_2), \end{aligned}$$

where the final equality holds because if $z'_2 \in \varphi^{-1}(z_2) \cap T'$, then there exists $z'_1 \in \varphi^{-1}(z_1) \cap T'$ such that $z'_2 \geq z'_1$, namely, the unique point of $[x', z'_2]$ mapping to z_1 . \blacksquare

Proposition 4.9. *Let $a \in \mathfrak{m}_R \setminus \{0\}$, let $A' = \mathbf{S}(a)_+$, let B be an open ball, and let $\varphi : A' \rightarrow B$ be a morphism with finite fibers. Suppose that each end of A' maps to the end of B or to a type-2 point*

of B under the induced map $\bar{A}' \rightarrow \bar{B}$. Let $\alpha = \varphi \circ \sigma : [0, \text{val}(a)] \rightarrow \bar{B}$. Then there exist finitely many numbers $r_0, r_1, r_2, \dots, r_n \in \Lambda$ with $0 = r_0 < r_1 < r_2 < \dots < r_n = \text{val}(a)$ such that α is an embedding with nonzero integer expansion factor when restricted to each interval $[r_i, r_{i+1}]$. In other words, α is piecewise affine with integer expansion factors. Moreover, the image of α is a geodesic segment between type-2 points of B .

Proof. Let x' be the end $\sigma(0)$ of A' and let $x = \varphi(x')$. Suppose first that x is the end of B . Choose an identification $B \cong \mathbf{B}(1)_+$, and let $r_+ = \min\{\text{val}(y') : \varphi(y') = 0\} > 0$. Let A'_+ be the open annulus $\text{val}^{-1}((0, r_+)) \subset A'$. Then $\varphi(A'_+) \subset \mathbf{B}(1)_+ \setminus \{0\} = \mathbf{S}(0)_+$ and the end x' of A'_+ maps to the end x of $\mathbf{S}(0)_+$, so by Lemma 4.2(2,3), α is an embedding with (nonzero) integer expansion factor when restricted to $[0, r_+]$.

Now suppose that x is not the end of B . Let $r = \min\{\text{val}(y') : \varphi(y') = x\} > 0$ and let $A'' = \text{val}^{-1}((0, r))$. Then $\varphi(A'')$ is contained in a connected component C of $B \setminus \{x\}$, and φ takes the end x' of A'' to the end x of C . If C is an open annulus, then α is an embedding with integer expansion factor when restricted to $[0, r]$ by Lemma 4.2(2,3), and if C is an open ball, then we proceed as above to find $r_+ \in (0, r) \cap \Lambda$ such that α is an embedding with integer expansion factor when restricted to $[0, r_+]$.

Applying the above argument to the morphism φ composed with the automorphism $t \mapsto a/t$ of $\mathbf{S}(a)_+$ (which interchanges the two ends), we find that there exists $r_- \in [0, \text{val}(a)) \cap \Lambda$ such that α is an embedding with integer expansion factor when restricted to $[r_-, \text{val}(a)]$. Let $s \in (0, \text{val}(a)) \cap \Lambda$. Replacing A' with the annulus $\text{val}^{-1}((0, s))$ (resp. $\text{val}^{-1}((s, \text{val}(a)))$), the above arguments then provide us with $s_+ \in (s, \text{val}(a)) \cap \Lambda$ (resp. $s_- \in [0, s) \cap \Lambda$) such that α is an embedding with integer expansion factor when restricted to $[s, s_+]$ (resp. $[s_-, s]$). The first assertions now follow because there is a finite subcover of the open covering

$$\{[0, r_+]\} \cup \{[r_-, \text{val}(a)]\} \cup \{(s_-, s_+) : s \in (0, r) \cap \Lambda\}$$

of the compact space $[0, r]$.

As for the final assertion, choose $0 = r_0 < r_1 < r_2 < \dots < r_n = \text{val}(a)$ such that α is an embedding with integer expansion factor on each $[r_i, r_{i+1}]$. Let $i_0 \in \{0, 1, \dots, n\}$ be the largest integer such that $\alpha(r_{i_0}) > \alpha(r_i)$ (in the canonical partial ordering on B) for all $i < i_0$. If $i_0 = n$, then we are done, so assume that $i_0 < n$. Let $y = \alpha(r_{i_0}) \in B$, and choose an identification $B \cong \mathbf{B}(1)_+$ such that $0 > y$. If $F = -\log|\varphi|$, then by [BPR11, Proposition 5.10], at every point $r \in (0, \text{val}(a))$ the change in slope of F at r is equal to the negative of the number of zeros of φ with valuation r (cf. the proof of Lemma 4.5); in particular, the slope of F can only *decrease*. By construction F is monotonically increasing on $[0, r_{i_0}]$. Since $\alpha([r_{i_0}, r_{i_0+1}])$ is a geodesic segment, it meets y only at $\alpha(r_{i_0})$. The image of (r_{i_0}, r_{i_0+1}) under α is not contained in an open ball connected component of $B \setminus \{y\}$ because $\alpha(r_{i_0+1}) \not\asymp y$; hence F is *decreasing* on an interval $[r_{i_0}, r_{i_0} + \varepsilon]$ for some $\varepsilon > 0$. Since the slope of F can only decrease, it follows that F is monotonically decreasing on $[r_{i_0}, r_n]$. It follows immediately from this that $\alpha([0, \text{val}(a)]) = [x, y]$ or $\alpha([0, \text{val}(a)]) = [\alpha(\text{val}(a)), y]$, whichever segment is larger. ■

4.10. Morphisms between curves and skeleta. In what follows we fix smooth, connected, proper algebraic K -curves X, X' and a finite morphism $\varphi : X' \rightarrow X$. Let $D \subset X(K)$ and $D' \subset X'(K)$ be finite sets of closed points. The map on analytifications $\varphi : X'^{\text{an}} \rightarrow X^{\text{an}}$ is finite and open by [Ber90, Lemma 3.2.4].

Proposition 4.11. *Let Σ' be a skeleton of (X', D') and let Σ be a skeleton of (X, D) . There exists a skeleton Σ_1 of $(X, D \cup \varphi(D'))$ containing $\Sigma \cup \varphi(\Sigma')$, and there is a minimal such Σ_1 with respect to inclusion.*

Proof. First we will prove the Proposition in the case $D = D' = \emptyset$. Let V be a vertex set for Σ and let V' be a vertex set for Σ' containing $\{\tau(y') : y' \in \varphi^{-1}(V)\}$. Let A' be an open annulus connected component of $X'^{\text{an}} \setminus V'$ and let $e' \subset \Sigma'$ be the associated edge. We claim that $\varphi(e')$ is a geodesic segment between type-2 points of X^{an} . Let C be the connected component of $X^{\text{an}} \setminus V$ containing $\varphi(A')$.

- (i) If C is an open ball then the claim follows immediately from Proposition 4.9.
- (ii) If C is an open annulus and $\varphi(A') \cap \Sigma(C) = \emptyset$, then A' is contained in an open ball in C because each connected component of $C \setminus \Sigma(C)$ is an open ball, so the claim is true as in (i).
- (iii) If C is an open annulus and $\varphi(A') \cap \Sigma(C) \neq \emptyset$, then $\varphi(e')$ is a geodesic segment in X^{an} by Lemma 4.2.

Applying the above to each edge e' of Σ' , we find that there exists a finite set of type-2 points $x_1, y_1, x_2, y_2, \dots, x_n, y_n \in X^{\text{an}}$ such that $\varphi(\Sigma') = \bigcup_{i=1}^n [x_i, y_i]$, where $[x_i, y_i]$ denotes a geodesic segment from x_i to y_i . Let

$$\Sigma_1 = \Sigma \cup \bigcup_{i=1}^n ([x_i, \tau_\Sigma(x_i)] \cup [y_i, \tau_\Sigma(y_i)]).$$

By Lemma 3.15(2) as applied to $W = \{x_1, y_1, \dots, x_n, y_n\}$, we have that Σ_1 is a skeleton of X , so by Proposition 3.12(5), Σ_1 contains $\Sigma \cup \varphi(\Sigma')$ since Σ_1 contains each geodesic segment $[x_i, y_i]$. Any other skeleton of X containing Σ and all x_i, y_i also contains the geodesics $[x_i, \tau_\Sigma(x_i)]$ and $[y_i, \tau_\Sigma(y_i)]$, so Σ_1 is the minimal such skeleton.

Now assume that $D \amalg D' \neq \emptyset$. By Lemma 3.15(3) there is a minimal skeleton Σ_1 of X containing Σ and $\varphi(D')$; replacing Σ by Σ_1 and D by $D \cup \varphi(D')$, we may assume without loss of generality that $\varphi(D') \subset D$. We will proceed by induction on the size of $D \amalg D'$. As above we let V be a vertex set for Σ and we let V' be a vertex set for Σ' containing $\{\tau(y') : y' \in \varphi^{-1}(V \cup D)\}$.

- (i) Suppose that D' is not empty. Let $x' \in D'$ and let Σ_1 be the minimal skeleton of (X, D) containing $\Sigma \cup \varphi(\Sigma(X', V' \cup D' \setminus \{x'\}))$. Let A' be the connected component of $X^{\text{an}} \setminus (V' \cup D')$ whose closure contains x' , let $x = \varphi(x') \in D$, and let A be the connected component of $X^{\text{an}} \setminus (V \cup D)$ whose closure contains x . Note that $\Sigma' = \Sigma(X', V' \cup D' \setminus \{x'\}) \cup \Sigma(A') \cup \{x'\}$. Clearly $\varphi(A') \subset A$, and $\varphi^{-1}(x) \cap A' = \emptyset$ by construction. Since φ takes the puncture of A' to the puncture of A , one shows as in the proof of Lemma 4.2 that $\varphi(\Sigma(A')) \subset \Sigma(A)$. Therefore Σ_1 contains $\Sigma \cup \varphi(\Sigma')$.
- (ii) Now suppose that $D' = \emptyset$ and $D \neq \emptyset$. Let $x \in D$ and let Σ_1 be the minimal skeleton of $(X, D \setminus \{x\})$ containing $\Sigma(X, V \cup D \setminus \{x\}) \cup \varphi(\Sigma')$. Let $\Sigma_2 = \Sigma_1 \cup [x, \tau(x)]$. It follows from Lemma 3.15(3) that Σ_2 is the minimal skeleton of (X, D) containing Σ_1 . ■

Let V be a semistable vertex set of (X, D) . Recall from (3.11) that a connected component C of $X^{\text{an}} \setminus (V \cup D)$ is *adjacent* to a vertex $x \in V$ provided that the closure of C in X^{an} contains x . Proposition 4.12(1) below is exactly [Ber90, Theorem 4.5.3] when $D = D' = \emptyset$.

Proposition 4.12. *Suppose that $D' = \varphi^{-1}(D)$ and that one of the following two conditions holds:*

- (1) $\chi(X \setminus D) \leq 0$ (hence also $\chi(X' \setminus D') \leq 0$), or
- (2) $\varphi^{-1}(V) \subset V'$.

In the situation of (1) let Σ (resp. Σ') be the minimal skeleton of (X, D) (resp. (X', D')), and in (2) let $\Sigma = \Sigma(X, V \cup D)$ and $\Sigma' = \Sigma(X', V' \cup D')$. Then $\varphi^{-1}(\Sigma) \subset \Sigma'$.

Proof. First suppose that (2) holds. Let $x' \in X^{\text{an}} \setminus \Sigma'$ and let B' be the connected component of $X^{\text{an}} \setminus \Sigma'$ containing x' , so B' is an open ball. By hypothesis, $\varphi(B')$ is contained in a connected component C of $X^{\text{an}} \setminus (V \cup D)$. If C is an open ball, then $\varphi(B') \cap \Sigma = \emptyset$ by the definition of $\Sigma(X, V \cup D)$. If C is a generalized open annulus, then $\varphi(B') \cap \Sigma = \varphi(B') \cap \Sigma(C) = \emptyset$ by Lemma 4.3. Therefore $\varphi(x') \notin \Sigma$.

Now suppose that (1) holds. Let V be a semistable vertex set of (X, D) such that $\Sigma = \Sigma(X, V \cup D)$. By subdividing edges of Σ and enlarging V if necessary, we may and do assume that Σ has no loop edges. By Proposition 3.18(2), no point in V of genus zero has valence one in Σ . First we claim that if $B' \subset X^{\text{an}} \setminus D'$ is an open analytic domain which is isomorphic to $\mathbf{B}(1)_+$, then $\varphi(B') \cap V = \emptyset$. If $\varphi^{-1}(V) \cap B'$ contains more than one point, then it is easy to see that there exists a smaller open ball $B'' \subset B'$ such that $\varphi^{-1}(V) \cap B''$ contains exactly one point. Replacing B' by B'' , we may assume that there is a unique point $y' \in \varphi^{-1}(V) \cap B'$. Let $y = \varphi(y') \in V$. Since $g(y') = 0$ we have $g(y) = 0$.

The open analytic domain $B' \setminus \{y'\}$ is the disjoint union of an open annulus A' and an infinite collection of open balls by Lemma 3.4. Each connected component C' of $B' \setminus \{y'\}$ maps into a connected component C of $X^{\text{an}} \setminus (V \cup D)$ adjacent to y , with the end y' of C' mapping to the end y of C . By Lemma 4.3, no open ball connected component of $B' \setminus \{y'\}$ can map to a generalized open annulus connected component of $X^{\text{an}} \setminus (V \cup D)$. There are at least two generalized open annulus connected components A of $X^{\text{an}} \setminus (V \cup D)$ adjacent to y , so some such A must satisfy $\varphi(B') \cap A = \emptyset$. But the map $\varphi : X'^{\text{an}} \rightarrow X^{\text{an}}$ is open by [Ber90, Lemma 3.2.4], so $\varphi(B')$ is an open neighborhood of y , which contradicts the fact that y is a limit point of A . This proves the claim.

Let V' be a semistable vertex set of (X', D') such that $\Sigma' = \Sigma(X', V' \cup D')$. Since $X' \setminus \Sigma'$ is a disjoint union of open balls, by the above we have $\varphi^{-1}(V) \subset \Sigma'$. Hence we may enlarge V' to contain $\varphi^{-1}(V)$ without changing Σ' , so we are reduced to (2). \blacksquare

Theorem 4.13. *Let Σ' be a skeleton of (X', D') and let Σ be a skeleton of (X, D) . Suppose that $\varphi(\Sigma') \subset \Sigma$, and if $X \cong \mathbf{P}^1$ assume in addition that there exists a type-2 point $z \in \Sigma$ such that $\varphi^{-1}(z) \subset \Sigma'$. Then $\varphi^{-1}(\Sigma)$ is a skeleton of $(X', \varphi^{-1}(D))$.*

We will require the following lemmas. Lemma 4.14 is similar to [Ber90, Corollary 4.5.4].

Lemma 4.14. *Let $B' \subset X'^{\text{an}}$ be an open analytic domain isomorphic to $\mathbf{B}(1)_+$ and let $B = \varphi(B') \subset X^{\text{an}}$. Then B is an open analytic domain of X^{an} , and one of the following is true:*

- (1) B is an open ball and $\varphi : B' \rightarrow B$ is finite and order-preserving.
- (2) $X \cong \mathbf{P}^1$ and $B = X^{\text{an}}$.

Proof. Suppose that the genus of X is at least one, so X' also has genus at least one. By Lemma 4.6 we only need to show that B is contained in an open ball in X^{an} . Let Σ (resp. Σ') be the minimal skeleton of X (resp. X'). By Proposition 3.18(1) we have $B' \subset X'^{\text{an}} \setminus \Sigma'$, so by Proposition 4.12(1) we have $B \subset X^{\text{an}} \setminus \Sigma$. But every connected component of $X^{\text{an}} \setminus \Sigma$ is an open ball, so B is contained in an open ball.

In the case $X = \mathbf{P}^1$, suppose first that $B'(K) \rightarrow \mathbf{P}^1(K)$ is not surjective, so we may assume that $\infty \notin B$ after choosing a suitable coordinate on \mathbf{P}^1 . Let x' be the end of B' and let $x = \varphi(x')$. Then x is a type-2 point, so $\varphi(B' \cup \{x'\}) = B \cup \{x\}$ is a compact subset of $\mathbf{A}^{1, \text{an}}$. Since $\mathbf{A}^{1, \text{an}}$ is covered by an increasing union of open balls, B is contained in an open ball.

Now suppose that $B'(K) \rightarrow \mathbf{P}^1(K)$ is surjective. Since $\mathbf{P}^1(K)$ is dense in $\mathbf{P}^{1, \text{an}}$ and $B \cup \{x\}$ is closed in $\mathbf{P}^{1, \text{an}}$, it follows that $B \cup \{x\} = \mathbf{P}^{1, \text{an}}$. If $x \notin B$, then B is contained in a connected component of $\mathbf{P}^{1, \text{an}} \setminus \{x\}$, which contradicts the surjectivity of $B'(K) \rightarrow \mathbf{P}^1(K)$. Therefore $B = \mathbf{P}^{1, \text{an}}$. \blacksquare

Remark 4.15. Case (2) of Lemma 4.14 does occur. For instance, let $\varphi : \mathbf{P}^1 \rightarrow \mathbf{P}^1$ be the finite morphism $t \mapsto t^2$, and let B' be the open ball in $\mathbf{P}^{1, \text{an}}$ obtained by deleting the closed ball of radius $1/2$ around $1 \in \mathbf{P}^1(K)$. If $\text{char}(k) \neq 2$, then for every point $x \in \mathbf{P}^1(K)$, either $x \in B'(K)$ or $-x \in B'(K)$, so $B' \rightarrow \mathbf{P}^{1, \text{an}}$ is surjective.

Lemma 4.16. *Let V be a semistable vertex set of (X, D) , let $\Sigma = \Sigma(X, V \cup D)$, and let $B \subset X$ be an open analytic domain isomorphic to $\mathbf{B}(1)_+$ whose end x is contained in Σ . Then*

$$\Sigma \cap \overline{B} = \bigcup_{y \in B \cap (V \cup D)} [x, y].$$

Proof. We may assume without loss of generality that $x \in V$. Let $V_0 = V \setminus (V \cap B)$ and $D_0 = D \setminus (D \cap B)$. Then V_0 is a semistable vertex set of (X, D_0) : indeed, every connected component of $X^{\text{an}} \setminus (V \cup D)$ is either a connected component of $X^{\text{an}} \setminus (V_0 \cup D_0)$ or is contained in B , so $X^{\text{an}} \setminus (V_0 \cup D_0)$ is a disjoint union of open balls and finitely many generalized open annuli. Let $\Sigma_0 = \Sigma(X, V_0 \cup D_0)$, so $x \in \Sigma_0$ but $B \cap \Sigma_0 = \emptyset$ by Proposition 3.12(1). By Lemma 4.6 we have that B is a connected component of $X^{\text{an}} \setminus \Sigma_0$, so the Lemma now follows from Lemma 3.15 as applied to $\Sigma = \Sigma_0$, $W = V \cap B$, and $E = D \cap B$. \blacksquare

Proof of Theorem 4.13. Note that $\varphi(\Sigma') \subset \Sigma$ implies $\varphi(D') \subset D$. Let V (resp. V') be a vertex set for Σ (resp. Σ'). We claim that

$$\varphi^{-1}(\Sigma) = \Sigma' \cup \bigcup_{x' \in \varphi^{-1}(V \cup D)} [x', \tau_{\Sigma'}(x')],$$

which by Lemma 3.15 is the minimal skeleton of $(X', \varphi^{-1}(D))$ containing Σ' and $\varphi^{-1}(V)$.

Let B' be a connected component of $X'^{\text{an}} \setminus \Sigma'$, let $x' \in \Sigma'$ be its end, and let $x = \varphi(x') \in \Sigma$. Then $B = \varphi(B')$ is an open ball and x is its end: if $X \not\cong \mathbf{P}^1$, then this follows directly from Lemma 4.14, and if $X \cong \mathbf{P}^1$, then $B \subset X^{\text{an}} \setminus \{z\}$, so $\varphi(B) \neq X^{\text{an}}$ and therefore B is an open ball in this case as well. Hence

$$\Sigma \cap B = \bigcup_{y \in B \cap (V \cup D)} [x, y]$$

by Lemma 4.16, so

$$\varphi^{-1}(\Sigma) \cap B' = \bigcup_{y' \in B' \cap \varphi^{-1}(V \cup D)} [x', y']$$

by Proposition 4.8. ■

Remark 4.17. When $X \cong \mathbf{P}^1$, the extra hypothesis on Σ' in Theorem 4.13 is necessary. Indeed, let $\varphi : \mathbf{P}^1 \rightarrow \mathbf{P}^1$ be a finite morphism and let $x' \in \mathbf{P}^{1, \text{an}}$ be a type-2 point such that $\varphi^{-1}(\varphi(x'))$ has more than one element. Then $\Sigma = \{\varphi(x')\}$ is a skeleton containing the image of the skeleton $\Sigma' = \{x'\}$, but $\varphi^{-1}(\Sigma)$ is not a skeleton.

Corollary 4.18. *Let Σ (resp. Σ') be a skeleton of (X, D) (resp. (X', D')). There exists a skeleton Σ_1 of $(X, D \cup \varphi(D'))$ such that $\Sigma_1 \supset \Sigma \cup \varphi(\Sigma')$ and such that $\varphi^{-1}(\Sigma_1)$ is a skeleton of $(X', \varphi^{-1}(D \cup \varphi(D')))$. Moreover, there is a minimal such Σ_1 with respect to inclusion.*

Proof. If $X \not\cong \mathbf{P}^1$ then this is an immediate consequence of Proposition 4.11 and Theorem 4.13: if Σ_1 is the minimal skeleton of $(X, D \cup \varphi(D'))$ containing $\Sigma \cup \varphi(\Sigma')$, then $\varphi^{-1}(\Sigma_1)$ is a skeleton of $(X', \varphi^{-1}(D \cup \varphi(D')))$. Suppose then that $X \cong \mathbf{P}^1$. Let $x' \in \Sigma'$ be a type-2 point, let $x = \varphi(x')$, and let $x', x'_1, x'_2, \dots, x'_n$ be the points of X' mapping to x . Let

$$\Sigma'_1 = \Sigma' \cup \bigcup_{i=1}^n [x'_i, \tau_{\Sigma'}(x'_i)].$$

This is the minimal skeleton of (X', D') containing Σ' and $\varphi^{-1}(x)$ by Lemma 3.15(3). Let Σ_1 be the minimal skeleton of $(X, D \cup \varphi(D'))$ containing $\Sigma \cup \varphi(\Sigma'_1)$. Then $\varphi^{-1}(\Sigma_1)$ is a skeleton of $(X', \varphi^{-1}(D \cup \varphi(D')))$ by Theorem 4.13. If Σ_2 is a skeleton of $(X, D \cup \varphi(D'))$ such that $\Sigma_2 \supset \Sigma \cup \varphi(\Sigma')$ and $\varphi^{-1}(\Sigma_2)$ is a skeleton of $(X', \varphi^{-1}(D \cup \varphi(D')))$, then $\varphi^{-1}(\Sigma_2)$ contains each geodesic $[x'_i, \tau_{\Sigma'}(x'_i)]$, so $\varphi^{-1}(\Sigma_2)$ contains Σ'_1 and therefore $\Sigma_2 \supset \Sigma_1$. ■

Remark 4.19. Suppose that $\varphi : X' \rightarrow X$ is a finite morphism such that $\varphi^{-1}(D) = D'$ and $\varphi^{-1}(V) = V'$. Let $\Sigma = \Sigma(X, V \cup D)$ and $\Sigma' = \Sigma(X', V' \cup D')$. In this case we can describe the skeleton Σ_1 of Corollary 4.18 more explicitly, as follows. Let e' be an open edge of Σ' with respect to the given choice of vertex set and let $A' = \tau^{-1}(e')$. This is a connected component of $X'^{\text{an}} \setminus (V' \cup D')$. Let A be the connected component of $X^{\text{an}} \setminus (V \cup D)$ containing $\varphi(A')$. Since A' is a connected component of $\varphi^{-1}(A)$, the map $A' \rightarrow A$ is finite, hence surjective. If A' is a punctured open ball, then so is A , so by Lemma 4.2, φ maps e' homeomorphically with nonzero integer expansion factor onto $e = \Sigma(A)$. If A' is an open annulus, then A could be an open annulus or an open ball; in the former case we again have e' mapping homeomorphically onto $e = \Sigma(A)$ with nonzero integer expansion factor. If A is an open ball, then by Proposition 4.9, e' maps onto a geodesic segment in A ; clearly both vertices of e' map to the end of A , so e' “goes straight and doubles back” under φ .

It follows from this and Proposition 4.12(2) that $\varphi(\Sigma')$ is equal to Σ union a finite number of geodesic segments T_1, \dots, T_r contained in open balls $B_1, \dots, B_r \subset X^{\text{an}} \setminus (V \cup D)$ and with one

endpoint at elements of V . These are precisely the images of the edges of Σ' not mapping to edges of Σ . By Lemma 3.15, $\Sigma \cup (T_1 \cup \dots \cup T_r)$ is again a skeleton of X , so in this case

$$\Sigma_1 = \varphi(\Sigma') = \Sigma \cup (T_1 \cup \dots \cup T_r)$$

(the inverse image of Σ_1 is a skeleton by Theorem 4.13).

4.20. Tangent directions and morphisms of curves. Our next goal is to prove that a finite morphism of curves induces a finite harmonic morphism of metrized complexes of curves for a suitable choice of skeleta. First we need to formulate and prove a relative version of Theorem 3.21. Let X, X' be smooth, proper, connected K -curves.

Definition 4.21. A continuous function $\varphi : X'^{\text{an}} \rightarrow X^{\text{an}}$ is *piecewise affine* provided that, for every geodesic segment $\alpha : [a, b] \hookrightarrow X'^{\text{an}}$, the composition $\varphi \circ \alpha$ is piecewise affine with integer expansion factors in the sense of Definition 4.4.

Let $\varphi : X'^{\text{an}} \rightarrow X^{\text{an}}$ be a piecewise affine function, let $x' \in X'^{\text{an}}$, and let $x = \varphi(x') \in X^{\text{an}}$. Let $v' \in T_{x'}$ and let $\alpha : [a, b] \hookrightarrow X'^{\text{an}}$ be a geodesic segment representing v' (so $a = -\infty$ if x' has type 1). Taking b small enough, we can assume that $\varphi \circ \alpha$ is an embedding with integer expansion factor m . Let $v \in T_x$ be the tangent direction represented by $\varphi \circ \alpha$. We define

$$d\varphi(x') : T_{x'} \rightarrow T_x \quad \text{by} \quad d\varphi(x')(v') = v;$$

this is independent of the choice of α . We call the expansion factor m the *outgoing slope of φ in the direction v'* and we write $m = d_{v'}\varphi(x')$.

Definition 4.22. A piecewise affine function $\varphi : X'^{\text{an}} \rightarrow X^{\text{an}}$ is *harmonic* at a point $x' \in X'^{\text{an}}$ provided that, for all $v \in T_{\varphi(x')}$, the integer

$$\sum_{\substack{v' \in T_{x'} \\ d\varphi(v')=v}} d_{v'}\varphi(x')$$

is independent of the choice of v .

Let $\varphi : X' \rightarrow X$ be a finite morphism of smooth, proper, connected K -curves, let $x' \in X'^{\text{an}}$ be a type-2 point, and let $x = \varphi(x')$. Let C_x and $C_{x'}$ denote the smooth proper connected k -curves with function fields $\mathcal{H}(x)$ and $\mathcal{H}(x')$, respectively. We denote by $\varphi_{x'}$ the induced morphism $C_{x'} \rightarrow C_x$. We have the following relative version of the slope formula of Theorem 3.21:

Theorem 4.23. *Let $\varphi : X' \rightarrow X$ be a finite morphism of smooth, proper, connected K -curves.*

- (1) *The analytification $\varphi : X'^{\text{an}} \rightarrow X^{\text{an}}$ is piecewise affine and harmonic.*
- (2) *Let $x' \in X'^{\text{an}}$ be a type-2 point, let $x = \varphi(x')$, let $v' \in T_{x'}$, let $v = d\varphi(v') \in T_x$, and let $\xi_{v'} \in C_{x'}$ and $\xi_v \in C_x$ be the closed points associated (3.22.1) to v' and v , respectively. Then $\varphi_{x'}(\xi_{v'}) = \xi_v$, and the ramification degree of $\varphi_{x'}$ at $\xi_{v'}$ is equal to $d_{v'}\varphi(x')$.*
- (3) *Let $x' \in X'^{\text{an}}$ be a type-1 point and let $v' \in T_{x'}$ be the unique tangent direction. Then $d_{v'}\varphi(x')$ is the ramification degree of φ at x' .*

Proof. First we claim that φ is piecewise affine. Let $[x', y'] \subset X'^{\text{an}}$ be a geodesic segment. Suppose first that x' and y' have type 2 or 3. Then there exists a skeleton Σ' of X' containing $[x', y']$ by [BPR11, Corollaries 5.56 and 5.64]. From this we easily reduce to the case that $[x', y']$ is an edge of Σ' with respect to some vertex set; now the claim follows from Lemma 4.2 and Proposition 4.9 as in the proof of Proposition 4.11. If x' has type 1 or 4, then there is an open neighborhood B of $x = \varphi(x')$ and an open neighborhood $B' \subset \varphi^{-1}(B)$ of x' such that B and B' are open balls. Shrinking B' if necessary we can assume that $y' \notin B'$, so the end z' of B' is contained in $[x', y']$ (removing z' disconnects X'^{an}). The restriction of φ to $[x', z']$ is piecewise affine by Lemma 4.5, so this proves the claim.

Next we prove (2). By functoriality of the reduction map, for any non-zero rational function f on X , we have $\varphi_{x'}^*(\tilde{f}_x) = \varphi^*(f)_{x'}$. By the slope formula (Theorem 3.21), the point ξ_v corresponds to the

discrete valuation on $\widetilde{\mathcal{H}}(x)$ given by $\text{ord}_{\xi_v}(\widetilde{f}_x) = d_v F(x)$, where $F = -\log |f|$. Similarly, the slope formula applied to $\varphi^*(f)$ on X' gives $\text{ord}_{\xi_{v'}}(\varphi^*(f)_{x'}) = d_{v'} \varphi(x') d_v F(x)$. Therefore, we get

$$\text{ord}_{\xi_{v'}}(\varphi_{x'}^*(\widetilde{f}_x)) = d_{v'} \varphi(x') \text{ord}_{\xi_v}(\widetilde{f}_x).$$

Since any non-zero element of $\widetilde{\mathcal{H}}(x)$ is of the form \widetilde{f}_x for some non-zero rational function f on X , this shows that the center of the valuation $\xi_{v'}$ on $\widetilde{\mathcal{H}}(x) \xrightarrow{\varphi_{x'}^*} \widetilde{\mathcal{H}}(x')$ coincides with the center of ξ_v , that is $\varphi_{x'}(\xi_{v'}) = \xi_v$, and at the same time, the ramification degree of $\varphi_{x'}$ at $\xi_{v'}$ is equal to $d_{v'} \varphi(x')$.

The proof of (3) proceeds in the same way as (2), applying the slope formula to a non-zero rational function f on X with a zero at $x = \varphi(x')$, and to $\varphi^*(f)$ on X' . ■

4.24. Morphisms of curves induce morphisms of metrized complexes. More precisely, this occurs when the morphism of curves respects a choice of triangulations, in the following sense.

Definition 4.25. Let $(X, V \cup D)$ and $(X', V' \cup D')$ be triangulated punctured curves. A *finite morphism* $\varphi : (X', V' \cup D') \rightarrow (X, V \cup D)$ is a finite morphism $\varphi : X' \rightarrow X$ such that $\varphi^{-1}(D) = D'$, $\varphi^{-1}(V) = V'$, and $\varphi^{-1}(\Sigma(X, V \cup D)) = \Sigma(X', V' \cup D')$ (as sets).

Corollary 4.26. Let $\varphi : X' \rightarrow X$ be a finite morphism of smooth, connected, proper K -curves, let $D \subset X(K)$ be a finite set, and let $D' = \varphi^{-1}(D)$. There exists a strongly semistable vertex set V of (X, D) such that $V' = \varphi^{-1}(V)$ is a strongly semistable vertex set of (X', D') and such that

$$\Sigma(X', V' \cup D') = \varphi^{-1}(\Sigma(X, V \cup D)).$$

In particular, φ extends to a finite morphism of triangulated punctured curves.

Proof. By Corollary 4.18, there is a skeleton Σ of (X, D) such that $\Sigma' = \varphi^{-1}(\Sigma)$ is a skeleton of (X', D') . Let V_0' be a strongly semistable vertex set of Σ' , let V be a strongly semistable vertex set of Σ containing $\varphi(V_0')$, and let $V' = \varphi^{-1}(V)$. Then V' is again a strongly semistable vertex set of Σ' . ■

4.27. Let $\varphi : (X', V' \cup D') \rightarrow (X, V \cup D)$ be a finite morphism of triangulated punctured curves. Let $\Sigma = \Sigma(X, V \cup D)$ and $\Sigma' = \Sigma(X', V' \cup D')$. Let e' be an open edge of Σ' and let $e = \varphi(e')$, an open edge of Σ . The image of the annulus $A' = \tau^{-1}(e')$ is contained in $A = \tau^{-1}(e)$, so the restriction of φ to e' is an embedding with integer expansion factor by Lemma 4.2(2). It is clear from this and Theorem 4.23 that $\varphi|_{\Sigma'} : \Sigma' \rightarrow \Sigma$ is a harmonic (V', V) -morphism of Λ -metric graphs. For $x' \in V'$ with image $x = \varphi(x')$ we have a finite morphism of k -curves $\varphi_{x'} : C_{x'} \rightarrow C_x$. It now follows from Theorem 4.23 that these extra data enrich $\varphi|_{\Sigma'} : \Sigma' \rightarrow \Sigma$ with the structure of a finite harmonic morphism of Λ -metrized complexes of curves:

Corollary 4.28. Let $\varphi : (X', V' \cup D') \rightarrow (X, V \cup D)$ be a finite morphism of triangulated punctured curves. Then φ naturally induces a finite harmonic morphism of Λ -metrized complexes of curves

$$\Sigma(X', V' \cup D') \longrightarrow \Sigma(X, V \cup D).$$

Remark 4.29. More generally, suppose that $(X, V \cup D)$ and $(X', V' \cup D')$ are triangulated punctured K -curves and that $\varphi : X' \rightarrow X$ is a finite morphism such that $\varphi^{-1}(D) = D'$ and $\varphi^{-1}(V) \subset V'$. Then $\varphi^{-1}(\Sigma) \subset \Sigma'$ by Proposition 4.12(2). One can show using Theorem 4.23 that the map $\Sigma' \rightarrow X^{\text{an}}$ followed by the retraction $X^{\text{an}} \rightarrow \Sigma$ is a (not necessarily finite) harmonic morphism of metric graphs $\Sigma' \rightarrow \Sigma$.

4.30. Tame coverings of triangulated curves. Many of the results in this paper involve constructing a curve X' and a morphism $\varphi : X' \rightarrow X$ inducing a given morphism of skeleta as in Corollary 4.28. For these purposes it is useful to introduce some mild restrictions on the morphism φ . Fix a triangulated punctured curve $(X, V \cup D)$ with skeleton $\Sigma = \Sigma(X, V \cup D)$, regarded as a metrized complex of curves as in (3.22).

Definition 4.31. Let $(X', V' \cup D')$ be a triangulated punctured curve with skeleton $\Sigma' = \Sigma(X', V' \cup D')$ and let $\varphi : (X', V' \cup D') \rightarrow (X, V \cup D)$ be a finite morphism. We say that φ is a *tame covering* of $(X, V \cup D)$ provided that:

- (1) D contains the branch locus of φ ,
- (2) if $\text{char}(k) = p > 0$, then for every edge $e' \in E(\Sigma')$ the expansion factor $d_{e'}(\varphi)$ is not divisible by p , and
- (3) $\varphi_{x'}$ is separable (= generically étale) for all $x' \in V'$.

Remark 4.32.

- (1) Since D is finite, it follows from (1) and Theorem 4.23(3) that a tame covering of $(X, V \cup D)$ is a tamely ramified morphism of curves.
- (2) If Σ has at least one edge, then (2) implies (3).
- (3) Let $\varphi : (X', V' \cup D') \rightarrow (X, V \cup D)$ be a tame covering, let $S \subset \Sigma$ be a finite set of type-2 points, and let $S' = \varphi^{-1}(S)$. Then $S' \subset \Sigma(X', V' \cup D')$ is also a finite set of type-2 points, so φ also defines a tame covering $(X', S' \cup V' \cup D') \rightarrow (X, S \cup V \cup D)$.

Lemma 4.33. *Let $\varphi : (X', V' \cup D') \rightarrow (X, V \cup D)$ be a tame covering and let $\Sigma = \Sigma(X, V \cup D)$ and $\Sigma' = \Sigma(X', V' \cup D')$. Then $\varphi|_{\Sigma'} : \Sigma' \rightarrow \Sigma$ is a tame covering of metrized complexes of curves.*

Proof. Let Γ (resp. Γ') be the augmented metric graph underlying Σ (resp. Σ'). By Proposition 2.22 we only have to show that $\varphi|_{\Gamma'}$ is generically étale. Let $R = \sum_{x' \in V' \cup D'} R_{x'}(x')$ be the ramification divisor of $\varphi|_{\Gamma'}$, as defined in (2.14.1), and let $S = \sum_{x' \in D'} S_{x'}(x')$ be the ramification divisor of $\varphi : X' \rightarrow X$. We will show that $R = S$. Since $\varphi_{x'}$ is generically étale for all $x' \in V'$, we see from (2.14.1) and the Riemann–Hurwitz formula as applied to $\varphi_{x'}$ that $R_{x'} \geq 0$. If $x' \in D'$, then there is a unique edge e' adjacent to x' and $d_{e'}(\varphi) = d_{x'}(\varphi)$, so

$$R_{x'} = 2d_{x'}(\varphi) - 2 - d_{x'}(\varphi) + 1 = d_{x'}(\varphi) - 1 = S_{x'},$$

where the final equality is Theorem 4.23(3). Since $R_{x'} \geq 0$ for all $x' \in V'$, it is enough to show that $\deg(R) = \deg(S)$. By (2.14.2) we have $K_{\Gamma'} = (\varphi|_{\Gamma'})^*(K_{\Gamma}) + R$, so counting degrees gives

$$\deg(R) = \deg(\varphi|_{\Gamma'}) (2 - 2g(\Gamma)) - (2 - 2g(\Gamma')).$$

The Lemma now follows from the equalities $\deg(\varphi) = \deg(\varphi|_{\Gamma'})$, $g(\Gamma) = g(X)$, $g(\Gamma') = g(X')$, and the Riemann–Hurwitz formula as applied to $\varphi : X' \rightarrow X$. \blacksquare

Remark 4.34. We showed in the proof of Lemma 4.33 that the ramification divisor of $\varphi|_{\Gamma'}$ coincides with the ramification divisor of φ . Moreover, it follows from Proposition 2.22 that for every $x' \in V'$, every ramification point $\bar{x}' \in C_{x'}$ is the reduction of a tangent direction in Γ' . In other words, for a tame covering all ramification points of all residue curves are “visible” in the morphism of underlying metric graphs $\Gamma' \rightarrow \Gamma$.

Proposition 4.35. *Let $\varphi : (X', V' \cup D') \rightarrow (X, V \cup D)$ be a finite morphism of triangulated punctured curves. Then φ is a tame covering if and only if*

- (1) D contains the branch locus of φ and
- (2) $\varphi_{x'} : C_{x'} \rightarrow C_{\varphi(x)}$ is tamely ramified for every type-2 point $x' \in X'^{\text{an}}$.

Moreover, if φ is a tame covering and $x' \in X'^{\text{an}}$ is a type-2 point not contained in Σ' , then φ is an isomorphism in a neighborhood of x' .

Proof. It is clear that conditions (1)–(2) imply that φ is a tame covering, so suppose that φ is a tame covering and $x' \in X'^{\text{an}}$ is a type-2 point. We have the following cases:

- If $x' \in V'$, then $\varphi_{x'}$ is tamely ramified by Lemma 4.33 and Proposition 2.22.
- Suppose that x' is contained in the interior of an edge e' of Σ' . Since φ is a tame covering of $(X, (V \cup \{\varphi(y')\}) \cup D)$ by Remark 4.32(3), we are reduced to the case treated above.
- Suppose that $x' \notin \Sigma'$ but $y' = \tau(x')$ is contained in V' . Let \mathfrak{X} (resp. \mathfrak{X}') be the semistable formal model of X (resp. X') corresponding to the semistable vertex set V (resp. V') and let

φ denote the unique finite morphism $\mathfrak{X}' \rightarrow \mathfrak{X}$ extending $\varphi : X' \rightarrow X$ (see Theorem 5.13). Let $\bar{x}' = \text{red}(x') \in \mathfrak{X}'(k)$ and let $x = \varphi(x')$ and $\bar{x} = \text{red}(x) = \varphi_k(\bar{x}')$. The connected component B' of x' in $X'^{\text{an}} \setminus \Sigma'$ is equal to $\text{red}^{-1}(\bar{x}')$, and the connected component B of x in $X^{\text{an}} \setminus \Sigma$ is equal to $\text{red}^{-1}(\bar{x})$. Since $\Sigma \cap B = \emptyset$, φ_k is not branched over \bar{x} by Lemma 4.33 and Proposition 2.22. Therefore $\varphi^{-1}(B)$ is a disjoint union of $\deg(\varphi)$ open balls (one of which is B') mapping isomorphically onto B . In particular, φ is an isomorphism in a neighborhood of x' .

- Suppose that $x' \notin \Sigma'$ but $y' = \tau(x')$ is contained in the interior of an edge of Σ' . We claim that φ is an isomorphism in a neighborhood of x' . Let $y = \varphi(y') \in \Sigma$. Then φ is a tame covering of $(X, (V \cup \{y\}) \cup D)$, so replacing V with $V \cup \{y\}$ and V' with $V' \cup \varphi^{-1}(y)$ (see Remark 4.32(3)), we are reduced to the previous case. ■

Remark 4.36. It follows from Proposition 4.35 that if φ is a tame covering then the set-theoretic branch locus of $\varphi : X'^{\text{an}} \rightarrow X^{\text{an}}$ (i.e. the set of all points in X^{an} with fewer than $\deg(\varphi)$ points in its fiber) is contained in Σ . See [Fab13a, Fab13b] for more on the topic of the Berkovich ramification locus in the case of self-maps of \mathbf{P}^1 .

5. APPLICATIONS TO MORPHISMS OF SEMISTABLE MODELS

In this section we show how Section 4 formally implies a large part of the results of Liu [Liu06] on simultaneous semistable reduction of morphisms of curves (see also [Col03]) over an arbitrary valued field. In addition to establishing these results over a more general ground field, we feel that the “skeletal” point of view on simultaneous semistable reduction is enlightening (see Remark 5.23). To this end, we let K_0 be any field equipped with a nontrivial non-Archimedean valuation $\text{val} : K_0 \rightarrow \mathbf{R} \cup \{\infty\}$. Its valuation ring will be denoted R_0 , its maximal ideal \mathfrak{m}_{R_0} , and its residue field k_0 .

Let X be a smooth, proper, geometrically connected algebraic K_0 -curve. By a (strongly) semistable R_0 -model of X we mean a flat, integral, proper relative curve $\mathcal{X} \rightarrow \text{Spec}(R_0)$ whose special fiber \mathcal{X}_{k_0} is a (strongly) semistable curve (i.e. \mathcal{X}_{k_0} is a reduced curve with at worst ordinary double point singularities; it is strongly semistable if its irreducible components are smooth) and whose generic fiber is equipped with an isomorphism $\mathcal{X}_{K_0} \cong X$. By properness of \mathcal{X} , any K_0 -point $x \in X(K_0)$ extends in a unique way to an R_0 -point $x \in \mathcal{X}(R_0)$; the special fiber of this point is the *reduction* of x and is denoted $\text{red}(x) \in \mathcal{X}(k_0)$. Let $D \subset X(K_0)$ be a finite set. A semistable model \mathcal{X} of X is a *semistable R_0 -model of (X, D)* provided that the points of D reduce to distinct smooth points of \mathcal{X}_{k_0} . The model \mathcal{X} is a *stable R_0 -model of (X, D)* provided that every rational (resp. genus-1) component of the normalization \mathcal{X}_k contains at least three points (resp. one point) mapping to a singular point of \mathcal{X}_k or to the reduction of a point of D .

If $K_0 = K$ is complete and algebraically closed, we define a (strongly) semistable formal R -model of X to be an integral, proper, admissible formal R -curve \mathfrak{X} whose analytic generic fiber \mathfrak{X}_K is equipped with an isomorphism to X^{an} and whose special fiber \mathfrak{X}_k is a (strongly) semistable curve. There is a natural map of sets $\text{red} : X^{\text{an}} \rightarrow \mathfrak{X}_k$, called the *reduction*; it is surjective and anti-continuous in the sense that the inverse image of a Zariski-open subset of \mathfrak{X}_k is a closed subset of X^{an} (and vice-versa). Using the reduction map, for a finite set $D \subset X(K)$ we define *semistable* and *stable formal R -models of (X, D)* as above.

As we will be passing between formal and algebraic R -models of K -curves, it is worth stating the following lemma; see [BPR11, Remark 5.30(2)].

Lemma 5.1. *Let X be a smooth, proper, connected K -curve. The ϖ -adic completion functor defines an equivalence from the category of semistable R -models of X to semistable formal R -models of X .*

The inverse of the ϖ -adic completion functor of Lemma 5.1 will be called *algebraization*.

5.2. Descent to a general ground field. We will use the following lemmas to descend the geometric theory of Section 4 to the valued field K_0 . Fix an algebraic closure \bar{K}_0 of K_0 and a valuation val on

\overline{K}_0 extending the given valuation on K_0 . For any field $K_1 \subset \overline{K}_0$ we consider K_1 as a valued field with respect to the restriction of val , and we write R_1 for the valuation ring of K_1 . Let K be the completion of \overline{K}_0 with respect to val . This field is algebraically closed by [BGR84, Proposition 3.4.1/3].

Lemma 5.3. *Let X_0 be a smooth, proper, geometrically connected curve over K_0 and let $X = X_0 \otimes_{K_0} K$. Let K_0^{sep} be the separable closure of K_0 in K . Then the image of $X_0(K_0^{\text{sep}})$ under the natural inclusion*

$$X_0(K_0^{\text{sep}}) \subset X_0(K) = X(K) \subset X^{\text{an}}$$

is dense in X^{an} .

Proof. Let $\widehat{K}_0 \subset K$ denote the completion of K_0 and let $(\widehat{K}_0)^{\text{sep}} \subset K$ be its separable closure. By [BGR84, Proposition 3.4.1/6], $(\widehat{K}_0)^{\text{sep}}$ is dense in K . Let $\widehat{K}_1 \subset (\widehat{K}_0)^{\text{sep}}$ be a finite, separable extension of \widehat{K}_0 . By Krasner's lemma (see §3.4.2 of loc. cit.), there exists a finite, separable extension K_1/K_0 contained in K_0^{sep} which is dense in \widehat{K}_1 . It follows that K_0^{sep} is dense in $(\widehat{K}_0)^{\text{sep}}$, so K_0^{sep} is dense in K . Since $\mathbf{P}_K^1(K)$ is dense in $\mathbf{P}_K^{1,\text{an}}$ and the subspace topology on $\mathbf{P}_K^1(K) \subset \mathbf{P}_K^{1,\text{an}}$ coincides with the ultrametric topology, this proves the lemma for $X_0 = \mathbf{P}_{K_0}^1$.

For general X_0 , choose a finite, generically étale morphism $\varphi : X_0 \rightarrow \mathbf{P}_{K_0}^1$. Let $D \subset \mathbf{P}_K^1(K)$ be the branch locus, so $\varphi : X^{\text{an}} \setminus \varphi^{-1}(D) \rightarrow \mathbf{P}_K^{1,\text{an}} \setminus D$ is étale, hence open by [Ber93, Corollary 3.7.4]. It follows that if $U \subset X^{\text{an}}$ is a nonempty open set, then $\varphi(U) \setminus D$ contains a nonempty open subset of $\mathbf{P}_K^{1,\text{an}} \setminus D$. Let $x \in \mathbf{P}_{K_0}^1(K_0^{\text{sep}})$ be a point contained in $\varphi(U) \setminus D$. Then $\varphi^{-1}(x) \subset X_0(K_0^{\text{sep}})$ and $\varphi^{-1}(x) \cap U \neq \emptyset$. ■

Lemma 5.4. *Let X_0 be a smooth, proper, geometrically connected curve over K_0 , let $D \subset X_0(K_0)$ be a finite subset, let $X = X_0 \otimes_{K_0} K$, and let \mathcal{X} be a semistable model of (X, D) . There exists a finite, separable extension K_1 of K_0 and a semistable model \mathcal{X}_1 of $X_1 = X_0 \otimes_{K_0} K_1$ with respect to D such that $\mathcal{X}_1 \otimes_{R_1} R \cong \mathcal{X}$.*

Proof. Let \mathfrak{X} be the ϖ -adic completion of \mathcal{X} . For $\bar{x} \in \mathfrak{X}_k(k)$ the subset $\text{red}^{-1}(\bar{x}) \subset X^{\text{an}}$ is open in X^{an} . By Lemma 5.3, there exists a point $x \in X_0(K_0^{\text{sep}})$ with $\text{red}(x) = \bar{x}$, so after passing to a finite, separable extension of K_0 if necessary, we may enlarge $D \subset X_0(K_0)$ to assume that \mathcal{X} is a stable model of (X, D) . On the other hand, by the stable reduction theorem (i.e. the properness of the Deligne-Mumford stack $\overline{\mathcal{M}}_{g,n}$ parameterizing stable marked curves) there exists a stable model \mathcal{X}_0 of (X_0, D_0) after potentially passing to a finite, separable extension of K_0 . By uniqueness of stable models we have $\mathcal{X}_0 \otimes_{R_0} R \cong \mathcal{X}$. ■

Lemma 5.5. *Let X_0, X'_0 be smooth, proper, geometrically connected K_0 -curves and let $\varphi : X'_0 \rightarrow X_0$ be a finite morphism. Let \mathcal{X}_0 (resp. \mathcal{X}'_0) be a semistable model of X_0 (resp. X'_0), let $X = X_0 \otimes_{K_0} K$ and $\mathcal{X} = \mathcal{X}_0 \otimes_{R_0} R$ (resp. $X' = X'_0 \otimes_{K_0} K$ and $\mathcal{X}' = \mathcal{X}'_0 \otimes_{R_0} R$), and suppose that $\varphi_K : X' \rightarrow X$ extends to a morphism $\mathcal{X}' \rightarrow \mathcal{X}$. Then φ extends to a morphism $\mathcal{X}'_0 \rightarrow \mathcal{X}_0$ in a unique way.*

Proof. The morphism φ gives rise to a section $X'_0 \rightarrow X_0 \times_{K_0} X'_0$ of $\text{pr}_2 : X_0 \times_{K_0} X'_0 \rightarrow X'_0$. Let \mathcal{Z} be the schematic closure of the image of X'_0 in $\mathcal{X}_0 \times_{R_0} \mathcal{X}'_0$. It is enough to show that $\text{pr}_2 : \mathcal{Z} \rightarrow \mathcal{X}'_0$ is an isomorphism. Since schematic closure respects flat base change, the schematic closure of the image of X' in $\mathcal{X} \times_R \mathcal{X}'$ is equal to $\mathcal{Z} \otimes_{R_0} R$. By properness of \mathcal{X}' , the image of \mathcal{X}' in $\mathcal{X} \times_R \mathcal{X}'$ is then equal to $\mathcal{Z} \otimes_{R_0} R$, so $\text{pr}_2 : \mathcal{Z} \otimes_{R_0} R \rightarrow \mathcal{X}'$ is an isomorphism. Since $R_0 \rightarrow R$ is faithfully flat, this implies that $\text{pr}_2 : \mathcal{Z} \rightarrow \mathcal{X}'_0$ is an isomorphism. ■

5.6. The relation between semistable models and skeleta. Let X be a smooth, proper, connected K -curve. The following theorem is due to Berkovich and Bosch-Lütkebohmert; see [BPR11, Theorem 5.34] for more precise references.

Theorem 5.7. *Let \mathfrak{X} be a semistable formal model of X and let $\bar{x} \in \mathfrak{X}_k$ be a point. Then*

- (1) \bar{x} is a generic point if and only if $\text{red}^{-1}(\bar{x})$ consists of a single type-2 point of X^{an} .
- (2) \bar{x} is a smooth closed point if and only if $\text{red}^{-1}(\bar{x})$ is an open ball.
- (3) \bar{x} is an ordinary double point if and only if $\text{red}^{-1}(\bar{x})$ is an open annulus.

It follows from Theorem 5.7 that if \mathfrak{X} is a semistable formal model of X , then

$$V(\mathfrak{X}) = \{ \text{red}^{-1}(\bar{\zeta}) : \zeta \in \mathfrak{X}_k \text{ is a generic point} \}$$

is a semistable vertex set of X : indeed,

$$(5.7.1) \quad X \setminus V(\mathfrak{X}) = \coprod_{\substack{\bar{x} \in \mathfrak{X}_k \\ \text{singular}}} \text{red}^{-1}(\bar{x}) \cup \coprod_{\substack{\bar{x} \in \mathfrak{X}_k \\ \text{smooth}}} \text{red}^{-1}(\bar{x})$$

is a disjoint union of open balls and finitely many open annuli. The following folklore theorem is proved in [BPR11, Theorem 5.38].

Theorem 5.8. *Let X be a smooth, proper, connected algebraic K -curve and let $D \subset X(K)$ be a finite set of closed points. The association $\mathfrak{X} \mapsto V(\mathfrak{X})$ sets up a bijection between the set of (strongly) semistable formal models of (X, D) (up to isomorphism) and the set of (strongly) semistable vertex sets of (X, D) .*

5.9. Let \mathfrak{X} be a semistable formal model of (X, D) . One can construct the metrized complex of curves $\Sigma(X, V(\mathfrak{X}) \cup D)$ directly from \mathfrak{X} , as follows. Let V be the set of irreducible components of \mathfrak{X}_k , regarded as a set of vertices of a graph. For $x \in V$ we will write C_x to denote the corresponding component of \mathfrak{X}_k . To every double point $\bar{x} \in \mathfrak{X}_k$ we associate an edge e connecting the irreducible components of \mathfrak{X}_k containing \bar{x} (this is a loop edge if there is only one such component); if C_x is such a component, then we set $\text{red}_x(e) = \bar{x}$. The completed local ring of \mathfrak{X} at \bar{x} is isomorphic to $R[[s, t]]/(st - \pi)$ for some $\pi \in \mathfrak{m}_R \setminus \{0\}$; we define the length of e to be $\ell(e) = \text{val}(\pi)$. We connect a point $x \in D$ to the vertex corresponding to the irreducible component of \mathfrak{X}_k containing $\text{red}(x)$. These data define a metrized complex of curves $\Sigma(\mathfrak{X}, D)$. By (5.7.1), the graphs underlying $\Sigma(\mathfrak{X}, D)$ and $\Sigma(X, V(\mathfrak{X}) \cup D)$ are naturally isomorphic. The edge lengths coincide — see for instance [BPR11, Proposition 5.37] — and the residue curves of the two metrized complexes are naturally isomorphic by [Ber90, Proposition 2.4.4]. From this it is straightforward to verify that $\Sigma(\mathfrak{X}, D)$ and $\Sigma(X, V(\mathfrak{X}) \cup D)$ are identified as metrized complexes of curves.

In particular, a semistable formal model \mathfrak{X} of (X, D) is stable if and only if $V(\mathfrak{X})$ is a stable vertex set of (X, D) .

As another consequence of Theorem 5.8, we obtain the following compatibility of skeleta and extension of scalars.

Proposition 5.10. *Let X be a smooth, proper, connected K -curve, let K' be a complete and algebraically closed valued field extension of K , and let $\pi : X_{K'}^{\text{an}} \rightarrow X_K^{\text{an}}$ be the canonical map. If \mathfrak{X} is a semistable formal model of X , then*

- (1) π maps $V(\mathfrak{X}_{K'})$ bijectively onto $V(\mathfrak{X})$, and
- (2) π maps $\Sigma(X_{K'}, V(\mathfrak{X}_{K'}) \cup D)$ bijectively onto $\Sigma(X, V(\mathfrak{X}) \cup D)$, with

$$\pi : \Sigma(X_{K'}, V(\mathfrak{X}_{K'}) \cup D) \xrightarrow{\sim} \Sigma(X, V(\mathfrak{X}) \cup D)$$

an isomorphism of augmented metric graphs.

Proof. The reduction map is compatible with extension of scalars, in that the following square commutes

$$\begin{array}{ccc} X_{K'}^{\text{an}} & \xrightarrow{\text{red}} & \mathfrak{X}_{k'} \\ \pi \downarrow & & \downarrow \bar{\pi} \\ X^{\text{an}} & \xrightarrow{\text{red}} & \mathfrak{X}_k \end{array}$$

with k' the residue field of K' . The first assertion is immediate because the extension of scalars morphism $\bar{\pi} : \mathfrak{X}_{k'} \rightarrow \mathfrak{X}_k$ is a bijection on generic points. It is also a bijection on singular points, and for any node $\bar{x}' \in \mathfrak{X}_{k'}$ the open annulus $A' = \text{red}^{-1}(\bar{x}')$ is identified with the extension of scalars of $A = \text{red}^{-1}(\bar{\pi}(\bar{x}'))$. It is easy to see that $\pi : A' \rightarrow A$ takes the skeleton of A' isomorphically onto the skeleton of A . \blacksquare

5.11. Extending morphisms to semistable models: analytic criteria. Let X be a smooth, proper, connected K -curve and let \mathfrak{X} be a semistable formal model of X . The *inverse image topology* on X^{an} is the topology $\mathcal{T}(\mathfrak{X})$ whose open sets are the sets of the form $\text{red}^{-1}(\mathfrak{U}_k)$, where \mathfrak{U}_k is a Zariski-open subset of \mathfrak{X}_k . Note that any such set is *closed* in the natural topology on X^{an} . If $\mathfrak{U}_k \subset \mathfrak{X}_k$ is an affine open subset, then \mathfrak{U}_k is the special fiber of a formal affine open $\mathfrak{U} = \text{Spf}(\mathcal{A}) \subset \mathfrak{X}$, the set underlying the generic fiber $\mathfrak{U}_K = \mathcal{M}(\mathcal{A}_K) \subset X^{\text{an}}$ is equal to $\text{red}^{-1}(\mathfrak{U}_k)$, and \mathcal{A} is equal to the ring $\hat{\mathcal{A}}_K$ of power-bounded elements in the affinoid algebra \mathcal{A}_K . This essentially means that \mathfrak{X} is a *formal analytic variety* in the sense of [BL85]; see [BPR11, Remark 5.30(3)] for an explanation. Therefore \mathfrak{X} is constructed by gluing the canonical models of the affinoid subdomains of X^{an} which are open in the inverse image topology $\mathcal{T}(\mathfrak{X})$, along the canonical models of their intersections; here the *canonical model* of an affinoid space $\mathcal{M}(\mathcal{B})$ is by definition $\text{Spf}(\hat{\mathcal{B}})$. The following general fact about formal analytic varieties follows from these observations and the functoriality of the reduction map.

Proposition 5.12. *Let X, X' be smooth, proper, connected algebraic curves over K , let $\varphi : X' \rightarrow X$ be a finite morphism, and let \mathfrak{X} and \mathfrak{X}' be semistable formal models of X and X' , respectively. Then φ extends to a morphism $\mathfrak{X}' \rightarrow \mathfrak{X}$ if and only if φ is continuous with respect to the inverse image topologies $\mathcal{T}(\mathfrak{X})$ and $\mathcal{T}(\mathfrak{X}')$, in which case there is exactly one morphism $\mathfrak{X}' \rightarrow \mathfrak{X}$ extending φ .*

Fix smooth, proper, connected algebraic K -curves X, X' and a finite morphism $\varphi : X' \rightarrow X$. The following theorem is well-known to experts, although no proof appears in the literature to the best of our knowledge.

Theorem 5.13. *Let \mathfrak{X} and \mathfrak{X}' be semistable formal models of X and X' , respectively. Then $\varphi : X' \rightarrow X$ extends to a morphism $\mathfrak{X}' \rightarrow \mathfrak{X}$ if and only if $\varphi^{-1}(V(\mathfrak{X})) \subset V(\mathfrak{X}')$, and $\mathfrak{X}' \rightarrow \mathfrak{X}$ is finite if and only if $\varphi^{-1}(V(\mathfrak{X})) = V(\mathfrak{X}')$.*

Remark 5.14. If \mathfrak{X} and \mathfrak{X}' are semistable formal models of the punctured curves (X, D) and (X', D') , respectively, and $\varphi : X' \rightarrow X$ is a finite morphism with $\varphi^{-1}(D) = D'$ which extends to a morphism $\mathfrak{X}' \rightarrow \mathfrak{X}$, then since $\varphi^{-1}(V(\mathfrak{X})) \subset V(\mathfrak{X}')$ it follows from Remark 4.29 that there is a natural harmonic morphism of metric graphs $\Sigma(X', V(\mathfrak{X}') \cup D') \rightarrow \Sigma(X, V(\mathfrak{X}) \cup D)$. The morphism $\mathfrak{X}' \rightarrow \mathfrak{X}$ is finite if and only if the local degree of this harmonic morphism at every $v' \in V(\mathfrak{X}')$ is positive.

Before giving a proof of Theorem 5.13, we mention the following consequences. Let \mathfrak{X}_1 and \mathfrak{X}_2 be semistable formal models of X . We say that \mathfrak{X}_1 *dominates* \mathfrak{X}_2 provided that there exists a (necessarily unique) morphism $\mathfrak{X}_1 \rightarrow \mathfrak{X}_2$ inducing the identity map on analytic generic fibers. Taking $X = X'$ in Theorem 5.13, we obtain the following corollary; this gives a different proof of the second part of [BPR11, Theorem 5.38].

Corollary 5.15. *Let \mathfrak{X}_1 and \mathfrak{X}_2 be semistable formal models of X . Then \mathfrak{X}_1 dominates \mathfrak{X}_2 if and only if $V(\mathfrak{X}_1) \supset V(\mathfrak{X}_2)$.*

In conjunction with Proposition 5.10, we obtain the following corollary. If K' is a complete and algebraically closed field extension of K , we say that a semistable formal model \mathfrak{X}' of $X_{K'}$ is *defined over R* provided that it arises as the extension of scalars of a (necessarily unique) semistable formal model of X .

Corollary 5.16. *Let K' be a complete and algebraically closed field extension of K , let \mathfrak{X}' be a semistable formal model of $X_{K'}$, and suppose that there exists a semistable formal model \mathfrak{X} of X such that $\mathfrak{X}_{R'}$ dominates \mathfrak{X}' . Then \mathfrak{X}' is defined over R .*

Proof. Let $\pi : X_{K'}^{\text{an}} \rightarrow X^{\text{an}}$ be the canonical map. By Corollary 5.15 we have $V(\mathfrak{X}_{R'}) \supset V(\mathfrak{X}')$, and by Proposition 5.10 the map π defines a bijection $V(\mathfrak{X}_{R'}) \xrightarrow{\sim} V(\mathfrak{X})$ and an isomorphism

$$\Sigma(X_{K'}, V(\mathfrak{X}_{R'})) \xrightarrow{\sim} \Sigma(X, V(\mathfrak{X})).$$

Let $V = V(\mathfrak{X})$ and $V' = \pi(V(\mathfrak{X}'))$. We claim that V' is a semistable vertex set. Granted this, letting \mathfrak{Y} be the semistable formal model of X associated to V' , we have $V(\mathfrak{Y}_{R'}) = V(\mathfrak{X}')$ since $V(\mathfrak{Y}_{R'}) \subset V(\mathfrak{X}_{R'})$ (as $\mathfrak{X}_{R'}$ dominates $\mathfrak{Y}_{R'}$) and $\pi(V(\mathfrak{Y}_{R'})) = V(\mathfrak{Y})$, so $\mathfrak{Y}_{R'} = \mathfrak{X}'$.

It remains to prove the claim that V' is a semistable vertex set. The point is that the question of whether or not a subset of V is a semistable vertex set is intrinsic to the augmented Λ -metric graph $\Sigma = \Sigma(X, V) \cong \Sigma(X_{K'}, V(\mathfrak{X}_{R'}))$. We leave the details to the reader. \blacksquare

We will use the following lemmas in the proof of Theorem 5.13. Recall from (3.11) that if V is a semistable vertex set of X , then a connected component C of $X^{\text{an}} \setminus V$ is *adjacent* to a point $x \in V$ provided that the closure of C in X^{an} contains x .

Lemma 5.17. *Let \mathfrak{X} be a semistable formal model of X , let $V = V(\mathfrak{X})$, let $\mathfrak{U}_k \subset \mathfrak{X}_k$ be a subset, and let $U = \text{red}^{-1}(\mathfrak{U}_k) \subset X^{\text{an}}$. Then U is open in the topology $\mathcal{T}(\mathfrak{X})$ if and only if the following conditions hold:*

- (1) U is closed in the ordinary topology on X^{an} , and
- (2) for every $x \in U \cap V$, all but finitely many connected components of $X^{\text{an}} \setminus V$ which are adjacent to x are contained in U .

Proof. Let $\bar{\zeta}$ be a generic point of \mathfrak{X}_k and let $\bar{y} \in \mathfrak{X}_k$ be a closed point. Let $x \in V$ be the unique point of X^{an} reducing to $\bar{\zeta}$ and let $B = \text{red}^{-1}(\bar{y})$. By the anti-continuity of the reduction map, \bar{y} is in the closure of $\{\bar{\zeta}\}$ if and only if x is adjacent to B . The lemma follows easily from this and the fact that the connected components of $X^{\text{an}} \setminus V$ are exactly the inverse images of the closed points of \mathfrak{X}_k under red . \blacksquare

Lemma 5.18. *Let V and V' be semistable vertex sets of X and X' , respectively, and suppose that $\varphi^{-1}(V) \subset V'$. Let C' be a connected component of $X'^{\text{an}} \setminus \varphi^{-1}(V)$. Then C' has the following form:*

- (1) If C' intersects $\Sigma(X', V')$, then $C' = \tau^{-1}(\Sigma(X', V') \cap C')$, and
- (2) otherwise C' is an open ball connected component of $X'^{\text{an}} \setminus V'$.

Proof. Let $\Sigma' = \Sigma(X', V')$. Suppose that there exists $y' \in C'$ such that $\tau(y') \notin C'$. Let B' be the connected component of $X'^{\text{an}} \setminus \Sigma'$ containing y' , so B' is an open ball contained in C' and $\tau(y')$ is the end of B' . It follows that $\tau(y')$ is contained in the closure of C' in X'^{an} . Since C' is a connected component of $X'^{\text{an}} \setminus \varphi^{-1}(V)$, its closure is contained in $C' \cup \varphi^{-1}(V)$, so $\tau(y') \in \varphi^{-1}(V)$. Therefore $B' = (B' \cup \{\tau(y')\}) \cap (X'^{\text{an}} \setminus \varphi^{-1}(V))$ is open and closed in $X'^{\text{an}} \setminus \varphi^{-1}(V)$, so B' is a connected component of $X'^{\text{an}} \setminus \varphi^{-1}(V)$ and hence $B' = C'$.

Now suppose that $\tau(C') \subset C'$. Then $\tau(C') \subset C' \cap \Sigma'$, so $C' \subset \tau^{-1}(C' \cap \Sigma')$. Let $y' \in \tau^{-1}(C' \cap \Sigma')$ and let B' be the connected component of $X'^{\text{an}} \setminus \Sigma'$ containing y' , as above. Since $\bar{B}' = B' \cup \{\tau(y')\}$ is a connected subset of $X'^{\text{an}} \setminus \varphi^{-1}(V)$ intersecting C' we have $\bar{B}' \subset C'$, so $y' \in C'$ and therefore $C' = \tau^{-1}(C' \cap \Sigma')$. \blacksquare

Proof of Theorem 5.13. Let $V = V(\mathfrak{X})$ and $V' = V(\mathfrak{X}')$. If there is an extension $\mathfrak{X}' \rightarrow \mathfrak{X}$ of φ , then the square

(5.18.1)

$$\begin{array}{ccc} X'^{\text{an}} & \xrightarrow{\varphi} & X^{\text{an}} \\ \text{red} \downarrow & & \downarrow \text{red} \\ \mathfrak{X}'_k & \longrightarrow & \mathfrak{X}_k \end{array}$$

commutes. Let $x' \in \varphi^{-1}(V)$, so $x = \varphi(x')$ reduces to a generic point $\bar{\zeta}$ of \mathfrak{X}_k . Since the reduction $\bar{\zeta}'$ of x' maps to $\bar{\zeta}$, the point $\bar{\zeta}'$ is generic, so $x' \in V'$. Therefore $\varphi^{-1}(V) \subset V'$. The morphism $\mathfrak{X}' \rightarrow \mathfrak{X}$ is finite if and only if every generic point of \mathfrak{X}'_k maps to a generic point of \mathfrak{X}_k ; as above, this is equivalent to $V' = \varphi^{-1}(V)$.

It remains to prove that if $\varphi^{-1}(V) \subset V'$, then φ extends to a morphism $\mathfrak{X}' \rightarrow \mathfrak{X}$. By Proposition 5.12, we must show that φ is continuous with respect to the topologies $\mathcal{T}(\mathfrak{X})$ and $\mathcal{T}(\mathfrak{X}')$. Let $U \subset X^{\text{an}}$ be $\mathcal{T}(\mathfrak{X})$ -open and let $U' = \varphi^{-1}(U)$. Clearly U is closed, so U' is closed (with respect to the ordinary topologies). We must show that condition (2) of Lemma 5.17 holds for U' . Let $x' \in U' \cap V'$ and let $x = \varphi(x')$. If $x \notin V$, then let $C \subset U$ be the connected component of $X^{\text{an}} \setminus V$ containing x and let C' be the connected component of $X'^{\text{an}} \setminus \varphi^{-1}(V)$ containing x' . Then $C' \subset U'$ because

$\varphi(C') \subset C$, and C' contains every connected component of $X'^{\text{an}} \setminus V'$ adjacent to x' since C' is an open neighborhood of x' .

Now suppose that $x \in V$. Any connected component of $X'^{\text{an}} \setminus V'$ which is adjacent to x' maps into a connected component of $X^{\text{an}} \setminus V$ which is adjacent to x . There are finitely many connected components $X^{\text{an}} \setminus V$ which are adjacent to x and not contained in U by Lemma 5.17. Let C be such a component. Since φ is finite, there are only finitely many connected components of $\varphi^{-1}(C)$; each of these is a connected component of $X'^{\text{an}} \setminus \varphi^{-1}(V)$. If C' is such a connected component, then either C' is an open ball connected component of $X'^{\text{an}} \setminus V'$ or C' intersects $\Sigma' = \Sigma(X', V')$ by Lemma 5.18. There are finitely many connected components of $X'^{\text{an}} \setminus V'$ which intersect Σ' — these are just the open annulus connected components of $X'^{\text{an}} \setminus V'$ — so there are only finitely many connected components of $X'^{\text{an}} \setminus V'$ contained in $\varphi^{-1}(C)$. Therefore all but finitely many connected components of $X'^{\text{an}} \setminus V'$ which are adjacent to x' map to connected components of $X^{\text{an}} \setminus V$ which are contained in U . \blacksquare

5.19. Simultaneous semistable reduction theorems. Recall that K_0 is a field equipped with a non-trivial non-Archimedean valuation $\text{val} : K_0 \rightarrow \mathbf{R} \cup \{\infty\}$. As above we fix an algebraic closure \overline{K}_0 of K_0 and a valuation val on \overline{K}_0 extending the given valuation on K_0 ; for any field $K_1 \subset \overline{K}_0$ we consider K_1 as a valued field with respect to the restriction of val , and we write R_1 for the valuation ring of K_1 . In what follows, X and X' are smooth, proper, geometrically connected K_0 -curves and $\varphi : X' \rightarrow X$ is a finite morphism.

Proposition 5.20. *Let \mathcal{X} be a semistable R_0 -model of X . If there exists a semistable R_0 -model \mathcal{X}' of X' such that $\varphi : X' \rightarrow X$ extends to a finite morphism $\mathcal{X}' \rightarrow \mathcal{X}$, then there is exactly one such model \mathcal{X}' up to (unique) isomorphism.*

Proof. Suppose first that $K = K_0$ is complete and algebraically closed. Let \mathfrak{X} (resp. \mathfrak{X}') be the ϖ -adic completion of \mathcal{X} (resp. \mathcal{X}'). In this case, the proposition follows from Theorem 5.8 and the fact that if the morphism $\mathfrak{X}' \rightarrow \mathfrak{X}$ of Theorem 5.13 is finite, then $V(\mathfrak{X}')$ is uniquely determined by $V(\mathfrak{X})$. The general case follows from this case after passing to the completion of the algebraic closure K of K_0 : if $\mathcal{X}', \mathcal{X}''$ are two semistable models of X' such that φ extends to finite morphisms $\mathcal{X}' \rightarrow \mathcal{X}$ and $\mathcal{X}'' \rightarrow \mathcal{X}$, then the isomorphism $\mathcal{X}''_R \xrightarrow{\sim} \mathcal{X}'_R$ descends to an isomorphism $\mathcal{X}'' \xrightarrow{\sim} \mathcal{X}'$ by Lemma 5.5. \blacksquare

Let $\mathcal{X}_1, \mathcal{X}_2$ be semistable R_0 -models of X . Recall that \mathcal{X}_1 *dominates* \mathcal{X}_2 if there exists a (necessarily unique) morphism $\mathcal{X}_1 \rightarrow \mathcal{X}_2$ inducing the identity on X .

Proposition 5.21. *Let \mathcal{X} and \mathcal{X}' be semistable R_0 -models of X and X' , respectively. Let $D' \subset X'(K_0)$ be a finite set of points. Then there exists a finite, separable extension K_1 of K_0 and a semistable R_1 -model \mathcal{X}'' of (X'_{K_1}, D') such that:*

- (1) \mathcal{X}'' dominates \mathcal{X}'_{R_1} ,
- (2) $\varphi_{K_1} : X'_{K_1} \rightarrow X_{K_1}$ extends to a morphism $\mathcal{X}'' \rightarrow \mathcal{X}_{R_1}$, and
- (3) any other semistable R_1 -model of (X'_{K_1}, D') satisfying the above two properties dominates \mathcal{X}'' .

Moreover, the formation of \mathcal{X}'' commutes with arbitrary valued field extensions $K_1 \rightarrow K'_1$.

Proof. Suppose first that $K_0 = K$ is complete and algebraically closed. Let \mathfrak{X} and \mathfrak{X}' be the ϖ -adic completions of \mathcal{X} and \mathcal{X}' , respectively. Let $V = V(\mathfrak{X})$ and $V' = V(\mathfrak{X}')$. By Lemma 3.15, there is a minimal semistable vertex set V'' of (X', D') which contains $\varphi^{-1}(V) \cup V'$. Let \mathfrak{X}'' be the semistable formal model of X' corresponding to V'' . Then \mathfrak{X}'' dominates \mathfrak{X}' by Corollary 5.15 and φ extends to a morphism $\mathfrak{X}'' \rightarrow \mathfrak{X}$ by Theorem 5.13. Part (3) follows from Corollary 5.15 and the minimality of V'' . Taking \mathcal{X}'' to be the algebraization of \mathfrak{X}'' yields (1)–(3) in this case.

For a general valued field K_0 , suppose that K is the completion of \overline{K}_0 . By Lemma 5.4 the model of X'_K constructed above descends to a model \mathcal{X}'' defined over the ring of integers of a finite, separable extension K_1 of K_0 . Properties (1)–(3) follow from Lemma 5.5 and the corresponding properties of \mathcal{X}''_R .

Now we address the behavior of this construction with respect to base change. Using Lemma 5.5 we immediately reduce to the case of an extension $K \rightarrow K'$ of complete and algebraically closed

valued fields. Let R' be the ring of integers of K' . Let \mathcal{X}'' (resp. \mathcal{X}''') be the minimal R -model of X' dominating \mathcal{X}' (resp. R' -model of $X'_{K'}$ dominating $\mathcal{X}'_{R'}$) mapping to \mathcal{X} (resp. $\mathcal{X}_{R'}$). By (3) as applied to \mathcal{X}''' , we have that $\mathcal{X}''_{R'}$ dominates \mathcal{X}''' . By Corollary 5.16, \mathcal{X}''' is defined over R , so by (3) as applied to \mathcal{X}'' , we have $\mathcal{X}''_{R'} = \mathcal{X}'''$. ■

Theorem 5.22. (Liu) *Let \mathcal{X} (resp. \mathcal{X}') be a semistable R_0 -model of X (resp. X'). Let $D \subset X(K_0)$ and $D' \subset X'(K_0)$ be finite sets, and suppose that $\varphi(D') \subset D$. Then there exists a finite, separable extension K_1 of K_0 and semistable R_1 -models $\mathcal{X}_1, \mathcal{X}'_1$ of (X_{K_1}, D) , (X'_{K_1}, D') , respectively, such that*

- (1) \mathcal{X}_1 dominates \mathcal{X}_{R_1} and \mathcal{X}'_1 dominates \mathcal{X}'_{R_1} ,
- (2) the morphism $\varphi_{K_1} : X'_{K_1} \rightarrow X_{K_1}$ extends to a finite morphism $\mathcal{X}'_1 \rightarrow \mathcal{X}_1$, and
- (3) if $\mathcal{X}_2, \mathcal{X}'_2$ are semistable formal models of (X_{K_1}, D) , (X'_{K_1}, D') , respectively, satisfying (1) and (2) above, then \mathcal{X}_2 dominates \mathcal{X}_1 and \mathcal{X}'_2 dominates \mathcal{X}'_1 .

Moreover, the formation of $\mathcal{X}'_1 \rightarrow \mathcal{X}_1$ commutes with arbitrary valued field extensions $K_1 \rightarrow K'_1$.

The morphism $\mathcal{X}'_1 \rightarrow \mathcal{X}_1$ is called the *stable marked hull* of $\mathcal{X}' \dashrightarrow \mathcal{X}$ in [Liu06].

Proof. First assume that $K_0 = K$ is complete and algebraically closed; let $\mathfrak{X}, \mathfrak{X}'$ be the ϖ -adic completions of $\mathcal{X}, \mathcal{X}'$, respectively. Let $V = V(\mathfrak{X})$ and $V' = V(\mathfrak{X}')$. By Theorem 5.8, Theorem 5.13, and Corollary 5.15, we may equivalently formulate the existence and uniqueness of $\mathfrak{X}'_1 \rightarrow \mathfrak{X}_1$ in terms of semistable vertex sets, as follows. We must prove that there exists a semistable vertex set V_1 of (X, D) such that

- (1) V_1 contains $V \cup \varphi(V')$,
- (2) $\varphi^{-1}(V_1)$ is a semistable vertex set of (X', D') , and
- (3) V_1 is minimal in the sense that if V_2 is another semistable vertex set of (X, D) satisfying (1) and (2) above, then $V_2 \supset V_1$.

First we will prove the existence of V_1 satisfying (1) and (2). By Lemma 3.15 we may enlarge V to assume that V is a semistable vertex set of (X, D) . Let $\Sigma = \Sigma(X, V \cup D)$ and $\Sigma' = \Sigma(X', V')$. By Corollary 4.18, there exists a skeleton Σ_1 of (X, D) such that $\Sigma_1 \supset \Sigma \cup \varphi(\Sigma')$ and such that $\Sigma'_1 = \varphi^{-1}(\Sigma_1)$ is a skeleton of $(X', \varphi^{-1}(D))$. Let V'_1 be a vertex set for Σ'_1 . Since any finite subset of type-2 points of Σ'_1 which contains V'_1 is again a vertex set for Σ'_1 by Proposition 3.12(4), we may and do assume that $V' \subset V'_1$. Let $V_1 \subset \Sigma_1$ be the union of a vertex set for Σ_1 with $V \cup \varphi(V'_1)$. Then V_1 is a semistable vertex set of (X, D) , and $\varphi^{-1}(V_1) \subset \Sigma'_1$ is a finite set of type-2 points containing V'_1 , thus is a semistable vertex set of $(X', \varphi^{-1}(D))$ (hence of (X', D') as well).

To prove that there exists a minimal such V_1 , we make the following recursive construction. Let $V(0) = V$, let $V'(0) = V'$, and for each $n \geq 1$ let $V(n)$ be the minimal semistable vertex set of (X, D) containing $V(n-1) \cup \varphi(V'(n-1))$ and let $V'(n)$ be the minimal semistable vertex set of (X', D') containing $\varphi^{-1}(V(n))$. These sets exist by Lemma 3.15. By induction it is clear that if V_2 is any semistable vertex set of (X, D) satisfying (1) and (2) above, then $V(n) \subset V_2$ for each n . Since V_2 is a finite set, for some n we have $V(n) = V(n+1)$, which is to say that $\varphi(V'(n)) \subset V(n)$; since $V'(n) \supset \varphi^{-1}(V(n))$, we have that $V'(n) = \varphi^{-1}(V(n))$ is a semistable vertex set of (X', D') . Then $V_1 = V(n)$ is the minimal semistable vertex set satisfying (1) and (2) above.

The case of a general ground field reduces to the geometric case handled above exactly as in the proof of Proposition 5.21, as does the statement about the behavior of the stable marked hull with respect to valued field extensions. ■

Remark 5.23. (The skeletal viewpoint on Liu's theorem) The statement of Theorem 5.22 is strongly analogous to Corollary 4.18, which is indeed the main ingredient in the proof. The difference is that whereas finite morphisms of semistable models correspond to pairs V, V' of semistable vertex sets such that $\varphi^{-1}(V) = V'$, for finite morphisms of triangulated punctured curves one requires in addition that $\varphi^{-1}(\Sigma) = \Sigma'$. The former condition (of Liu's theorem) does not imply the latter: for instance, let X' be the Tate curve $y^2 = x^3 - x + \varpi$, let $X = \mathbf{P}^1$, and let $\varphi : X' \rightarrow X$ be the cover $(x, y) \mapsto x$. This extends to a finite morphism of semistable models given by the same equations; however, the associated skeleton of X' (resp. of \mathbf{P}^1) is a circle (resp. a point), so $\varphi^{-1}(\Sigma) \subsetneq \Sigma'$.

In general, if $\varphi^{-1}(V) = V'$ and $\varphi^{-1}(D) = D'$, then by Remark 4.19 the image of the skeleton $\Sigma(X', V' \cup D')$ is equal to $\Sigma(X, V \cup D)$ union a finite number of geodesic segments T_1, \dots, T_r , and by Theorem 4.13 the saturation $\varphi^{-1}(\varphi(\Sigma(X', V' \cup D')))$ is a skeleton of X' .

Theorem 5.25 below follows the same philosophy in deriving a simultaneous semistable reduction theorem of Liu–Lorenzini from Proposition 4.12.

Remark 5.24. Liu in fact works over an arbitrary Dedekind scheme (a connected Noetherian regular scheme of dimension 1), which includes discrete valuation rings but not more general valuation rings. In his statement of Theorem 5.22 the given models $\mathcal{X}, \mathcal{X}'$ are allowed to be any integral, projective $\text{Spec}(R_0)$ -schemes with generic fibers X and X' , respectively. Although we restrict to semistable models $\mathcal{X}, \mathcal{X}'$ in this paper, using a more general notion of a triangulation our methods can be extended to treat relatively normal models $\mathcal{X}, \mathcal{X}'$.

The following simultaneous stable reduction theorem can be found in [LL99]. Theorem 5.25 is to Proposition 4.12 as Theorem 5.22 is to Corollary 4.18

Theorem 5.25. (Liu–Lorenzini) *Suppose that (X, D) and (X', D') are both stable and that $\varphi^{-1}(D) = D'$. Assume that (X, D) and (X', D') admit stable models \mathcal{X} and \mathcal{X}' , respectively, defined over R_0 . Then $\varphi : X' \rightarrow X$ extends to a (not necessarily finite) morphism $\mathcal{X}' \rightarrow \mathcal{X}$.*

Proof. By Lemma 5.5 we may assume that $K_0 = K$ is complete and algebraically closed. Let V (resp. V') be the minimal semistable vertex set of (X, D) (resp. (X', D')). By Theorem 5.13, we must show that $\varphi^{-1}(V) \subset V'$. Let $\Sigma = \Sigma(X, V)$ and $\Sigma' = \Sigma(X', V')$. Let V_1 be a vertex set for Σ with respect to which Σ has no loop edges. Recall that

$$V = \{x \in V_1 : g(x) \geq 1 \text{ or } x \text{ has valency at least } 3\}$$

by Proposition 3.18(2). Let $V'_1 = \varphi^{-1}(V_1) \cup V'$; this is a vertex set for Σ' since $\varphi^{-1}(\Sigma) \subset \Sigma'$ by Proposition 4.12(2). If $x \in V$ has genus at least 1, then any $x' \in \varphi^{-1}(x)$ has genus at least 1, so $\varphi^{-1}(x) \subset V'$. Let $x \in V$ have genus 0, so x has valency at least 3 in Σ . Let $x' \in \varphi^{-1}(x) \subset \Sigma'$ and let U' be the set containing x' and all of the connected components of $X'^{\text{an}} \setminus V'_1$ adjacent to x' . Then U' is an open neighborhood of x' , so $\varphi(U')$ is an open neighborhood of x by [Ber90, Lemma 3.2.4]. Let A be an open annulus connected component of $X^{\text{an}} \setminus V_1$ adjacent to x . Any open neighborhood of x intersects A , so there exists $y' \in U'$ with $\varphi(y') \in A$. Let A' be the connected component of $U' \setminus \{x'\}$ containing y' , so $\varphi(A') \subset A$. Since x' is an end of A' and $\varphi(x')$ is an end of A , Lemma 4.3 implies that A' is an open annulus. Since x has valency at least 3 in Σ (and since Σ has no loop edges), there are at least 3 distinct open annulus connected components of $X^{\text{an}} \setminus V_1$ adjacent to x . By the above, the same is true of x' , so x' has valency at least 3 in Σ' , so $x' \in V'$, as desired. \blacksquare

6. A LOCAL LIFTING THEOREM

We now begin with the classification of lifts of harmonic morphisms of metrized complexes of curves to finite morphisms of algebraic curves. We begin by working in the neighborhood of a vertex of a metrized complex.

6.1. Let X be a smooth, proper, connected curve over K and let $x \in X^{\text{an}}$ be a type-2 point. Let $V \subset X^{\text{an}}$ be a strongly semistable vertex set of X containing x , let $\Sigma = \Sigma(X, V)$, and let $\tau : X^{\text{an}} \rightarrow \Sigma$ be the retraction. Let e_1, \dots, e_r be the edges of Σ adjacent to x and let $\Sigma_0 = \{x\} \cup e_1^\circ \cup \dots \cup e_r^\circ$, where for a (closed) edge e we let e° denote the corresponding open edge, i.e. the edge without its endpoints. Then Σ_0 is an open neighborhood of x in Σ and $\tau^{-1}(\Sigma_0)$ is an open neighborhood of x in X^{an} . Following [BPR11, 5.54] and [Ber93], we define a *simple neighborhood* of x to be an open neighborhood of this form (for some choice of V). The connected components of $\tau^{-1}(\Sigma_0) \setminus \{x\}$ are open balls and the open annuli $\tau^{-1}(e_1^\circ), \dots, \tau^{-1}(e_r^\circ)$.

Definition 6.2. A *star-shaped curve* is a pointed K -analytic space (Y, y) which is isomorphic to (U, x) where x is a type-2 point in the analytification of a smooth, proper, connected curve over K and U is a simple neighborhood of x . The point y is called the *central vertex* of Y .

6.3. Let (Y, y) be a star-shaped curve, so $Y \setminus \{y\}$ is a disjoint union of open balls and finitely many open annuli A_1, \dots, A_r . The *skeleton* of Y is defined to be the set $\Sigma(Y, \{y\}) = \{y\} \cup \bigcup_{i=1}^r \Sigma(A_i)$. A *compatible divisor in Y* is a finite set $D \subset Y(K)$ whose points are contained in distinct open ball connected components of $Y \setminus \{y\}$, so the connected components of $Y \setminus (\{y\} \cup D)$ are open balls, the open annuli A_1, \dots, A_r , and (finitely many) open balls B_1, \dots, B_s punctured at a point of D . The data (Y, y, D) of a star-shaped curve along with a compatible divisor is called a *punctured star-shaped curve*. The *skeleton of (Y, y, D)* is the set

$$\Sigma(Y, \{y\} \cup D) = \{y\} \cup D \cup \bigcup_{i=1}^r \Sigma(A_i) \cup \bigcup_{j=1}^s \Sigma(B_j).$$

Fix a compatible divisor D in Y , and let $\Sigma_0 = \Sigma(Y, \{y\} \cup D)$. There is a canonical continuous retraction map $\tau : Y \rightarrow \Sigma_0$ defined exactly as for skeleta of algebraic curves (3.11). The connected components of $\Sigma_0 \setminus \{y\}$ are called the *edges* of Σ_0 ; an edge is called *infinite* if it contains a point of D and *finite* otherwise.

If Y is the simple neighborhood $\tau^{-1}(\Sigma_0)$ of $x \in X^{\text{an}}$ as above, then $\Sigma(Y, \{x\}) = \Sigma_0 = \Sigma \cap Y$. A finite set $D \subset Y(K)$ is compatible with Y if and only if V is a semistable vertex set for (X, D) , in which case $\Sigma(Y, \{x\} \cup D) = Y \cap \Sigma(X, V \cup D)$. The retraction $\tau : Y \rightarrow \Sigma(Y, \{x\} \cup D)$ is the restriction of the canonical retraction $\tau : X^{\text{an}} \rightarrow \Sigma(X, V \cup D)$.

6.4. Let (Y, y) be a star-shaped curve. Then Y is proper as a K -analytic space if and only if all connected components of $Y \setminus \{y\}$ are open balls, i.e. if and only if $\Sigma(Y, \{y\}) = \{y\}$. If Y is proper, then there is a smooth, proper, connected curve X over K and an isomorphism $f : Y \xrightarrow{\sim} X^{\text{an}}$. Let $x = f(y)$. Then $\{x\}$ is a semistable vertex set of X , so by Theorem 5.8 there is a unique smooth formal model \mathfrak{X} of X^{an} such that x reduces to the generic point of \mathfrak{X}_k . Let $D \subset Y(K)$ be a finite set. Then D is compatible with Y if and only if the points of $f(D)$ reduce to distinct closed points of \mathfrak{X}_k .

Conversely, let \mathfrak{X} be a smooth, proper, connected formal curve over $\text{Spf}(R)$. If $x \in X^{\text{an}}$ is the point reducing to the generic point of \mathfrak{X}_k , then (\mathfrak{X}_K, x) is a proper star-shaped curve.

Proposition 6.5. *Let (Y, y) be a star-shaped curve. Then (Y, y) is isomorphic to (U, x) where x is the central vertex of a proper star-shaped curve X and U is a simple neighborhood of x .*

Proof. Let A_1, \dots, A_r be the open annulus connected components of $Y \setminus \{y\}$. Choose isomorphisms $f_i : A_i \xrightarrow{\sim} \mathbf{S}(a_i)_+$ with standard open annuli such that $f_i(x)$ approaches the Gauss point of $\mathbf{B}(1)$ as x approaches y . Let X be the curve obtained from Y by gluing an open ball $\mathbf{B}(1)_+$ onto each A_i via the inclusions $f_i : A_i \xrightarrow{\sim} \mathbf{S}(a_i)_+ \subset \mathbf{B}(1)_+$. Then X is a smooth, proper, connected K -analytic curve, and it is clear from the construction that (X, y) is star-shaped and that Y is a simple neighborhood of y in X . ■

6.6. A proper star-shaped curve X and an inclusion $i : Y \xrightarrow{\sim} U \subset X$ as in Proposition 6.5 is called a *compactification* of Y (as a star-shaped curve). Note that $X \setminus i(Y)$ is a disjoint union of finitely many closed balls, one for each open annulus connected component of $Y \setminus \{y\}$.

6.7. Let (Y, y) be a star-shaped curve. The smooth, proper, connected k -curve C_y with function field $\mathcal{H}(y)$ is called the *residue curve* of Y . The tangent vectors in T_y are naturally in bijective correspondence with the connected components of $Y \setminus \{y\}$. We define a *reduction map* $\text{red} : Y \rightarrow C_y$ by sending y to the generic point of C_y , and sending every point in a connected component B of $Y \setminus \{y\}$ to the closed point of C_y corresponding to the tangent vector determined by B . This sets up a one-to-one correspondence between the connected components of $Y \setminus \{y\}$ and the closed points of C_y .

Remark 6.8.

- (1) When Y is proper, so $Y \cong \mathfrak{X}_K$ for a smooth, proper, connected formal curve over $\text{Spf}(R)$, then $C_y \cong \mathfrak{X}_k$ and the reduction map $Y \rightarrow C_y$ coincides with the canonical reduction map $\mathfrak{X}_K \rightarrow \mathfrak{X}_k$.

- (2) Let $D \subset Y(K)$ be a compatible divisor and let $\Sigma_0 = \Sigma(Y, \{y\} \cup D)$. Every edge e of Σ_0 is contained in a unique connected component of $Y \setminus \{y\}$, and $\text{red}(e) \in C_y(k)$ is the closed point corresponding to the tangent direction represented by e .

6.9. Tame coverings. We now study a class of morphisms of star-shaped curves analogous to (4.30). We begin with the following technical result.

Lemma 6.10. *Let A, A' be open annuli or punctured open balls and let $\varphi : A' \rightarrow A$ be a finite morphism of degree δ . Suppose that δ is prime to $\text{char}(k)$ if $\text{char}(k) > 0$. Fix an isomorphism $A \cong \mathbf{S}(a)_+$ (where we allow $a = 0$).*

- (1) *There is an isomorphism $A' \cong \mathbf{S}(a')_+$ such that the composition*

$$(6.10.1) \quad \mathbf{S}(a')_+ \cong A' \xrightarrow{\varphi} A \cong \mathbf{S}(a)_+$$

is $t \mapsto t^\delta$.

- (2) *There is an isomorphism $A' \cong \mathbf{S}(a')_+$ such that (6.10.1) extends to a morphism $\psi : \mathbf{B}(1)_+ \rightarrow \mathbf{B}(1)_+$ with $\psi^{-1}(0) = 0$.*
(3) *If (6.10.1) extends to a morphism $\psi : \mathbf{B}(1)_+ \rightarrow \mathbf{B}(1)_+$ for a given isomorphism $A' \cong \mathbf{S}(a')_+$, then the extension is unique.*

Proof. Let u be a parameter on A' , i.e. an isomorphism $u : A' \xrightarrow{\sim} \mathbf{S}(a')_+ \subset \mathbf{B}(1)_+$ with a standard open annulus. By Lemma 4.2, φ restricts to an affine map on skeleta $\Sigma(A') \rightarrow \Sigma(A)$ with degree δ . If $B = \mathbf{S}(b, c)$ is a closed sub-annulus of $\mathbf{S}(a')_+$, then by [Thu05, Lemme 2.2.1], after potentially replacing u by u^{-1} , $\varphi^*(t)$ has the form $\alpha u^\delta(1 + g(u))$ on B , where $\alpha \in R^\times$ and $|g|_{\text{sup}} < 1$. Since δ is not divisible by $\text{char}(k)$ if $\text{char}(k) > 0$, the Taylor expansion for $\sqrt[\delta]{1 + g}$ has coefficients contained in R , hence converges to a δ th root of $1 + g(u)$ on A . Choosing a δ th root of α as well and letting $u_B = u \sqrt[\delta]{\alpha} \sqrt[\delta]{1 + g}$, we have that u_B is a parameter on B such that $\varphi^*(t) = u_B^\delta$. Hence for each such B there are exactly δ choices of a parameter u_B on B such that $\varphi^*(t) = u_B^\delta$; choosing a compatible set of such parameters for all B yields a parameter on A' satisfying (1).

Part (2) follows immediately from (1). A morphism from a K -analytic space X to $\mathbf{B}(1)_+$ is given by a unique analytic function f on X such that $|f(x)| < 1$ for all $x \in X$, so (3) follows from the fact that the restriction homomorphism $\mathcal{O}(\mathbf{B}(1)_+) \rightarrow \mathcal{O}(\mathbf{S}(a')_+)$ is injective. \blacksquare

6.11. Let A be an open annulus. Let δ be a positive integer, and assume that δ is prime to the characteristic of k if $\text{char}(k) > 0$. Let A' be an open annulus and let $\varphi : A' \rightarrow A$ be a finite morphism of degree δ . By Lemma 6.10, the group $\text{Aut}_A(A')$ is isomorphic to $\mathbf{Z}/\delta\mathbf{Z}$. Moreover, if A'' is another open annulus and $\varphi' : A'' \rightarrow A$ is a finite morphism of degree δ , then there exists an A -isomorphism $\psi : A' \xrightarrow{\sim} A''$; the induced isomorphism $\text{Aut}_A(A') \xrightarrow{\sim} \text{Aut}_A(A'')$ is independent of the choice of ψ since both groups are abelian. Therefore the group $\text{Aut}_A(\delta) := \text{Aut}_A(A') \cong \mathbf{Z}/\delta\mathbf{Z}$ is canonically determined by A and δ .

Definition 6.12. Let (Y, y, D) be a punctured star-shaped curve with skeleton $\Sigma_0 = \Sigma(Y, \{y\} \cup D)$ and let C_y be the residue curve of Y . A *tame covering* of (Y, y, D) consists of a punctured star-shaped curve (Y', y', D') and a finite morphism $\varphi : Y' \rightarrow Y$ satisfying the following properties:

- (1) $\varphi^{-1}(y) = \{y'\}$,
- (2) $D' = \varphi^{-1}(D)$, and
- (3) if $C_{y'}$ denotes the residue curve of Y' and $\varphi_{y'} : C_{y'} \rightarrow C_y$ is the morphism induced by φ , then $\varphi_{y'}$ is tamely ramified and is branched only over the points of C_y corresponding to tangent directions at y represented by edges in Σ_0 .

Remark 6.13. The degree of φ is equal to the degree of $\varphi_{y'}$.

Example 6.14. Let $\varphi : (X', V', D') \rightarrow (X, V, D)$ be a tame covering of triangulated punctured curves. Let $\Sigma = \Sigma(X, V \cup D)$ and $\Sigma' = \Sigma(X', V' \cup D') = \varphi^{-1}(\Sigma)$. Let $x \in V$ be a finite vertex of Σ , let e_1, \dots, e_r be the finite edges of Σ adjacent to x , let $\Sigma_0 = \{x\} \cup \bigcup_{i=1}^r e_i^\circ$, let $Y = \tau^{-1}(\Sigma_0)$, and let $D_0 = D \cap Y$. Then (Y, x, D_0) is a punctured star-shaped curve. Let Σ'_0 be a connected component of $\varphi^{-1}(\Sigma_0)$, let

$x' \in \Sigma'_0$ be the unique inverse image of x , let $Y' = \tau^{-1}(\Sigma'_0)$, and let $D'_0 = D' \cap Y'$. Then (Y', x', D'_0) is also a punctured star-shaped curve, and φ restricts to a finite morphism $\varphi : Y' \rightarrow Y$. This is in fact a tame covering of punctured star-shaped curves by Remark 4.34 and Proposition 4.35.

Proposition 6.15. *Let $\varphi : (Y', y', D') \rightarrow (Y, y, D)$ be a degree- δ tame covering of punctured star-shaped curves.¹ Let $\Sigma_0 = \Sigma(Y, \{y\} \cup D)$, let $\Sigma'_0 = \Sigma(Y', \{y'\} \cup D')$, let C_y (resp. $C_{y'}$) be the residue curve of Y (resp. Y'), and let $\varphi_{y'} : C_{y'} \rightarrow C_y$ be the induced morphism.*

- (1) *Let B be a connected component of $Y \setminus \{y\}$ disjoint from Σ_0 . Then φ^{-1} is a disjoint union of δ open balls mapping isomorphically onto B .*
- (2) *Let B be a connected component of $Y \setminus \{y\}$ meeting D , and choose an isomorphism of B with $\mathbf{B}(1)_+$ which identifies the unique point of $B \cap D$ with 0. Let B' be a connected component of $\varphi^{-1}(B)$. Then $\varphi|_{B'} : B' \rightarrow B$ is a finite morphism, the degree of $\varphi|_{B'}$ is the ramification degree δ' of $\varphi_{y'}$ at $\text{red}(B')$, and there is an isomorphism $B' \cong \mathbf{B}(1)_+$ sending the unique point of $B' \cap D'$ to 0 such that the composition*

$$\mathbf{B}(1)_+ \cong B' \xrightarrow{\varphi} B \cong \mathbf{B}(1)_+$$

is $t \mapsto t^{\delta'}$.

- (3) *Let A be an open annulus connected component of $Y \setminus \{y\}$, and choose an isomorphism $A \cong \mathbf{S}(a)_+$. Let A' be a connected component of $\varphi^{-1}(A)$. Then $\varphi|_{A'} : A' \rightarrow A$ is a finite morphism, the degree of $\varphi|_{A'}$ is the ramification degree δ' of $\varphi_{y'}$ at $\text{red}(A')$, and there is an isomorphism of A' with an open annulus $\mathbf{S}(a')$ such that the composition*

$$\mathbf{S}(a')_+ \cong A' \xrightarrow{\varphi} A \cong \mathbf{S}(a)_+$$

is $t \mapsto t^{\delta'}$.

- (4) *φ is étale over $Y \setminus D$.*
- (5) *$\varphi^{-1}(\Sigma_0) = \Sigma'_0$.*

Proof. In the situation of (1), let $\bar{x} = \text{red}(B)$. Since $\varphi_{y'}$ is not branched over \bar{x} , there are δ distinct points of $C_{y'}$ mapping to \bar{x} ; hence there are δ connected components B'_1, \dots, B'_δ of $Y' \setminus \{y'\}$ mapping onto B . The restriction of φ to each B'_i is finite of degree 1.

Next we prove (2). It is clear that $\varphi|_{B'} : B' \rightarrow B$ is finite, hence surjective; since B' is a connected component of $Y' \setminus \{y'\}$ containing a point of $D' = \varphi^{-1}(D)$, it is an open ball. Let x (resp. x') be the unique point of D (resp. D') contained in B (resp. B') and let $e \subset \Sigma_0$ (resp. $e' \subset \Sigma'_0$) be the edge adjacent to x (resp. x'). Then $B' \setminus \{x'\} \rightarrow B \setminus \{x\}$ is a finite morphism of punctured open balls, so by Lemma 3.3, Proposition 4.8, and Theorem 4.23(2), the restriction of φ to e' is affine morphism $e' \rightarrow e$ of degree δ' , and δ' is the degree of $B' \rightarrow B$. So by Lemma 6.10 there is an isomorphism $B' \cong \mathbf{B}(1)_+$ as described in the statement of the Theorem.

In the situation of (3), we claim that A' is an open annulus. Clearly $\varphi|_{A'} : A' \rightarrow A$ is finite, hence surjective. If A' is not an open annulus, then $A' \cong \mathbf{B}(1)_+$ is an open ball. The morphism $\mathbf{B}(1)_+ \cong A' \rightarrow A \cong \mathbf{S}(a)_+$ is given by a unit on $\mathbf{B}(1)_+$, which has constant absolute value; this contradicts surjectivity, so A' is in fact an open annulus. The proof now proceeds exactly as above.

Parts (4) and (5) follow immediately from parts (1)–(3). ■

The following proposition is reminiscent of [Sai04, Proposition 3.3.2].

Corollary 6.16. *Let $\varphi : (Y', y', D'_0) \rightarrow (Y, y, D_0)$ be a tame covering of punctured star-shaped curves and let $i : (Y, y) \hookrightarrow (X, x)$ be a compactification of Y . Let D_1 be the union of $i(D_0)$ with a choice of K -point from every connected component of $X \setminus i(Y)$. Then there exists a compactification $i' : (Y', y') \hookrightarrow (X', x')$ and a tame covering $\psi : (X', x', D'_1) \rightarrow (X, x, D_1)$ such that $\psi \circ i' = i \circ \varphi$.*

Proof. We compactify (Y', y') as in the proof of Proposition 6.5, gluing balls onto the annulus connected components of $Y' \setminus \{y'\}$. By Proposition 6.15(3), if $A \cong \mathbf{S}(a)_+$ is an open annulus connected component of $Y \setminus \{y\}$ and $A' \subset Y' \setminus \{y'\}$ is a connected component mapping to A , then we can choose

¹This δ need not be prime to $\text{char}(k)$.

an isomorphism $A' \cong \mathbf{S}(a')_+$ such that $\mathbf{S}(a')_+ \cong A' \rightarrow A \cong \mathbf{S}(a)_+$ is of the form $t \mapsto t^\delta$; this map extends to a morphism $\mathbf{B}(1)_+ \rightarrow \mathbf{B}(1)_+$, and these maps glue to give a tame covering $X' \rightarrow X$. \blacksquare

6.17. A local lifting theorem. Let (Y, y, D) be a punctured star-shaped curve with skeleton $\Sigma_0 = \Sigma(Y, \{y\} \cup D)$ and residue curve C_y . Let C' be a smooth, proper, connected k -curve and let $\bar{\varphi} : C' \rightarrow C_y$ be a finite, tamely ramified morphism branched only over the points of C_y corresponding to tangent directions at y represented by edges in Σ_0 . A *lifting of C' to a punctured star-shaped curve over (Y, y, D)* is the data of a punctured star-shaped curve (Y', y', D') , a tame covering $\varphi : (Y', y', D') \rightarrow (Y, y, D)$, and an isomorphism of the residue curve $C_{y'}$ with C' which identifies $\bar{\varphi}$ with the morphism $\varphi_{y'} : C_{y'} \rightarrow C_y$ induced by φ . An *isomorphism between two liftings (Y', y') and (Y'', y'')* is a Y -isomorphism $Y' \rightarrow Y''$ such that the induced morphism $C_{y'} \xrightarrow{\sim} C_{y''}$ respects the identifications $C_{y'} \cong C'$ and $C_{y''} \cong C'$.

Theorem 6.18. *Let (Y, y, D) be a punctured star-shaped curve with skeleton $\Sigma_0 = \Sigma(Y, \{y\} \cup D)$ and residue curve C_y , let C' be a smooth, proper, connected k -curve, and let $\bar{\varphi} : C' \rightarrow C_y$ be a finite, tamely ramified morphism branched only over the points of C_y corresponding to tangent directions at y represented by edges in Σ_0 . Then there exists a lifting of C' to a punctured star-shaped curve over (Y, y, D) , and this lifting is unique up to unique isomorphism.*

Before giving the proof of this theorem, we need a technical lemma. Let \mathcal{X} be a finitely presented, flat, separated R -scheme and let \mathfrak{X} be its ϖ -adic completion; this is an admissible formal R -scheme by [BPR11, Proposition 3.12]. There is a canonical open immersion $i_{\mathcal{X}} : \mathfrak{X}_K \rightarrow \mathcal{X}_K^{\text{an}}$ defined in [Con99, §A.3] which is functorial in \mathcal{X} and respects the formation of fiber products, and is an isomorphism when \mathcal{X} is proper over R . (This fact is implicitly contained in the statement of Lemma 5.1.)

Lemma 6.19. *Let $\mathcal{X}, \mathcal{X}'$ be finitely presented, flat, separated R -schemes and let $\mathfrak{X}, \mathfrak{X}'$ denote their ϖ -adic completions. Let $\mathcal{X}' \rightarrow \mathcal{X}$ be a finite and flat morphism. Then the square*

$$(6.19.1) \quad \begin{array}{ccc} \mathfrak{X}'_K & \xrightarrow{i_{\mathcal{X}'}} & \mathcal{X}'_K^{\text{an}} \\ f \downarrow & & \downarrow g \\ \mathfrak{X}_K & \xrightarrow{i_{\mathcal{X}}} & \mathcal{X}_K^{\text{an}} \end{array}$$

is Cartesian.

Proof. The vertical arrows f, g of (6.19.1) are finite and the horizontal arrows $i_{\mathcal{X}'}, i_{\mathcal{X}}$ are open immersions. Let $Y = \mathcal{X}'_K^{\text{an}} \times_{\mathcal{X}_K^{\text{an}}} \mathfrak{X}_K$, so $Y \rightarrow \mathcal{X}'_K^{\text{an}}$ is an open immersion and $Y \rightarrow \mathfrak{X}_K$ is finite. Let $h : \mathfrak{X}'_K \rightarrow Y$ be the canonical morphism. Then h is an open immersion because its composition with $Y \rightarrow \mathcal{X}'_K^{\text{an}}$ is an open immersion, and h is finite because its composition with $Y \rightarrow \mathfrak{X}_K$ is finite. Therefore h is an isomorphism of \mathfrak{X}'_K onto an open and closed subspace of Y . It suffices to show that h is surjective, and since $Y(K)$ is dense in Y , we only need to check that $Y(K)$ is in the image of h .

Let $x \in \mathfrak{X}_K(K)$ and $y \in \mathcal{X}'_K^{\text{an}}(K)$ be points with the same image $z \in \mathcal{X}_K^{\text{an}}(K)$, so $(x, y) \in Y(K)$. Choose an open formal affine $\mathfrak{U} \subset \mathfrak{X}$ such that $x \in \mathfrak{U}_K(K)$, and assume without loss of generality that there is an affine open $\mathcal{U} = \text{Spec}(A) \subset \mathcal{X}$ such that $\mathfrak{U} = \text{Spf}(\hat{A})$, where \hat{A} is the ϖ -adic completion of A . Then x corresponds to a continuous K -homomorphism $\eta : \hat{A} \otimes_R K \rightarrow K$. We have $\eta(\hat{A}) \subset R$, so composing with the completion homomorphism $A \rightarrow \hat{A}$ we obtain an R -homomorphism $A \rightarrow R$. Let $\xi : \text{Spec}(R) \rightarrow \mathcal{U} \subset \mathcal{X}$ denote the induced morphism. The corresponding morphism $\text{Spf}(R)_K \rightarrow \mathfrak{X}_K$ is the point x .

Let \mathcal{Z} be the fiber product of $\mathcal{X}' \rightarrow \mathcal{X}$ with $\eta : \text{Spec}(R) \rightarrow \mathcal{X}$ and let \mathfrak{Z} be its ϖ -adic completion. Note that \mathcal{Z} is finite and flat over $\text{Spec}(R)$. In particular, \mathcal{Z} is proper over R , so $i_{\mathcal{Z}} : \mathfrak{Z}_K \rightarrow \mathcal{Z}_K^{\text{an}}$ is an

isomorphism. We have a commutative cube

$$\begin{array}{ccccc}
 & & \mathcal{Z}_K^{\text{an}} & \longrightarrow & \mathcal{X}'_K{}^{\text{an}} \\
 & \nearrow \cong & \downarrow & \nearrow & \downarrow g \\
 \mathfrak{Z}_K & \longrightarrow & \mathfrak{X}'_K & \longrightarrow & \mathcal{X}'_K{}^{\text{an}} \\
 & \searrow \cong & \downarrow f & \searrow & \downarrow \\
 & & \text{Spec}(K)^{\text{an}} & \xrightarrow{z} & \mathcal{X}_K{}^{\text{an}} \\
 \text{Spf}(R)_K & \xrightarrow{x} & \mathfrak{X}_K & \longrightarrow & \mathcal{X}_K{}^{\text{an}}
 \end{array}$$

where the front and back faces are Cartesian and the diagonal arrows are the canonical open immersions. Since $g(y) = z$ and $\mathcal{Z}_K^{\text{an}} = f^{-1}(z)$, the point y lifts to a point in $\mathcal{Z}_K^{\text{an}}(K)$, which then lifts to a point in $\mathfrak{Z}_K(K)$ whose image in $\mathfrak{X}'_K(K)$ maps to x in $\mathfrak{X}_K(K)$ and $y \in \mathcal{X}'_K{}^{\text{an}}(K)$. \blacksquare

6.20. Proof of Theorem 6.18. We first prove the theorem when Y is proper. In this case we may and do assume that Y is the analytic generic fiber of a smooth, proper, connected formal curve \mathfrak{X} over $\text{Spf}(R)$. Let $\mathcal{X} \rightarrow \text{Spec}(R)$ be the algebraization of \mathfrak{X} (Lemma 5.1); this is a smooth, proper relative curve of finite presentation and with connected fibers whose ϖ -adic completion is isomorphic to \mathfrak{X} . Note that $\mathcal{X}_k = \mathfrak{X}_k = C_y$. Let $X = \mathcal{X}_K$, so $X^{\text{an}} = \mathfrak{X}_K = Y$. By the valuative criterion of properness, every point x of D extends uniquely to a section $\text{Spec}(R) \rightarrow \mathcal{X}$ which sends the closed point to the reduction of x ; hence the closure \mathcal{D} of D in \mathcal{X} is a disjoint union of sections. Let $\mathcal{U} = \mathcal{X} \setminus \mathcal{D}$.

The theory of the *tamely ramified étale fundamental group* π_1^t of a morphism of schemes with a relative normal crossings divisor is developed in [SGA1, Exposé XIII]. The finite-index subgroups of π_1^t classify so-called tamely ramified étale covers. The subscheme $\mathcal{D} \subset \mathcal{X}$ is a relative normal crossings divisor relative to $\text{Spec}(R)$, so the proof of Corollaire 2.12 of *loc. cit.* shows that the specialization homomorphism $\pi_1(\mathcal{U}_K) \rightarrow \pi_1^t(\mathcal{U}_k)$ is surjective (we suppress the base points). What this means concretely is that every tamely ramified cover $\overline{U}' \rightarrow \mathcal{U}_k$ over \mathcal{X}_k relative to \mathcal{D}_k extends to a finite étale morphism $\mathcal{U}' \rightarrow \mathcal{U}$, unique up to unique isomorphism, and the generic fiber of \mathcal{U}' is connected.

Let $\overline{D}' = \overline{\varphi}^{-1}(\mathcal{D}_k)$ and let $\overline{U}' = C' \setminus \overline{D}'$, so $\overline{U}' \rightarrow \mathcal{U}_k$ is a tamely ramified cover of \mathcal{U}_k over \mathcal{X}_k relative to \mathcal{D}_k . Let $\varphi : \mathcal{U}' \rightarrow \mathcal{U}$ be the unique étale covering extending $\overline{\varphi}|_{\overline{U}'}$; this is equipped with an isomorphism $\mathcal{U}'_k \cong \overline{U}'$ identifying $\overline{\varphi}|_{\overline{U}'}$ with φ_k . Let $U = \mathcal{U}_K = X \setminus D$, let $U' = \mathcal{U}'_K$, let X' be the smooth compactification of U' , and let $\varphi_K : X' \rightarrow X$ denote the unique morphism extending $\varphi_K : U' \rightarrow U$. We will show that there is a simple neighborhood W of y in X^{an} such that $\varphi_K^{-1}(W) \rightarrow W$ is a tame covering, and conclude that a lifting exists using Corollary 6.16.

6.20.1. First we claim that $\varphi_K^{-1}(y)$ consists of a unique point $y' \in X'^{\text{an}}$. Letting \mathfrak{U} and \mathfrak{U}' denote the ϖ -adic completions of \mathcal{U} and \mathcal{U}' , respectively, we have a Cartesian square

$$\begin{array}{ccc}
 \mathfrak{U}'_K & \xrightarrow{i_{\mathfrak{U}'}} & U'^{\text{an}} \\
 \downarrow & & \downarrow \\
 \mathfrak{U}_K & \xrightarrow{i_{\mathfrak{U}}} & U^{\text{an}}
 \end{array}$$

by Lemma 6.19. In other words, \mathfrak{U}'_K is the inverse image of \mathfrak{U}_K under $\varphi_K : U'^{\text{an}} \rightarrow U^{\text{an}}$. Since the reduction map $\mathfrak{U}_K \rightarrow \mathfrak{U}_k = \mathcal{U}_k$ is surjective and y is the unique point of X^{an} reducing to the generic point of $\mathfrak{U}_k \subset \mathfrak{X}_k$, we have $y \in \mathfrak{U}_K$, so it suffices to show that there is a unique point $y' \in \mathfrak{U}'_K$ mapping to y . It follows easily from the functoriality of the reduction map that the only point y' mapping to y is the unique point of \mathfrak{U}'_K reducing to the generic point of \mathfrak{U}'_k . In particular, the residue curve of X'^{an} at y' is identified with C' , which is the smooth completion of $\mathfrak{U}'_k = \overline{U}'$.

6.20.2. Choose a strongly semistable vertex set V' of X' containing y' , let $\Sigma' = \Sigma(X', V')$, and let $\tau : X'^{\text{an}} \rightarrow \Sigma'$ be the retraction. Let $B \subset X^{\text{an}}$ be the formal fiber of a point $\bar{x} \in \mathfrak{U}_k(k)$; equivalently, B is a connected component of $X^{\text{an}} \setminus \{y\}$ not meeting D . Let B' be a connected component of

$X'^{\text{an}} \setminus \{y'\}$ contained in $\varphi_K^{-1}(B)$. We have $B \subset \mathfrak{U}_K$; therefore $B' \subset \mathfrak{U}'_K$, so B' is a connected component of $\mathfrak{U}'_K \setminus \{y'\}$, hence B' is the formal fiber of a point $\bar{x}' \in \mathfrak{U}'_k(k)$, so $B' \cong \mathbf{B}(1)_+$. It follows that $V' \setminus (V' \cap B')$ is again a semistable vertex set, so we may and do assume that $\Sigma' \cap B' = \emptyset$ for all such B' .

Let $e' \subset \Sigma'$ be an edge adjacent to y' and let B be the connected component of $X^{\text{an}} \setminus \{y\}$ containing $\varphi_K(e^\circ)$. By the above, B is the formal fiber of a point $\bar{x} \in \mathcal{D}_k$. Fix an isomorphism $B \cong \mathbf{B}(1)_+$ taking the unique point in $D \cap B$ to 0. Let $v' \in T_{y'}$ be the tangent vector in the direction of e' and let $v = d\varphi_K(y')(v') \in T_y$. The point in $\mathfrak{X}_k(k)$ corresponding to v is \bar{x} ; let $\bar{x}' \in \bar{D}' \subset C'(k)$ be the point corresponding to v' , so $\bar{\varphi}(\bar{x}') = \bar{x}$. By Theorem 4.23, the ramification degree δ of $\bar{\varphi}$ at \bar{x}' is equal to $d_{v'}\varphi_K(x')$.

Let B' be the connected component of $X'^{\text{an}} \setminus \{y'\}$ containing e'° . Viewing $\varphi_K|_{B'}$ as a morphism $B' \rightarrow \mathbf{B}(1)_+ \subset \mathbf{A}_K^{1,\text{an}}$ via our chosen isomorphism $B \cong \mathbf{B}(1)_+$, the map $x' \mapsto \log|\varphi_K(x')|$ is a piecewise affine function on e'° which changes slope on the retractions of the zeros of φ_K . Shrinking e' if necessary, we may and do assume that $\log|\varphi_K|$ is affine on e'° ; its slope has absolute value δ , and $\log|\varphi_K(x')| \rightarrow 0$ as $x' \rightarrow y'$. By (4.24) the restriction of φ_K to the open annulus $\tau^{-1}(e'^\circ)$ is a finite morphism of degree δ onto an open annulus $\mathbf{S}(a)_+ \subset \mathbf{B}(1)_+$, where $\text{val}(a)$ is δ times the length of e'° .

6.20.3. Enlarging Σ' if necessary, we may and do assume that every tangent direction $v' \in T_{y'}$ corresponding to a point in \bar{D}' is represented by an edge in Σ' . Let e'_1, \dots, e'_r be the edges of Σ' adjacent to y' , for $i = 1, \dots, r$ let B_i be the connected component of $X^{\text{an}} \setminus \{y\}$ containing $\varphi_K(e_i^\circ)$, and choose isomorphisms $B_i \xrightarrow{\sim} \mathbf{B}(1)_+$ sending the unique point of $D \cap B_i$ to 0. (The balls B_1, \dots, B_r are not necessarily distinct; we mean that one should choose a single isomorphism for each distinct ball.) Let $\bar{x}'_i \in \bar{D}'$ be the point corresponding to the tangent direction at y' in the direction of e'_i , and let δ_i be the ramification degree of $\bar{\varphi}$ at \bar{x}'_i . Applying the procedure of (6.20.2) for each e'_i , and shrinking if necessary, we may and do assume that for every i , φ_K induces a degree- δ_i morphism of $\tau^{-1}(e_i^\circ)$ onto an open annulus $\mathbf{S}(a)_+ \subset \mathbf{B}(1)_+ \cong B_i$, with a independent of i .

Let $W \subset X^{\text{an}}$ be the union of $\tau^{-1}(y)$ with the annuli $\mathbf{S}(a)_+ \subset B_i$, so W is obtained from X^{an} by removing a closed ball around each point of D . This is a simple neighborhood of y in X^{an} , hence is a star-shaped curve. Let $\Sigma'_0 = \{y'\} \cup \bigcup_{i=1}^r e_i^\circ$ and let $W' = \tau^{-1}(\Sigma'_0)$, so W' is a simple neighborhood of y' in X'^{an} and is hence a star-shaped curve. We claim that $W' = \varphi_K^{-1}(W)$. Clearly $\varphi_K(W') \subset W$, so it suffices to show that the fibers of φ_K have length equal to $\deg(\varphi_K)$. This is certainly the case for $\varphi_K^{-1}(y) = \{y'\}$. If $B \subset X^{\text{an}}$ is the formal fiber of a point in $\mathfrak{U}_k(k)$, then as in the proof of Proposition 6.15(1), $\varphi_K^{-1}(B) \subset W'$ is a disjoint union of $\deg(\varphi_K)$ open balls mapping isomorphically onto B . For each i the inverse image of $\mathbf{S}(a)_+ \subset \mathbf{B}(1)_+ \cong B_i$ in W' is a disjoint union of annuli, one for each point in the fiber of $\bar{\varphi}$ containing \bar{x}'_i , and the degree of φ_K restricted to each annulus is the ramification degree of $\bar{\varphi}$ at the corresponding point. The sum of the ramification degrees of $\bar{\varphi}$ at the points in any fiber is equal to $\deg(\bar{\varphi}) = \deg(\varphi)$, which proves the claim. Therefore φ_K induces a finite morphism $W' \rightarrow W$, which is a tame covering of (W, y) (really of (W, y, \emptyset)).

By construction, (Y, y) is a compactification of (W, y) , so by Corollary 6.16, the tame covering $W' \rightarrow W$ lifts to a tame covering $Y' \rightarrow Y$ relative to D . It is clear that $(Y', y', \varphi_K^{-1}(D))$ is a lifting of C' to a punctured star-shaped curve over (Y, y, D) . (One can show that in fact $Y' \cong X'^{\text{an}}$, although this is not clear *a priori*.)

6.20.4. It remains to prove (still in the case when Y is proper) that liftings are unique up to unique isomorphism. Let $\varphi_K : (Y', y', D') \rightarrow (Y, y, D)$ and $\varphi'_K : (Y'', y'', D'') \rightarrow (Y, y, D)$ be two liftings of C' to punctured star-shaped curves over (Y, y, D) . Then Y' and Y'' are also proper, hence we may and do assume that they are the analytic generic fibers of smooth, proper, connected formal curves \mathfrak{X}' and \mathfrak{X}'' as in (6.20). Since $\varphi_K(y') = \varphi'_K(y'') = y$, there are unique finite morphisms $\varphi : \mathfrak{X}' \rightarrow \mathfrak{X}$ and $\varphi' : \mathfrak{X}'' \rightarrow \mathfrak{X}$ extending φ_K and φ'_K , respectively, and φ_K and φ'_K are identified with $\bar{\varphi}$ under the isomorphisms $C_{y'} = \mathfrak{X}'_k \cong C'$ and $C_{y''} = \mathfrak{X}''_k \cong C'$. Let \mathcal{X}' and \mathcal{X}'' be the algebraizations of \mathfrak{X}' and \mathfrak{X}'' , respectively, and let $\mathcal{U}' = \mathcal{X}' \setminus \varphi^{-1}(D)$ and $\mathcal{U}'' = \mathcal{X}'' \setminus \varphi'^{-1}(D)$. Then $\mathcal{U}', \mathcal{U}''$ are finite étale coverings of \mathcal{U} lifting $\bar{U}' \rightarrow \mathcal{U}_k$, so there is a unique isomorphism $\psi : \mathcal{U}' \xrightarrow{\sim} \mathcal{U}''$ over \mathcal{U} . Since Y' (resp. Y'') is the

analytification of the smooth completion of \mathcal{U}'_K (resp. \mathcal{U}''_K), ψ_K extends uniquely to a Y -isomorphism $\psi_K : Y' \xrightarrow{\sim} Y''$; we have $\psi_K(y') = y''$, and $\psi_{y'} : C_{y'} \xrightarrow{\sim} C_{y''}$ is a C_y -isomorphism because $\mathcal{U}'_k \xrightarrow{\sim} \mathcal{U}''_k$ is a \mathcal{U}_k -isomorphism. Hence ψ_K is an isomorphism of liftings.

As for uniqueness, suppose that $\psi_K : (Y', y') \rightarrow (Y', y')$ is an automorphism of Y' as a lifting of C' . Then ψ_K extends uniquely to an \mathcal{X} -automorphism $\psi : \mathcal{X}' \xrightarrow{\sim} \mathcal{X}'$ that is the identity on \mathcal{X}'_k , which restricts to a \mathcal{U} -automorphism of \mathcal{U}' that is the identity on \mathcal{U}'_k . It follows from the uniqueness of \mathcal{U}' up to unique isomorphism that ψ_K is the identity when restricted to \mathcal{U}'_K , so since Y' is the analytification of the smooth completion of \mathcal{U}'_K , the automorphism ψ_K is the identity. This concludes the proof for proper Y .

6.20.5. Now suppose that (Y, y) is not proper. Let $Y \hookrightarrow X$ be a compactification (6.6) of Y ; we will identify Y with its image in X . Let D_1 be the union of D with a choice of one K -point from every connected component of $X \setminus Y$. Let $\varphi : (X', y', D'_1) \rightarrow (X, y, D_1)$ be a lifting of C' to a punctured star-shaped curve over (X, y, D_1) and let $Y' = \varphi^{-1}(Y)$. Then $(Y', y', \varphi^{-1}(D))$ is a lifting of C' to a punctured star-shaped curve over (Y, y, D) .

Let $\psi : Y' \rightarrow Y'$ be an automorphism of Y' as a lifting of C' . Since ψ induces the identity map $C_{y'} \rightarrow C_{y'}$ on residue curves, ψ takes each connected component of $Y' \setminus \{y'\}$ to itself. Recall that $X' \setminus Y'$ consists of a disjoint union of closed balls around the points of $D'_1 \setminus D'$, where $D'_1 = \varphi^{-1}(D_1)$. Let $x' \in D'_1 \setminus D'$, let B' be the connected component of $X' \setminus \{y'\}$ containing x' , and let $B = \varphi(B') \subset X$. By Proposition 6.15(2), we can choose isomorphisms $B \cong \mathbf{B}(1)_+$ and $B' \cong \mathbf{B}(1)_+$ sending x' and $\varphi(x')$ to 0, such that the composition

$$\mathbf{B}(1)_+ \cong B' \xrightarrow{\varphi} B \cong \mathbf{B}(1)_+$$

is of the form $t \mapsto t^\delta$. Let $A' = Y' \cap B'$ and let $A = Y \cap B = \varphi(A')$. The isomorphism $B \cong \mathbf{B}(1)_+$ (resp. $B' \cong \mathbf{B}(1)_+$) identifies A (resp. A') with an open annulus $\mathbf{S}(a)_+$ (resp. $\mathbf{S}(a')_+$) in $\mathbf{B}(1)_+$. Since $\psi|_{A'} : A' \rightarrow A'$ is an A -morphism, the composition $\mathbf{S}(a')_+ \cong A' \rightarrow A' \cong \mathbf{S}(a')_+$ is of the form $t \mapsto \zeta t$, where $\zeta \in R^\times$ is a δ th root of unity. Therefore $\psi|_{A'}$ extends uniquely to a B -morphism $\psi|_{B'} : B' \rightarrow B'$ fixing 0. Gluing these morphisms together, we obtain an X -morphism $\psi : X' \rightarrow X'$ extending $\psi : Y' \rightarrow Y'$. By construction this is an automorphism of X' as a lifting of C' to X , which is thus the identity. Therefore $\psi : Y' \rightarrow Y'$ is the identity.

Let $\varphi' : (Y'', y'', D'') \rightarrow (Y, y, D)$ be another lifting of C' to a punctured star-shaped curve over (Y, y, D) . By Corollary 6.16, there exists a compactification $Y'' \hookrightarrow X''$ of Y'' and an extension of φ' to a tame covering $\varphi' : (X'', y'', \varphi'^{-1}(D_1)) \rightarrow (X, y, D_1)$. This is another lifting of C' to a punctured star-shaped curve over (X, y, D_1) , so there is an isomorphism $\psi : (X', y') \xrightarrow{\sim} (X'', y'')$ of liftings. Since $Y' = \varphi^{-1}(Y)$ and $Y'' = \varphi'^{-1}(Y)$, the isomorphism ψ restricts to an isomorphism $(Y', y') \xrightarrow{\sim} (Y'', y'')$ of liftings. ■

Corollary 6.21. *With the notation in Theorem 6.18, let $(Y', y', D') \rightarrow (Y, y, D)$ be a lifting of C' to a punctured star-shaped curve over (Y, y, D) . Then the natural homomorphism*

$$\mathrm{Aut}_Y(Y') \longrightarrow \mathrm{Aut}_{C_y}(C')$$

is bijective. If $(Y'', y'', D'') \rightarrow (Y, y, D)$ is a second lifting of C' to a punctured star-shaped curve over (Y, y, D) then the natural map

$$\mathrm{Isom}_Y(Y'', Y') \longrightarrow \mathrm{Isom}_{C_y}(C_{y''}, C_{y'})$$

is bijective.

7. CLASSIFICATION OF LIFTINGS OF HARMONIC MORPHISMS OF METRIZED COMPLEXES

Fix a triangulated punctured curve $(X, V \cup D)$ with skeleton $\Sigma = \Sigma(X, V \cup D)$ and let $\tau : X^{\mathrm{an}} \rightarrow \Sigma$ be the canonical retraction. Throughout this section, we **assume that Σ has no loop edges**. Let $\varphi : \Sigma' \rightarrow \Sigma$ be a tame covering of metrized complexes of curves. A *lifting of Σ' to a tame covering of $(X, V \cup D)$* is a tame covering of triangulated punctured curves $\varphi : (X', V' \cup D') \rightarrow (X, V \cup D)$

(see (4.30)) equipped with a Σ -isomorphism $\varphi^{-1}(\Sigma) \cong \Sigma'$ of metrized complexes of curves. Since $V' = \varphi^{-1}(V)$ and $D' = \varphi^{-1}(D)$, we will often denote a lifting simply by X' . Let $\varphi : X' \rightarrow X$ and $\varphi' : X'' \rightarrow X$ be two liftings of Σ' and let $\psi : X' \xrightarrow{\sim} X''$ be an X -isomorphism of curves. Then ψ restricts to a Σ -automorphism

$$\psi|_{\Sigma'} : \Sigma' \cong \varphi^{-1}(\Sigma) \xrightarrow{\sim} \varphi'^{-1}(\Sigma) \cong \Sigma'.$$

We will consider liftings up to X -isomorphism preserving Σ' , i.e. such that $\psi_{\Sigma'}$ is the identity, and we will also consider liftings up to isomorphism as curves over X .

7.1. For every finite vertex $x \in V$, let $\Sigma(x)$ be the connected component of x in $\{x\} \cup (\Sigma \setminus V)$, let $Y(x) = \tau^{-1}(\Sigma(x))$, and let $D(x) = D \cap Y(x)$, as in Example 6.14. Then $(Y(x), x, D(x))$ is a punctured star-shaped curve. By Proposition 2.22, for every finite vertex $x' \in V(\Sigma')$ lying above x the morphism $\varphi_{x'} : C_{x'} \rightarrow C_x$ is finite, tamely ramified, and branched only over the points of C_x corresponding to tangent directions at x represented by edges of $\Sigma(x)$. Let $\psi(x') : (Y'(x'), x', D(x')) \rightarrow (Y(x), x, D(x))$ be the unique lifting of $C_{x'}$ to a punctured star-shaped curve over $(Y(x), x, D(x))$ provided by Theorem 6.18 and let $\Sigma'(x') = \psi(x')^{-1}(\Sigma(x)) = \Sigma(Y', \{x'\} \cup D'(x'))$. Then $\Sigma'(x')$ is canonically identified with the connected component of x' in $\{x'\} \cup (\Sigma' \setminus V_f(\Sigma'))$ in such a way that for every edge e' of $\Sigma'(x')$, the point $\text{red}_{x'}(e') \in C_{x'}(k)$ is identified with the point $\text{red}(e')$ defined in (6.7). This induces an identification of $D'(x')$ with $\varphi^{-1}(D) \cap \Sigma'(x')$. Let $\tau_{x'}$ be the canonical retraction $Y'(x') \rightarrow \Sigma'(x')$ defined in (6.3).

7.2. Let $e' \in E_f(\Sigma')$ and $e = \varphi(e')$. Choose $a, a' \in K^\times$ with $\text{val}(a) = \ell(e)$ and $\text{val}(a') = \ell(e')$. Let x', y' be the endpoints of e' . By Proposition 6.15(3), we can choose isomorphisms $\tau_{x'}^{-1}(e'^{\circ}) \cong \mathbf{S}(a)_+$ and $\tau_{y'}^{-1}(e'^{\circ}) \cong \mathbf{S}(a')_+$ in such a way that the finite morphisms $\tau_{x'}^{-1}(e'^{\circ}) \rightarrow \tau^{-1}(e^{\circ})$ and $\tau_{y'}^{-1}(e'^{\circ}) \rightarrow \tau^{-1}(e^{\circ})$ are given by $t \mapsto t^{d_{e'}(\varphi)}$, where $d_{e'}(\varphi)$ is the degree of the edge map $e' \rightarrow e$. In particular, there exists a $\tau^{-1}(e^{\circ})$ -isomorphism $\tau_{x'}^{-1}(e'^{\circ}) \xrightarrow{\sim} \tau_{y'}^{-1}(e'^{\circ})$. As explained in (6.11), there are canonical identifications

$$\text{Aut}_{\tau^{-1}(e^{\circ})}(d_{e'}(\varphi)) := \text{Aut}_{\tau^{-1}(e^{\circ})}(\tau_{x'}^{-1}(e'^{\circ})) = \text{Aut}_{\tau^{-1}(e^{\circ})}(\tau_{y'}^{-1}(e'^{\circ})) \cong \mathbf{Z}/d_{e'}(\varphi)\mathbf{Z};$$

hence the set of isomorphisms $\tau_{x'}^{-1}(e'^{\circ}) \xrightarrow{\sim} \tau_{y'}^{-1}(e'^{\circ})$ is a principal homogeneous space under the pre- or post-composition action of $\text{Aut}_{\tau^{-1}(e^{\circ})}(d_{e'}(\varphi))$.

Let $E_f^{\pm}(\Sigma')$ denote the set of oriented finite edges of Σ' , and for $e' \in E_f^{\pm}(\Sigma')$ let \bar{e}' denote the same edge with the opposite orientation. Let $\mathcal{G}(\Sigma', X)$ denote the set of tuples $(\Theta_{e'})_{e' \in E_f^{\pm}(\Sigma')}$ of isomorphisms

$$(7.2.1) \quad \Theta_{e'} : \tau_{x'}^{-1}(e'^{\circ}) \xrightarrow{\sim} \tau_{y'}^{-1}(e'^{\circ})$$

such that $\Theta_{\bar{e}'} = \Theta_{e'}^{-1}$, where $e' = \overrightarrow{x'y'}$. We call $\mathcal{G}(\Sigma', X)$ the set of *gluing data* for a lifting of Σ' to a tame covering of $(X, V \cup D)$, and we emphasize that $\mathcal{G}(\Sigma', X)$ is *nonempty*.

7.3. Let $\alpha \in \text{Aut}_{\Sigma}(\Sigma')$, so α is a degree-1 finite harmonic morphism $\Sigma' \xrightarrow{\sim} \Sigma'$ preserving $\Sigma' \rightarrow \Sigma$. Let $x' \in V_f(\Sigma')$ and let $x'' = \alpha(x')$ and $x = \varphi(x') = \varphi(x'')$. Part of the data of α is a C_x -isomorphism $\alpha_{x'} : C_{x'} \xrightarrow{\sim} C_{x''}$. By Corollary 6.21 there is a unique lift of $\alpha_{x'}$ to a $Y(x)$ -isomorphism $Y'(x') \xrightarrow{\sim} Y'(x'')$ of punctured star-shaped curves inducing the isomorphism $C_{x'} \xrightarrow{\sim} C_{x''}$ on residue curves. Let $e' \in E_f(\Sigma')$ be an edge adjacent to x' , let $e'' = \alpha(e')$, and let $e = \varphi(e') = \varphi(e'')$. Then $\alpha_{x'}$ restricts to an isomorphism

$$\alpha_{x'} : \tau_{x'}^{-1}(e'^{\circ}) \xrightarrow{\sim} \tau_{x''}^{-1}(e''^{\circ}).$$

Define the *conjugation action* of $\text{Aut}_{\Sigma}(\Sigma')$ on $\mathcal{G}(\Sigma', X)$ by the rule

$$(7.3.1) \quad \alpha \cdot (\Theta_{e'})_{e' \in E_f^{\pm}(\Sigma')} = (\alpha_{y'}^{-1} \circ \Theta_{\alpha(e')} \circ \alpha_{x'})_{e' \in E_f^{\pm}(\Sigma')}$$

where $e' = \overrightarrow{x'y'}$.

Theorem 7.4. (Classification of lifts of harmonic morphisms) *Let $(X, V \cup D)$ be a triangulated punctured curve with skeleton $\Sigma = \Sigma(X, V \cup D)$. Assume that Σ has no loop edges. Let $\varphi : \Sigma' \rightarrow \Sigma$ be a tame covering of metrized complexes of curves.*

- (1) *There is a canonical bijection between the set of gluing data $\mathcal{G}(\Sigma', X)$ and the set of liftings of Σ' to a tame covering of $(X, V \cup D)$, up to X -isomorphism preserving Σ' . In particular, there exists a lifting of Σ' . Any such lifting has no nontrivial automorphisms which preserve Σ' .*
- (2) *Two tuples of gluing data determine X -isomorphic curves if and only if they are in the same orbit under the conjugation action (7.3.1). The stabilizer in $\text{Aut}_\Sigma(\Sigma')$ of an element of $\mathcal{G}(\Sigma', X)$ is canonically isomorphic to the X -automorphism group of the associated curve.*

Proof. Given $(\Theta_{e'}) \in \mathcal{G}(\Sigma', X)$ one can glue the local lifts $\{Y(x')\}_{x' \in V_f(\Sigma')}$ via the isomorphisms (7.2.1) to obtain an analytic space which one easily verifies is smooth and proper, hence arises as the analytification of an algebraic curve X' . Moreover the morphisms $Y'(x') \rightarrow Y(x)$ glue to give a morphism $X'^{\text{an}} \rightarrow X^{\text{an}}$, which is the analytification of a morphism $\varphi : X' \rightarrow X$. By construction, if $V' = \varphi^{-1}(V)$ and $D' = \varphi^{-1}(D)$, then $(X', V' \cup D') \rightarrow (X, V \cup D)$ is a lifting of Σ' to a tame covering of $(X, V \cup D)$.

Now let $\varphi : (X', V' \cup D') \rightarrow (X, V \cup D)$ be any lifting of Σ' to a tame covering of $(X, V \cup D)$. As explained in Example 6.14, for every $x \in V$ the inverse image $\varphi^{-1}(Y(x))$ is a disjoint union of tame covers of the punctured star-shaped curve $(Y(x), x, D(x))$, one for each $x' \in \varphi^{-1}(x)$. By Theorem 6.18, we have canonical identification $\varphi^{-1}(Y(x)) = \coprod_{x' \mapsto x} Y(x')$. For $x', y' \in V'$ we have $Y(x') \cap Y(y') = \coprod_{e' \ni x', y'} \tau^{-1}(e'^{\circ})$, where $\tau : X'^{\text{an}} \rightarrow \Sigma'$ is the retraction. Hence X'^{an} is obtained by pasting the local liftings $Y(x')$ via a choice of isomorphisms $(\Theta_{e'}) \in \mathcal{G}(\Sigma', X)$.

Let $\psi : X' \xrightarrow{\sim} X'$ be an X -automorphism preserving Σ' . Since ψ is the identity on the set $\Sigma' \subset X'^{\text{an}}$, for $x' \in V'$ we have $\psi(Y(x')) = Y(x')$. We also have that for $x' \in V'$ the map of residue curves $\psi_{x'} : C_{x'} \rightarrow C_{x'}$ is the identity, so by Corollary 6.21, ψ restricts to the identity morphism $Y(x') \rightarrow Y(x')$. Since $X'^{\text{an}} = \bigcup_{x' \in V'} Y(x')$ we have that ψ is the identity. It follows from this that the tuple $(\Theta_{e'})$ can be recovered from the class of X' modulo X -isomorphisms preserving Σ' , so different gluing data give rise to curves which are not equivalent under such isomorphisms. This proves (1).

Let $\varphi : (X', V' \cup D') \rightarrow (X, V \cup D)$ and $\varphi'' : (X'', V'' \cup D'') \rightarrow (X, V \cup D)$ be the liftings of Σ' associated to the tuples of gluing data $(\Theta'_{e'}), (\Theta''_{e'}) \in \mathcal{G}(\Sigma', X)$, respectively. Suppose that there exists $\alpha \in \text{Aut}_\Sigma(\Sigma')$ such that $\alpha \cdot (\Theta'_{e'}) = (\Theta''_{e'})$. Then for all $e' = \overrightarrow{x'y'} \in E_f^\pm(\Sigma')$ we have

$$\Theta''_{\alpha(e')} \circ \alpha_{x'} = \alpha_{y'} \circ \Theta'_{e'},$$

so the $Y(\varphi(x'))$ -isomorphisms $\alpha_{x'} : Y(x') \xrightarrow{\sim} Y(\alpha(x'))$ glue to give an X -isomorphism $\alpha : X' \xrightarrow{\sim} X''$. Conversely, the restriction of an X -isomorphism $\alpha : X' \xrightarrow{\sim} X''$ to Σ' is a Σ -automorphism of Σ' . It is easy to see that these are inverse constructions. Taking $X' = X''$ we have an injective homomorphism $\text{Aut}_X(X') \hookrightarrow \text{Aut}_\Sigma(\Sigma')$; it follows formally from the above considerations that its image is the stabilizer of $(\Theta'_{e'})$. \blacksquare

Remark 7.5. Let X' be a lifting of Σ' to a tame covering of $(X, V \cup D)$. It follows from Theorem 7.4(1) that the natural homomorphism

$$\text{Aut}_X(X') \longrightarrow \text{Aut}_\Sigma(\Sigma')$$

is injective. It is not in general surjective: see Example 7.8 (where all the stabilizer groups of elements of $\mathcal{G}(\Sigma', X)$ are proper subgroups of $\text{Aut}_\Sigma(\Sigma')$).

Remark 7.6. It is worth mentioning that the question of lifting (and classification of all possible liftings) in the wildly ramified case is more subtle and cannot be guaranteed in general. Some lifting results in the wildly ramified case are known: e.g., Liu proves in [Liu03, Proposition 5.4] that any finite surjective generically étale admissible cover of proper semistable curves over an algebraically closed field k lifts to a finite morphism of smooth proper curves over K . However, the same statement for metrized complexes cannot be true in general: consider the metrized complex \mathcal{C} consisting of a single finite vertex v and an infinite vertex u attached to v by an infinite edge e with $C_v \cong \mathbf{P}_k^1$ and

$\text{red}_v(e) = \infty$. Consider the degree p generically étale morphism of metrized complexes $\mathcal{C} \rightarrow \mathcal{C}$ which restricts to a degree p Artin-Schreier cover $\varphi : \mathbf{P}_k^1 \rightarrow \mathbf{P}_k^1$, étale over $\mathbf{A}_k^1 \rightarrow \mathbf{A}_k^1$ and ramified at ∞ (with ramification index p), and which is linear of slope p on the infinite edge e . This map cannot be lifted to a degree p cover $\mathbf{P}_K^1 \rightarrow \mathbf{P}_K^1$ in characteristic zero, otherwise, we would obtain a (connected) étale degree p cover of \mathbf{A}_K^1 which is impossible since \mathbf{A}_K^1 is simply connected.

We now state a variant of Theorem 7.4 in which we allow loop edges.

Theorem 7.7. *Let $(X, V \cup D)$ be a triangulated punctured K -curve, let $\Sigma = \Sigma(X, V \cup D)$, and let $\bar{\varphi} : \Sigma' \rightarrow \Sigma$ be a tame covering of Λ -metrized complexes of k -curves. Then there exists a tame covering $\varphi : (X', V' \cup D') \rightarrow (X, V \cup D)$ of triangulated punctured K -curves lifting $\bar{\varphi}$.*

Proof. The only issue is that in Theorem 7.7, we merely require that V is a semistable vertex set whereas in Theorem 7.4 we require it to be strongly semistable. However, we can modify Σ by inserting a (valence 2) vertex v along each loop edge, with $C_v \cong \mathbf{P}_k^1$, and with the marked points on C_v being 0 and ∞ . Call the resulting metrized complex $\tilde{\Sigma}$, and let \bar{V}_0 denote the set of vertices which have been added to the vertex set \bar{V} for Σ . We construct a new metrized complex $\tilde{\Sigma}'$ from Σ' by adding $\bar{V}'_0 := \bar{\varphi}^{-1}(\bar{V}_0)$ to the vertex set \bar{V}' for Σ' and letting $C_{v'} \cong \mathbf{P}_k^1$ for $v' \in \bar{V}'_0$, with the marked points on $C_{v'}$ being 0 and ∞ . We can extend the tame covering $\bar{\varphi} : \Sigma' \rightarrow \Sigma$ to a tame covering $\tilde{\Sigma}' \rightarrow \tilde{\Sigma}$ by letting the map from $C_{v'}$ to $C_{\bar{\varphi}(v')}$, for $v' \in \bar{V}'_0$, be $z \mapsto z^d$, where d is the local degree of $\bar{\varphi}$ along the loop edge corresponding to v' . By Theorem 7.4, there is a tame covering $(X', V' \cup V'_0 \cup D') \rightarrow (X, V \cup V_0 \cup D)$ lifting $\tilde{\Sigma}' \rightarrow \tilde{\Sigma}$, where V'_0 (resp. V_0) corresponds to \bar{V}'_0 (resp. \bar{V}_0). Removing the vertices in V'_0 and V_0 gives a tame covering $(X', V' \cup D') \rightarrow (X, V \cup D)$ lifting $\bar{\varphi}$. \blacksquare

Example 7.8. In this example we suppose that $\text{char}(k) \neq 2$. Let E be a Tate curve over K , let Σ be its (set-theoretic) skeleton, and let $\tau : E^{\text{an}} \rightarrow \Sigma$ be the canonical retraction. Let $U : \mathbf{G}_m^{\text{an}}/q^{\mathbf{Z}} \xrightarrow{\sim} E^{\text{an}}$ be the Tate uniformization of E^{an} . The 2-torsion subgroup of E is $U(\{\pm 1, \pm\sqrt{q}\})$; choose a square root of q , let $y = U(1)$ and $z = U(\sqrt{q})$, and let $V = \{y, z\} \subset \Sigma$. This is a semistable vertex set of E and $\Sigma = \Sigma(E, V)$ is the circle with circumference $\text{val}(q)$; the points y and z are antipodal on Σ . Let e_1, e_2 be the edges of Σ ; orient e_1 so that y is the source vertex and e_2 so that z is the source vertex. The residue curves C_y and C_z are both isomorphic to \mathbf{P}_k^1 ; fix isomorphisms $C_y \cong \mathbf{P}_k^1$ and $C_z \cong \mathbf{P}_k^1$ such that the tangent direction at a vertex in the direction of the outgoing (resp. incoming) edge corresponds to ∞ (resp. 0).

Let Σ' be a circle of circumference $\frac{1}{2} \text{val}(q)$, let $V' = \{y', z'\}$ be a pair of antipodal points on Σ' , and let $e'_1 = \overrightarrow{y'z'}$ and $e'_2 = \overrightarrow{z'y'}$ be the two edges of Σ' , with the indicated orientations. Enrich Σ' with the structure of a metrized complex of curves by setting $C_{y'} = C_{z'} = \mathbf{P}_k^1$ and letting the tangent direction at a vertex in the direction of the outgoing (resp. incoming) edge correspond to ∞ (resp. 0). Define a morphism $\varphi : \Sigma' \rightarrow \Sigma$ of metric graphs by

$$\varphi(y') = y, \quad \varphi(z') = z, \quad \varphi(e'_1) = e_1, \quad \varphi(e'_2) = e_2,$$

with both degrees equal to 2. See Figure 3.

Topologically, $\Sigma' \rightarrow \Sigma$ is a homeomorphism. Let $\varphi_{y'} : C_{y'} \rightarrow C_y$ and $\varphi_{z'} : C_{z'} \rightarrow C_z$ both be the morphism $t \mapsto t^2 : \mathbf{P}_k^1 \rightarrow \mathbf{P}_k^1$. This makes φ into a degree-2 tame covering of metrized complexes of curves.

For $x \in V$ the analytic space $Y(x)$ is an open annulus of logarithmic modulus $\text{val}(q)$; fix an isomorphism $Y(x) \cong \mathbf{S}(q)_+$ for each x . For $x' \in V'$ the analytic space $Y'(x')$ is an open annulus of logarithmic modulus $\frac{1}{2} \text{val}(q)$; fix isomorphisms $Y'(x') \cong \mathbf{S}(\sqrt{q})_+$ such that the morphisms $Y'(x') \rightarrow Y(\varphi(x'))$ are given by $t \mapsto t^2 : \mathbf{S}(\sqrt{q})_+ \rightarrow \mathbf{S}(q)_+$. For $e' \in E(\Sigma')$ we have $d_{\varphi}(e') = 2$ by definition. Hence $\mathcal{G}(\Sigma', E)$ has four elements, which we label $(\pm 1, \pm 1)$. By Theorem 7.4(1) there are four corresponding classes of liftings of Σ' to a tame covering of (E, V) up to isomorphism preserving Σ' .

Any Σ -automorphism of Σ' is the identity on the underlying topological space. Hence an element of $\text{Aut}_{\Sigma}(\Sigma')$ is a pair of automorphisms $(\psi_{y'}, \psi_{z'}) \in \text{Aut}_{C_y}(C_{y'}) \times \text{Aut}_{C_z}(C_{z'})$ such that $\psi_{y'}, \psi_{z'}$ fix the




FIGURE 3. This figure illustrates Example 7.8. The morphism φ is a homeomorphism of underlying sets but has a degree of 2 on e'_1 and e'_2 . The arrows on e_1, e_2, e'_1, e'_2 represent the chosen orientations and the indicated tangent vectors at y, z, y', z' represent the tangent direction corresponding to ∞ in the corresponding k -curve.

points $0, \infty \in \mathbf{P}_k^1$. Thus $\psi_{y'}, \psi_{z'} = \pm 1$, so $\text{Aut}_\Sigma(\Sigma') = \{\pm 1\} \times \{\pm 1\}$. For $x' = y', z'$ the automorphism $-1 : C_{x'} \xrightarrow{\sim} C_{x'}$ lifts to the automorphism $-1 : \mathbf{S}(\sqrt{q})_+ \xrightarrow{\sim} \mathbf{S}(\sqrt{q})_+$. Therefore the conjugation action of $\text{Aut}_\Sigma(\Sigma')$ on $\mathcal{G}(\Sigma', E)$ is given as follows:

$$\begin{aligned} (1, 1) \cdot (\pm 1, \pm 1) &= (-1, -1) \cdot (\pm 1, \pm 1) = (\pm 1, \pm 1) \\ (-1, 1) \cdot (\pm 1, \pm 1) &= (1, -1) \cdot (\pm 1, \pm 1) = -(\pm 1, \pm 1). \end{aligned}$$

By Theorem 7.4(2), there are two isomorphism classes of lifts of Σ' to a tame covering of (E, V) , and each such lift has two automorphisms.

These liftings can be described concretely as follows. Fix a square root of q , let E_\pm be the algebraization of the analytic elliptic curve $\mathbf{G}_m^{\text{an}}/(\pm\sqrt{q})^{\mathbf{Z}}$, and let $\psi_\pm : E_\pm \rightarrow E$ be the morphism $t \mapsto t^2$ on uniformizations. Let $\Sigma_\pm = \psi_\pm^{-1}(\Sigma)$. Then Σ_\pm is isomorphic to Σ' as a tame covering of Σ and E_\pm is a lifting of Σ' to a tame covering of (E, D) . The elliptic curves E_\pm are not isomorphic (as K -schemes) because they have different q -invariants, so they represent the two isomorphism classes of liftings of Σ' . The nontrivial automorphism of E_\pm is given by translating by the image of $-1 \in \mathbf{G}_m^{\text{an}}(K)$ (this is not a homomorphism). In fact, since $\psi_\pm : E_\pm \rightarrow E$ is an étale Galois cover of degree 2, this is the only nontrivial automorphism of E_\pm as an E -curve, so the homomorphism

$$\text{Aut}_E(E_\pm) \longrightarrow \text{Aut}_\Sigma(\Sigma') \cong \{\pm 1\} \times \{\pm 1\}$$

is injective but not surjective (its image is $\{\pm(1, 1)\}$).

7.9. Consider now the automorphism group $\text{Aut}_\Sigma^0(\Sigma') \subset \text{Aut}_\Sigma(\Sigma')$ consisting of all degree-1 finite harmonic morphisms $\alpha : \Sigma' \rightarrow \Sigma'$ respecting $\varphi : \Sigma' \rightarrow \Sigma$ and inducing the identity on the metric graph Γ' underlying Σ' . The restriction of the conjugacy action of $\text{Aut}_\Sigma(\Sigma')$ on $\mathcal{G}(\Sigma', X)$ to the subgroup $\text{Aut}_\Sigma^0(\Sigma')$ admits a simplified description that we describe now. Combining this with arguments similar to those in the proof of Theorem 7.4 yields a classification of the set of liftings of Σ' up to isomorphism as liftings of the metric graph underlying Σ' ; see Theorem 7.10.

First, for each $x' \in V(\Sigma')$ with image $x = \varphi(x')$, let $\text{Aut}_{C_x}^0(C_{x'})$ be the subgroup of $\text{Aut}_{C_x}(C_{x'})$ that fixes every point of $C_{x'}$ of the form $\text{red}(e')$ for some edge e' of Σ' adjacent to x' . Then

$$\text{Aut}_\Sigma^0(\Sigma') = \prod_{x' \in V_f(\Sigma')} \text{Aut}_{C_{\varphi(x)}}^0(C_{x'}) =: \mathcal{E}^0.$$

Denote by \mathcal{E}^1 the finite abelian group

$$\mathcal{E}^1 = \prod_{e' \in E_f(\Sigma')} \text{Aut}_{\tau^{-1}(\varphi(e')^\circ)}(d_{e'}(\varphi)).$$

The discussion preceding Theorem 7.4 shows that the set of gluing data $\mathcal{G}(\Sigma', X)$ is canonically a principal homogeneous space under \mathcal{E}^1 .

The subgroup $\text{Aut}_{C_x}^0(C_{x'}) \subset \text{Aut}_{C_x}(C_{x'})$ corresponds to the subgroup $\text{Aut}_{Y(x)}^0(Y'(x'))$ of automorphisms in $\text{Aut}_{Y(x)}(Y'(x'))$ which act trivially on the skeleton $\Sigma'(x')$. Restriction of a $Y(x)$ -automorphism of $Y'(x')$ to $\tau_{x'}^{-1}(e'^{\circ})$ defines a homomorphism

$$\rho_{x',e'} : \text{Aut}_{C_x}^0(C_{x'}) = \text{Aut}_{Y(x)}^0(Y'(x')) \longrightarrow \text{Aut}_{\tau^{-1}(e^{\circ})}(d_{e'}(\varphi)).$$

Fix an orientation of each finite edge of Σ' . For $x' \in V_f(\Sigma')$ with image $x = \varphi(x')$, let $\rho_{x'} : \text{Aut}_{C_x}^0(C_{x'}) \rightarrow \mathcal{E}^1$ be the homomorphism whose e' -coordinate is

$$(\rho_{x'}(\alpha))_{e'} = \begin{cases} \rho_{x',e'}(\alpha) & \text{if } x' \text{ is the source vertex of } e' \\ \rho_{x',e'}(\alpha)^{-1} & \text{if } x' \text{ is the target vertex of } e' \\ 1 & \text{if } x' \text{ is not an endpoint of } e'. \end{cases}$$

Taking the product over all $x' \in V_f(\Sigma')$ yields a homomorphism $\rho : \mathcal{E}^0 \rightarrow \mathcal{E}^1$. The kernel and cokernel of ρ are independent of the choice of orientations. Viewing $\mathcal{E}^0 \rightarrow \mathcal{E}^1$ as a two-term complex \mathcal{E}^\bullet of groups, its cohomology groups are

$$H^0(\mathcal{E}^\bullet) = \ker(\rho) \quad \text{and} \quad H^1(\mathcal{E}^\bullet) = \text{coker}(\rho).$$

Theorem 7.10. *Let $(X, V \cup D)$ be a triangulated punctured curve with skeleton $\Sigma = \Sigma(X, V \cup D)$. Assume that Σ has no loop edges. Let $\varphi : \Sigma' \rightarrow \Sigma$ be a tame covering of metrized complexes of curves.*

- (1) $\mathcal{G}(\Sigma', X)$ is canonically a principal homogeneous space under \mathcal{E}^1 and the conjugacy action of $\text{Aut}_{\Sigma'}^0(\Sigma')$ on $\mathcal{G}(\Sigma', X)$ is given by the action of the subgroup $\rho(\text{Aut}_{\Sigma'}^0(\Sigma')) = \rho(\mathcal{E}^0) \subseteq \mathcal{E}^1$ on $\mathcal{G}(\Sigma', X)$.
- (2) The set of liftings of Σ' up to isomorphism as liftings of the metric graph underlying Σ' is a principal homogeneous space under $H^1(\mathcal{E}^\bullet)$, and the group of automorphisms of a given lifting as a lifting of the metric graph underlying Σ' is isomorphic to $H^0(\mathcal{E}^\bullet)$.

7.11. Descent to a general ground field. Let K_0 be a subfield of K , let X_0 be a smooth, projective, geometrically connected K_0 -curve, and let $D \subset X_0(K_0)$ be a finite set. Let $X = X_0 \otimes_{K_0} K$, let V be a strongly semistable vertex set of (X, D) , let $\Sigma = \Sigma(X, V \cup D)$, and let $\varphi : \Sigma' \rightarrow \Sigma$ be a tame covering of metrized complexes of curves, as in the statement of Theorem 7.4. Let $\varphi : X' \rightarrow X$ be a lifting of φ to a tame covering of $(X, V \cup D)$. Whereas we take the data of the morphism $\Sigma' \rightarrow \Sigma$ to be geometric, i.e. only defined over K , the covering $X' \rightarrow X$ is in fact defined over a finite, separable extension of K_0 . This follows from the fact that if $U = X \setminus D$ and $U' = X' \setminus \varphi^{-1}(D)$, then $\varphi : U' \rightarrow U$ is a tamely ramified cover of U over X relative to D (see Remarks 2.20(1) and 4.32(1)), along with the following lemma.

Lemma 7.12. *Let K_0 be any field, let K be a separably closed field containing K_0 , let X_0 be a smooth, projective, geometrically connected K_0 -curve, and let $D \subset X_0(K_0)$ be a finite set. Let $X = X_0 \otimes_{K_0} K$, let $\varphi : X' \rightarrow X$ be a finite morphism with X' smooth and (geometrically) connected, and suppose that φ is branched only over D , with all ramification degrees prime to the characteristic of K . Then there exists a finite, separable extension K_1 of K and a morphism $\varphi_1 : X'_1 \rightarrow X_0 \otimes_{K_0} K_1$ descending φ .*

Proof. Let $U_0 = X_0 \setminus D$, and let $U = X \setminus D$ and $U' = X' \setminus \varphi^{-1}(D)$. First suppose that K_0 is separably closed. By [SGA1, Exposé XIII, Corollaire 2.12], the tamely ramified étale fundamental groups $\pi_1^t(U_0)$ and $\pi_1^t(U)$ are isomorphic (with respect to some choice of base point). Since $\varphi : U' \rightarrow U$ is a tamely ramified cover of U over X relative to D , it is classified by a finite-index subgroup of $\pi_1^t(U) = \pi_1^t(U_0)$, so there exists a tamely ramified cover $\varphi_0 : U'_0 \rightarrow U_0$ of U_0 over X_0 relative to D descending φ .

Now we drop the hypothesis that K_0 is separably closed. By the previous paragraph we may assume that K is a separable closure of K_0 . By general principles the projective morphism $X' \rightarrow X$ descends to a subfield of K which is finitely generated (i.e. finite) over K_0 . ■

Remark 7.13. With the notation in (7.11), suppose that K_0 is a complete valued field with value group $\Lambda_0 = \text{val}(K_0^\times)$ and algebraically closed residue field k , that Σ is “rational over K_0 ” in that it comes from a (split) semistable formal R_0 -model of X_0 in the sense of Section 5, and that Σ' has

edge lengths contained in Λ_0 . With some extra work it is possible to carry out the gluing arguments of Theorem 7.7 directly over the field K_0 (in this context Lemma 6.10(1) still holds), which shows that the cover $X' \rightarrow X$ is in fact defined over K_0 . In the case of a discrete valuation this also follows from [Wew99], or from [Sai97] if $D = \emptyset$.

7.14. Liftings of tame harmonic morphisms. Theorem 7.7 implies the existing of liftings for (finite) tame harmonic morphisms of metrized complexes which are not necessarily tame coverings, the difference being generic étaleness. See Definition 2.21 for both definitions.

Proposition 7.15. *Let $(X, V \cup D)$ be a triangulated punctured K -curve, let $\Sigma = \Sigma(X, V \cup D)$, and let $\varphi : \Sigma' \rightarrow \Sigma$ be a tame harmonic morphism of metrized complexes of curves. Then there exists a triangulated punctured K -curve $(X', V' \cup D')$ and a finite morphism $\psi : (X', V' \cup D') \rightarrow (X, V \cup D)$ lifting φ .*

Proof. For any finite vertex x' of Σ' at which φ , seen as a morphism of augmented metric graphs, is ramified, let $q_1, \dots, q_r \in C_v$ be all the branch points of $\varphi_{x'}$ which do not correspond to any edge of Σ , and let p_{ij} denote the preimages of q_i under $\varphi_{x'}$. We modify Σ' and Σ by attaching infinite edges e_i to Σ at $x = \varphi(x')$ for each q_i , infinite edges e'_{ij} to Σ' at x' for each p_{ij} , and defining $\text{red}_x(e_i) = q_i$ and $\text{red}_{x'}(e'_{ij}) = p_{ij}$. Since each $\varphi_{x'}$ is tamely ramified, the harmonic morphism φ naturally extends to a tame covering $\tilde{\varphi} : \tilde{\Sigma}' \rightarrow \tilde{\Sigma}$ between the resulting modifications. Enlarging D to \tilde{D} by choosing points in $X(K) \setminus D$ with reduction q_i , we can assume that $\tilde{\Sigma} = \Sigma(X, V \cup \tilde{D})$. The result now follows from Theorem 7.7 by first lifting $\tilde{\varphi}$ to a tame covering $\psi : (X', V' \cup \tilde{D}') \rightarrow (X, V \cup \tilde{D})$ and then taking the restriction of ψ to $(X', V' \cup D') \rightarrow (X, V \cup D)$, where $D' = \psi^{-1}(D)$. \blacksquare

Remark 7.16. As mentioned above, in Proposition 7.15 we do not require that $\varphi : \Sigma' \rightarrow \Sigma$ be generically étale. This corresponds to not requiring that $D \subset X(K)$ contain the branch locus of the lift $\psi : X' \rightarrow X$: indeed, by Lemma 4.33, if D contains the branch locus then φ is a tame covering. In the situation of Proposition 7.15 the set of liftings of Σ' to a tame covering of $(X, V \cup D)$ can be infinite. For example, if $\Gamma' = \{p'\}$ and $\Gamma = \{p\}$ are both points and the morphism $\varphi_{p'} : C_{p'} \cong \mathbf{P}^1 \rightarrow C_p \cong \mathbf{P}^1$ is $z \mapsto z^2$, with (X, V) a minimal triangulation of \mathbf{P}^1 and $D = \emptyset$, then there are infinitely many such lifts, corresponding to the different ways of lifting the critical points and critical values of $\varphi_{p'}$ from k to K .

8. APPLICATION: COMPONENT GROUPS OF NÉRON MODELS

8.1. In contrast to the rest of the paper, we assume in this section that R is a complete *discrete valuation ring* with fraction field K and algebraically closed residue field k . In this case we take the value group Λ to be \mathbf{Z} .

There is a natural notion of harmonic 1-forms on a metric graph Γ of genus g (see [MZ08]). The space $\Omega^1(\Gamma)$ is a g -dimensional real vector space which can be canonically identified with $H^1(\Gamma, \mathbf{R})$, but we write elements of $\Omega^1(\Gamma)$ as $\omega = \sum_e \omega_e de$ as in [BF11], where the sum is over all edges with respect to a fixed vertex set for Γ . There is a canonical lattice $\Omega_{\mathbf{Z}}^1(\Gamma)$ of *integer harmonic 1-forms* inside $\Omega^1(\Gamma)$; these are the harmonic 1-forms for which every ω_e is an integer. A harmonic morphism $\varphi : \Gamma' \rightarrow \Gamma$ induces a natural pullback map on harmonic 1-forms via the formula

$$\varphi^*\left(\sum_e \omega_e de\right) = \sum_{e'} d_{e'}(\varphi) \cdot \omega_{\varphi(e')} de'.$$

Recall that a \mathbf{Z} -metric graph (i.e. a Λ -metric graph for $\Lambda = \mathbf{Z}$) with no infinite vertices is a (compact and finite) metric graph whose edge lengths are all positive integers (or equivalently, having a vertex set with respect to which all edge lengths are 1). If X/K is a smooth, proper, geometrically connected analytic curve and \mathfrak{X} is a semistable R -model for X , the skeleton $\Gamma_{\mathfrak{X}}$ of \mathfrak{X} is naturally a \mathbf{Z} -metric graph. Moreover, as we have seen earlier in this paper, a finite morphism of semistable models induces in a natural way a harmonic morphism of \mathbf{Z} -metric graphs.

Let Γ be a \mathbf{Z} -metric graph with no infinite vertices. We define the *regularized Jacobian* $\text{Jac}_{\text{reg}}(\Gamma)$ of Γ to be the group $\text{Jac}(G)$, where G is the “regular model” for Γ (the unique graph induced by a vertex set with all edge lengths equal to 1). The group $\text{Jac}(G)$ can be described explicitly as

$$\text{Jac}(G) = \Omega^1(G)^\# / H_1(G, \mathbf{Z}),$$

where $\Omega^1(G)^\#$ denotes the linear functionals $\Omega^1(G, \mathbf{R}) \rightarrow \mathbf{R}$ of the form \int_α with $\alpha \in C_1(G, \mathbf{Z})$ (see [BF11]). There is a canonical isomorphism between $\text{Jac}(G)$ and $\text{Pic}^0(G)$, the group of divisors of degree 0 on G modulo the principal divisors (see [BdlHN97]) as well as a canonical isomorphism

$$\text{Jac}(G) \cong H^1(G, \mathbf{Z})^\# / \Omega_{\mathbf{Z}}^1(G),$$

where

$$H^1(G, \mathbf{Z})^\# = \left\{ \omega \in \Omega^1(G, \mathbf{R}) : \int_\gamma \omega \in \mathbf{Z} \forall \gamma \in H_1(G, \mathbf{Z}) \right\}.$$

We recall (in our own terminology) the following result of Raynaud (cf. [Ray70] and [Bak08, Appendix A]):

Theorem 8.2. (Raynaud) *If X/K is a semistable curve, then the component group of the Néron model of $\text{Jac}(X)$ over R is canonically isomorphic to $\text{Jac}_{\text{reg}}(\Gamma_{\mathfrak{X}})$ (for any semistable model \mathfrak{X} of X).*

A harmonic morphism $\varphi : \Gamma' \rightarrow \Gamma$ of \mathbf{Z} -metric graphs induces in a functorial way homomorphisms $\varphi_* : \text{Jac}_{\text{reg}}(\Gamma') \rightarrow \text{Jac}_{\text{reg}}(\Gamma)$, defined by

$$\varphi_* \left(\left[\int_{\alpha'} \right] \right) = \left[\int_{\varphi_*(\alpha')} \right],$$

where

$$\varphi_* \left(\sum \alpha_{e'} e' \right) = \sum \alpha_{e'} \varphi(e'),$$

and $\varphi^* : \text{Jac}_{\text{reg}}(\Gamma) \rightarrow \text{Jac}_{\text{reg}}(\Gamma')$, defined by

$$\varphi^*([\omega]) = [\varphi^* \omega],$$

where

$$\varphi^* \left(\sum \omega_e de \right) = \sum_{e'} d_{e'}(\varphi) \omega_{\varphi(e')} de'.$$

We have the following elementary result, whose proof we omit:

Lemma 8.3. *Under the canonical isomorphism between $\text{Jac}(G)$ and $\text{Pic}^0(G)$, the homomorphism $\varphi_* : \text{Jac}(G') \rightarrow \text{Jac}(G)$ corresponds to the map $[D'] \mapsto [\varphi_*(D')]$ from $\text{Pic}^0(G')$ to $\text{Pic}^0(G)$, where $\varphi_*(\sum a_{v'}(v')) = \sum a_{v'} \varphi(v')$.*

The following is a “relative” version of Raynaud’s theorem; it implies that the covariant functor which takes a semistable R -model \mathfrak{X} for a K -curve X to the component group of the Néron model of the Jacobian of X factors as the “reduction graph functor” $\mathfrak{X} \mapsto \Gamma_{\mathfrak{X}}$ followed by the “regularized Jacobian functor” $\Gamma \mapsto \text{Jac}_{\text{reg}}(\Gamma)$. It is a straightforward consequence of the analytic description of Raynaud’s theorem given in [Bak08, Appendix A] (see also [BR13]):

Theorem 8.4. *If $f_K : X' \rightarrow X$ is a finite morphism of curves over K , the induced maps $f_* : \Phi_{J(X')} \rightarrow \Phi_{J(X)}$ and $f^* : \Phi_{J(X)} \rightarrow \Phi_{J(X')}$ on component groups coincide with the induced maps $\varphi_* : \text{Jac}_{\text{reg}}(\Gamma_{\mathfrak{X}'}) \rightarrow \text{Jac}_{\text{reg}}(\Gamma_{\mathfrak{X}})$ and $\varphi^* : \text{Jac}_{\text{reg}}(\Gamma_{\mathfrak{X}}) \rightarrow \text{Jac}_{\text{reg}}(\Gamma_{\mathfrak{X}'})$ for any morphism $f : \mathfrak{X}' \rightarrow \mathfrak{X}$ of semistable models extending f_K . (Here φ denotes the harmonic morphism of skeleta induced by f ; see Remark 5.14.)*

It follows easily from Lemma 8.3 and Theorem 8.4 that if $f_K : X' \rightarrow X$ is regularizable, i.e., if f_K extends to a morphism of regular semistable models, then the induced map $f_* : \Phi_{X'} \rightarrow \Phi_X$ on component groups is surjective. Thus whenever f_* is not surjective, it follows that f_K does not extend to a morphism of regular semistable models. One can obtain a number of concrete examples of this

situation from modular curves (e.g. the map $X_0(33) \rightarrow E$ over \mathbf{Q}_3^{nr} , where E is the optimal elliptic curve of level 33.)

Remark 8.5. One can show that if $\varphi : \Gamma' \rightarrow \Gamma$ is a harmonic morphism of \mathbf{Z} -metric graphs, then $\varphi_* : \Phi_{X'} \rightarrow \Phi_X$ is surjective iff $\varphi^* : \Phi_X \rightarrow \Phi_{X'}$ is injective. Indeed, it is not hard to check that the maps φ_* and φ^* are adjoint with respect to the *combinatorial monodromy pairing*, the non-degenerate symmetric bilinear form $\langle \cdot, \cdot \rangle : \text{Jac}(G) \times \text{Jac}(G) \rightarrow \mathbf{Q}/\mathbf{Z}$ defined by $\langle [\omega], [\int_\alpha] \rangle = [\int_\alpha \omega]$, where $\omega \in H_1(G, \mathbf{Z})^\#$ and $\int_\alpha \in \Omega^1(G)^\#$. Thus the groups $\ker(\varphi^*)$ and $\text{coker}(\varphi_*)$ are canonically dual. This is a combinatorial analogue of results proved by Grothendieck in SGA7 on the (usual) monodromy pairing.

As an application of Theorem 8.4 and our results on lifting harmonic morphisms, one can construct many examples of harmonic morphisms of \mathbf{Z} -metric graphs for which φ_* is not surjective. For example, consider the following question posed by Ken Ribet in a 2007 email correspondence with the second author (Baker):

Suppose $f : X' \rightarrow X$ is a finite morphism of semistable curves over a complete discretely valued field K with $g(X) \geq 2$. Assume that the special fiber of the minimal regular model of X' consists of two projective lines intersecting transversely. Is the induced map $f_* : \Phi_{X'} \rightarrow \Phi_X$ on component groups of Néron models necessarily surjective?

We now show that the answer to Ribet’s question is **no**.

Example 8.6. Consider the “banana graph” $B(\ell_1, \dots, \ell_{g+1})$ consisting of two vertices and $g + 1$ edges of length ℓ_i . This is the reduction graph of a semistable curve whose reduction has two \mathbf{P}^1 ’s crossing transversely at singular points of thickness $\ell_1, \dots, \ell_{g+1}$. If we set $G' = B(1, 1, 1, 1)$ and $G = B(1, 2, 2)$ and let Γ', Γ be the geometric realizations of G' and G , respectively, then there is a degree 2 harmonic morphism of \mathbf{Z} -metric graphs $\varphi : \Gamma' \rightarrow \Gamma$ taking e'_1 and e'_2 to e_1 , e'_3 to e_2 , and e'_4 to e_3 . The homomorphism φ_* is non-surjective since $|\text{Jac}_{\text{reg}}(\Gamma)| = 4$ and $|\text{Jac}_{\text{reg}}(\Gamma')| = 8$. The map φ can be enriched to a harmonic morphism $\tilde{\varphi}$ of metrized complexes of curves by attaching a \mathbf{P}^1 to each vertex and letting each morphism $\mathbf{P}^1 \rightarrow \mathbf{P}^1$ be $z \mapsto z^2$, with $0, \pm 1$, and ∞ being the marked points upstairs. By Remark 7.13, the morphism $\tilde{\varphi}$ lifts to a morphism $\psi : X' \rightarrow X$ of curves over K . Since all edges of G' have length 1, the minimal proper regular model of X' has a special fiber consisting of two \mathbf{P}^1 ’s crossing transversely. By Theorem 8.4, the induced covariant map on component groups of Néron models is not surjective.

REFERENCES

- [AB12] O. Amini and M. Baker, *Linear series on metrized complexes of algebraic curves*, Preprint available at <http://arxiv.org/abs/1204.3508>.
- [ABBR14] O. Amini, M. Baker, E. Brugallé, and J. Rabinoff, *Lifting harmonic morphisms II: tropical curves and metrized complexes*, <http://arxiv.org/abs/1404.3390> (2014).
- [ACP12] S. Abramovich, L. Caporaso, and S. Payne, *The tropicalization of the moduli space of curves*, Preprint available at <http://arxiv.org/abs/1212.0373>.
- [Ana00] C.K. Anand, *A discrete analogue of the harmonic morphism*, Harmonic morphisms, harmonic maps, and related topics (Brest, 1997), Chapman & Hall/CRC Res. Notes Math., vol. 413, Chapman & Hall/CRC, Boca Raton, FL, 2000, pp. 109–112.
- [Bak08] M. Baker, *Specialization of linear systems from curves to graphs*, Algebra Number Theory **2** (2008), no. 6, 613–653, With an appendix by Brian Conrad.
- [BBM11] B. Bertrand, E. Brugallé, and G. Mikhalkin, *Tropical open Hurwitz numbers*, Rend. Semin. Mat. Univ. Padova **125** (2011), 157–171.
- [BdlHN97] R. Bacher, P. de la Harpe, and T. Nagnibeda, *The lattice of integral flows and the lattice of integral cuts on a finite graph*, Bull. Soc. Math. France **125** (1997), no. 2, 167–198.
- [Ber90] V. G. Berkovich, *Spectral theory and analytic geometry over non-Archimedean fields*, Mathematical Surveys and Monographs, vol. 33, American Mathematical Society, Providence, RI, 1990.
- [Ber93] ———, *Étale cohomology for non-Archimedean analytic spaces*, Inst. Hautes Études Sci. Publ. Math. **78** (1993), 5–161.
- [Ber99] ———, *Smooth p -adic analytic spaces are locally contractible*, Invent. Math. **137** (1999), no. 1, 1–84.

- [BF11] M. Baker and X. Faber, *Metric properties of the tropical Abel-Jacobi map*, J. Algebraic Combin. **33** (2011), no. 3, 349–381.
- [BGR84] S. Bosch, U. Güntzer, and R. Remmert, *Non-Archimedean analysis*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 261, Springer-Verlag, Berlin, 1984.
- [BL85] S. Bosch and W. Lütkebohmert, *Stable reduction and uniformization of abelian varieties. I*, Math. Ann. **270** (1985), no. 3, 349–379.
- [BL93] ———, *Formal and rigid geometry. I. Rigid spaces*, Math. Ann. **295** (1993), no. 2, 291–317.
- [BN07] M. Baker and S. Norine, *Riemann-Roch and Abel-Jacobi theory on a finite graph*, Adv. Math. **215** (2007), no. 1, 766–788.
- [BN09] ———, *Harmonic morphisms and hyperelliptic graphs*, International Math. Research Notices **15** (2009), 2914–2955.
- [BPR11] M. Baker, S. Payne, and J. Rabinoff, *Non-Archimedean geometry, tropicalization, and metrics on curves*, 2011, Preprint available at <http://arxiv.org/abs/1104.0320>.
- [BR10] M. Baker and R. Rumely, *Potential theory and dynamics on the Berkovich projective line*, Mathematical Surveys and Monographs, vol. 159, American Mathematical Society, Providence, RI, 2010.
- [BR13] M. Baker and J. Rabinoff, *An analytic version of Raynaud’s description of component groups of Néron models*, in preparation (2013).
- [Cha12] M. Chan, *Tropical hyperelliptic curves*, J. Alg. Combin. **15** (2012), 2914–2955.
- [CKK] G. Cornelissen, F. Kato, and J. Kool, *A combinatorial Li-Yau inequality and rational points on curves*, To appear in Math. Ann., Preprint available at <http://arxiv.org/abs/1211.2681>.
- [Col03] R. F. Coleman, *Stable maps of curves*, Documenta Math. **Extra volume Kato** (2003), 217–225.
- [Con99] B. Conrad, *Irreducible components of rigid spaces*, Ann. Inst. Fourier (Grenoble) **49** (1999), no. 2, 473–541.
- [Duc08] A. Ducros, *Triangulations et cohomologie étale sur une courbe analytique p -adique*, J. Algebraic Geom. **17** (2008), no. 3, 503–575.
- [Fab13a] X. Faber, *Topology and geometry of the Berkovich ramification locus for rational functions*, Manuscripta Math. (2013), 439–474.
- [Fab13b] ———, *Topology and geometry of the Berkovich ramification locus for rational functions, II*, Math. Ann. **356** (2013), no. 3, 819–844.
- [HM82] J. Harris and D. Mumford, *On the Kodaira dimension of the moduli space of curves*, Invent. Math. **67** (1982), no. 1, 23–88, With an appendix by W. Fulton.
- [KS01] M. Kontsevich and Y. Soibelman, *Homological mirror symmetry and torus fibrations*, Symplectic geometry and mirror symmetry (Seoul, 2000), World Sci. Publishing, River Edge, NJ, 2001, pp. 203–263.
- [KS06] ———, *Affine structures and non-Archimedean analytic spaces*, The unity of mathematics, Progr. Math., vol. 244, Birkhäuser Boston, Boston, MA, 2006, pp. 321–385.
- [Liu03] Q. Liu, *Reduction and lifting of finite covers of curves*, Proceedings of the 2003 Workshop on Cryptography and Related Mathematics, Chuo University, 2003, pp. 161–180.
- [Liu06] ———, *Stable reduction of finite covers of curves*, Compositio Math. **142** (2006), no. 1, 101–118.
- [LL99] Q. Liu and D. Lorenzini, *Models of curves and finite covers*, Compositio Math. **118** (1999), no. 1, 61–102.
- [Mik] G. Mikhalkin, *Phase-tropical curves I. Realizability and enumeration*, In preparation.
- [Mik05] ———, *Enumerative tropical algebraic geometry in \mathbb{R}^2* , J. Amer. Math. Soc. **18** (2005), no. 2, 313–377.
- [Mik06] ———, *Tropical geometry and its applications*, International Congress of Mathematicians. Vol. II, Eur. Math. Soc., Zürich, 2006, pp. 827–852.
- [MZ08] G. Mikhalkin and I. Zharkov, *Tropical curves, their Jacobians and theta functions*, Curves and abelian varieties, Contemp. Math., vol. 465, Amer. Math. Soc., Providence, RI, 2008, pp. 203–230.
- [Ray70] M. Raynaud, *Spécialisation du foncteur de Picard*, Inst. Hautes Études Sci. Publ. Math. (1970), no. 38, 27–76.
- [Sai96] M. Saïdi, *Revêtements étales abéliens, courants sur les graphes et réductions semi-stable des courbes*, Manuscripta Mathematica **89** (1996), 245–265.
- [Sai97] ———, *Revêtements modérés et groupe fondamental de graphe de groupes*, Compos. Math. **107** (1997), 319–338.
- [Sai04] ———, *Wild ramification and a vanishing cycles formula*, J. Algebra **273** (2004), no. 1, 108–128.
- [SGA1] A. Grothendieck and M. Raynaud, *Revêtements étales et groupe fondamental (SGA 1)*, Séminaire de Géométrie Algébrique de l’Institut des Hautes Études Scientifiques (SGA 1), Société Mathématique de France, Paris, 2003.
- [Tem] M. Temkin, *Introduction to Berkovich analytic spaces*, Preprint available at <http://arxiv.org/abs/1010.2235v1>.
- [Thu05] A. Thuillier, *Théorie du potentiel sur les courbes en géométrie analytique non archimédienne. Applications à la théorie d’Arakelov*, Ph.D. thesis, University of Rennes, 2005, Preprint available at <http://tel.ccsd.cnrs.fr/documents/archives0/00/01/09/90/index.html>.
- [Ura00] H. Urakawa, *A discrete analogue of the harmonic morphism and green kernel comparison theorems*, Glasg. Math. J. **42** (2000), 319–334.
- [Wel13] J. Welliaveetil, *A Riemann-Hurwitz formula for skeleta in non-Archimedean geometry*, 2013, Preprint available at <http://arxiv.org/abs/1303.0164>.
- [Wew99] S. Wewers, *Deformation of tame admissible covers of curves*, Aspects of Galois theory (Gainesville, FL, 1996), London Math. Soc. Lecture Note Ser., vol. 256, Cambridge Univ. Press, Cambridge, 1999, pp. 239–282.

E-mail address: oamini@math.ens.fr

CNRS-DMA, ÉCOLE NORMALE SUPÉRIEURE, 45 RUE D'ULM, PARIS

E-mail address: mbaker@math.gatech.edu

SCHOOL OF MATHEMATICS, GEORGIA INSTITUTE OF TECHNOLOGY, ATLANTA GA 30332-0160, USA

E-mail address: brugalle@math.jussieu.fr

UNIVERSITÉ PIERRE ET MARIE CURIE, PARIS 6, 4 PLACE JUSSIEU, 75 005 PARIS, FRANCE

E-mail address: jrabinoff@math.gatech.edu

SCHOOL OF MATHEMATICS, GEORGIA INSTITUTE OF TECHNOLOGY, ATLANTA GA 30332-0160, USA