

ON GENERALIZED GAUSSIAN FREE FIELDS AND STOCHASTIC HOMOGENIZATION

YU GU, JEAN-CHRISTOPHE MOURRAT

ABSTRACT. We study a generalization of the notion of Gaussian free field (GFF). Although the extension seems minor, we first show that a generalized GFF does not satisfy the spatial Markov property, unless it is a classical GFF. In stochastic homogenization, the scaling limit of the corrector is a possibly generalized GFF described in terms of an “effective fluctuation tensor” that we denote by \mathbf{Q} . We prove an expansion of \mathbf{Q} in the regime of small ellipticity ratio. This expansion shows that the scaling limit of the corrector is not necessarily a classical GFF, and in particular does not necessarily satisfy the Markov property.

MSC 2010: 60G60, 35R60, 35B27.

KEYWORDS: Gaussian free field, Markov property, stochastic homogenization, corrector.

1. INTRODUCTION

This paper is motivated by recent results in the stochastic homogenization of operators of the form $-\nabla \cdot \mathbf{a} \nabla$, where $x \mapsto \mathbf{a}(x)$ is a random stationary field over \mathbb{R}^d with strong decorrelation properties, and valued in the set Sym_d^+ of d -by- d symmetric positive-definite matrices. In this context, we aim to obtain a precise description of the large-scale behavior of solutions of equations involving this operator, and in particular of the corrector.

The corrector in the direction $\xi \in \mathbb{R}^d$ can be defined, up to an additive constant, as the unique sub-linear function $\phi_\xi : \mathbb{R}^d \rightarrow \mathbb{R}$ such that $-\nabla \cdot \mathbf{a}(\xi + \nabla \phi_\xi) = 0$. The minimizer in $H_0^1(U)$ of the mapping $v \mapsto \int_U (\xi + \nabla v) \cdot \mathbf{a}(\xi + \nabla v)$, where U is a large domain, offers a finite-volume approximation of the corrector. This minimization of a random quadratic functional suggests a parallel with gradient Gibbs measures, where a deterministic functional of quadratic type is used to produce a probability measure via the Gibbs principle. Such gradient Gibbs measures are known to rescale to Gaussian free fields (GFFs), see [20, 6, 16]. It is therefore natural to conjecture the same scaling limit for the corrector [1].

Under assumptions described more precisely below, the scaling limit of the corrector was identified in [19, 18] as an element of a slightly larger class of random fields, which we will call *generalized* GFFs (see also [8]). Intuitively, we define the generalized GFF as the field Φ such that $\nabla \Phi$ is “as close as possible” to a white noise vector field. More precisely, a generalized GFF is a random field Φ solving

$$(1.1) \quad -\nabla \cdot \bar{\mathbf{a}} \nabla \Phi = \nabla \cdot W,$$

where $\bar{\mathbf{a}} \in \text{Sym}_d^+$ and W is a vector-valued white noise field with covariance matrix $\mathbf{Q} \in \text{Sym}_d^+$. For simplicity, we will only consider such equations in the full space \mathbb{R}^d with $d \geq 3$, but our arguments also cover domains with suitable boundary condition, or the addition of a massive term, with only minor modifications. We recall that a classical GFF is a Gaussian field whose covariance function is the Green function of $-\nabla \cdot \mathbf{b} \nabla$, for some $\mathbf{b} \in \text{Sym}_d^+$. When $\bar{\mathbf{a}}$ and \mathbf{Q} are proportional, the field Φ defined by (1.1) is a classical GFF, and thus satisfies the spatial Markov property [25].

The goal of this paper is twofold. In Section 2, we show that the Markov property does *not* extend to the wider class of generalized GFFs. In fact, no generalized GFF satisfies the Markov property, unless it is a classical GFF. Therefore, our apparently mild extension of the notion of GFF in fact wipes out one of its fundamental properties. In Section 3, we turn to the context of homogenization, and study the effective parameters describing the generalized GFF arising as the scaling limit of the corrector. We prove an asymptotic expansion for these coefficients, in the regime of small ellipticity contrast. This expansion is of independent interest, and shows that the scaling limit of the corrector is generically *not* a classical GFF, thereby justifying the relevance of the notion of generalized GFF.

We close this introduction with two remarks. First, in order to substantiate that the definition (1.1) is consistent with the intuition that $\nabla\Phi$ is chosen to be “as close as possible” to a white noise vector field, it is simpler to discuss the case where the underlying state space is the torus $\mathbb{T}^d := [0, 1]^d$. Endowing the space $L^2(\mathbb{T}^d, \mathbb{R}^d)$ with the scalar product

$$(F, G) \mapsto \int_{\mathbb{T}^d} F \cdot \bar{\mathbf{a}}^{-1} G,$$

we denote by $L^2_{\text{pot}}(\mathbb{T}^d)$ the closure in $L^2(\mathbb{T}^d, \mathbb{R}^d)$ of the set

$$\{\bar{\mathbf{a}}\nabla f : f \in C^\infty(\mathbb{T}^d)\}$$

and denote by $L^2_{\text{sol}}(\mathbb{T}^d)$ its orthogonal complement

$$L^2_{\text{sol}}(\mathbb{T}^d) := \{\mathbf{g} \in L^2(\mathbb{T}^d, \mathbb{R}^d) : \forall f \in C^\infty(\mathbb{T}^d), \int_{\mathbb{T}^d} \nabla f \cdot \mathbf{g} = 0\},$$

which is the set of solenoidal (i.e. divergence-free) vector fields. This provides us with the Helmholtz-Hodge decomposition of $L^2(\mathbb{T}^d, \mathbb{R}^d)$ into the orthogonal sum of $L^2_{\text{pot}}(\mathbb{T}^d)$ and $L^2_{\text{sol}}(\mathbb{T}^d)$. If W were smooth, this would allow us to interpret $\bar{\mathbf{a}}\nabla\Phi$ as the orthogonal projection of W onto $L^2_{\text{pot}}(\mathbb{T}^d)$. For non-smooth W , the interpretation remains valid by testing and duality.

Second, we point out that although the generalized GFF fails to satisfy the Markov property, it does satisfy a related domain decomposition property: there exists a family of random fields (Φ_A) on \mathbb{R}^d indexed by all borel subsets A of \mathbb{R}^d such that the following holds for every Borel set $A \subseteq \mathbb{R}^d$:

- we have $\Phi = \Phi_A + \Phi_{A^c}$, (A^c denotes the complement of A in \mathbb{R}^d);
- the random fields Φ_A and Φ_{A^c} are independent;
- the random field Φ_A is an $\bar{\mathbf{a}}$ -harmonic function in the interior of A^c .

Indeed, these properties are easy to verify from the definition of Φ_A as solving

$$-\nabla \cdot \bar{\mathbf{a}}\nabla\Phi_A = \nabla \cdot (W\mathbf{1}_A).$$

2. NON-MARKOV PROPERTY

In this section, we show that a generalized GFF satisfies the Markov property if and only if it is a classical GFF.

For k, l positive integers, we write $C_c^\infty(\mathbb{R}^k, \mathbb{R}^l)$ for the space of infinitely differentiable functions from \mathbb{R}^k to \mathbb{R}^l with compact support. For every $U \subseteq \mathbb{R}^k$, we let $C_c^\infty(U, \mathbb{R}^l) = \{f \in C_c^\infty(\mathbb{R}^k, \mathbb{R}^l) : \text{Supp } f \subseteq U\}$, where $\text{Supp } f$ denotes the support of f . We simply write $C_c^\infty(U) = C_c^\infty(U, \mathbb{R})$ and $C_c^\infty = C_c^\infty(\mathbb{R})$. We fix $d \geq 3$ and $\bar{\mathbf{a}}, \mathbf{Q} \in \text{Sym}_d^+$. We say that the random distribution $W = (W_1, \dots, W_d)$ is a white noise vector field with covariance matrix \mathbf{Q} if for every $\mathbf{f} = (\mathbf{f}_1, \dots, \mathbf{f}_d) \in C_c^\infty(\mathbb{R}^d, \mathbb{R}^d)$, the random variable

$$W(\mathbf{f}) := W_1(\mathbf{f}_1) + \dots + W_d(\mathbf{f}_d)$$

is a centered Gaussian with variance $\int_{\mathbb{R}^d} \mathbf{f} \cdot \mathbf{Q} \mathbf{f}$. We denote by $(\Omega, \mathbb{F}, \mathbb{P})$ the underlying probability space, and by \mathbb{E} the associated expectation. Informally,

$$\mathbb{E}[W_i(x) W_j(y)] = \mathbf{Q}_{ij} \delta(x - y),$$

where δ is the Dirac distribution. One can extend the set of admissible test functions for W to every element of $L^2(\mathbb{R}^d, \mathbb{R}^d)$ by density. We define the solution Φ to (1.1) to be the random distribution such that for every $f \in C_c^\infty(\mathbb{R}^d)$,

$$(2.1) \quad \Phi(f) = -W(\nabla(-\nabla \cdot \bar{\mathbf{a}} \nabla)^{-1} f),$$

where $(-\nabla \cdot \bar{\mathbf{a}} \nabla)^{-1} f$ is the unique function u tending to zero at infinity and such that $-\nabla \cdot \bar{\mathbf{a}} \nabla u = f$. Formally integrating by parts shows the consistency between (1.1) and (2.1). The latter makes sense since for $f \in C_c^\infty(\mathbb{R}^d)$, the function $\nabla(-\nabla \cdot \bar{\mathbf{a}} \nabla)^{-1} f$ is in $L^2(\mathbb{R}^d, \mathbb{R}^d)$, as can be checked for instance using the Green representation formula.

For every open $U \subseteq \mathbb{R}^d$, we define

$$\mathcal{F}(U) := \sigma\{\Phi(f), f \in C_c^\infty(U)\}.$$

We simply write $\mathcal{F} := \mathcal{F}(\mathbb{R}^d)$. For every closed $A \subseteq \mathbb{R}^d$, we define

$$\mathcal{F}(A) := \bigcap_{\text{open } U \supseteq A} \mathcal{F}(U).$$

We enlarge the σ -algebras defined above (without changing the notation) so that they contain all \mathbb{P} -negligible measurable sets. We say that the field Φ is *Markovian with respect to the open set U* if conditionally on $\mathcal{F}(\partial U)$, the σ -algebras $\mathcal{F}(\bar{U})$ and $\mathcal{F}(U^c)$ are independent (where \bar{U} denotes the closure of U , U^c the complement of U , and $\partial U = \bar{U} \cap U^c$).

Theorem 2.1. *Let U be an open subset of \mathbb{R}^d , $U \notin \{\emptyset, \mathbb{R}^d\}$. The field Φ is Markovian with respect to U if and only if Φ is a classical GFF.*

Let $\mathcal{L} := (-\nabla \cdot \bar{\mathbf{a}} \nabla)(-\nabla \cdot \mathbf{Q} \nabla)^{-1}(-\nabla \cdot \bar{\mathbf{a}} \nabla)$. It follows from (2.1) that for every $f, g \in C_c^\infty$,

$$(2.2) \quad \mathbb{E}[\Phi(f) \Phi(g)] = \int f \mathcal{L}^{-1} g.$$

In particular, if $\bar{\mathbf{a}}$ and \mathbf{Q} are proportional, then the field Φ is a classical GFF, and we recall that in this case the Markov property is well-known. In order to prove the theorem, it thus suffices to show that if Φ is Markovian with respect to U , then $\bar{\mathbf{a}}$ and \mathbf{Q} are proportional.

To our knowledge, the rigorous study of the Markov property of random fields was initiated with [15], where Lévy's "Brownian motion" indexed by a multidimensional parameter [14] is shown to be Markovian if and only if the space dimension is odd.

For random fields on discrete graphs, the Markov property is equivalent to the locality of the "energy function" (in the Gaussian case, this is the Dirichlet form, and more generally, we mean the logarithm of the probability density, up to a constant). This equivalence can be checked by a direct computation for Gaussian fields, and we refer to [10, Theorems 4.1 and 4.2] for a more general statement. It is natural to expect a similar phenomenon in the continuum. However, a counter-example to this conjecture was given in [23]. We will recall this counter-example in subsection 2.2. In spite of this, relying on the fact that the field Φ is "at least as regular as white noise", we will be able to justify this conjecture in our context.

Prior to the counter-example of [23], incorrect proofs of the general conjecture were published. In [22, 13], the arguments are laid down pretending that the field is

defined pointwise, and therefore the difficulty caused by the possibly low regularity of the field is missed. The paper [9] proceeds more carefully, but is also flawed¹.

The equivalence between locality of \mathcal{L} and the Markov property was also investigated in the framework of Dirichlet form theory [4, 24, 11]. The arguments given there rely on potential theory, and only apply to operators satisfying the maximum principle. This is unfortunately not the case of the operator \mathcal{L} we consider here. In order to see this, we can use the general result of [3], which identifies the class of integro-differential operators satisfying the maximum principle. In our case, the operator is moreover invariant under translation. In this setting, the results of [3] can be understood as follows: if the operator satisfies the maximum principle, then it is the generator of a Lévy process. Moreover, this Lévy process must have the same scale invariance as Brownian motion, in view of the scaling properties of \mathcal{L} . By the Lévy-Khintchine formula, it follows that such a Lévy process must be a multiple of Brownian motion. The operator \mathcal{L} must therefore be of the form $-\nabla \cdot \mathbf{b} \nabla$ for some matrix \mathbf{b} , but we show below that this can only happen when $\bar{\mathbf{a}}$ and \mathbf{Q} are proportional.

2.1. Proof of Theorem 2.1. To sum up: using some mild regularity property of the field Φ defined by (2.1), we show that the Markov property implies that \mathcal{L} is a local operator. We then observe that \mathcal{L} is not a local operator, unless $\bar{\mathbf{a}}$ and \mathbf{Q} are proportional. As announced in the theorem, we will be sufficiently careful to actually only require the Markov property with respect to one non-trivial open set.

We denote by \mathcal{H} the Hilbert space obtained as the closure of $\{\Phi(f), f \in C_c^\infty\}$ in $L^2(\Omega, \mathcal{F}, \mathbb{P})$. We denote by \mathcal{K} the Hilbert space obtained by completing $\{\mathcal{L}^{-1}f, f \in C_c^\infty\}$ with respect to the scalar product

$$\langle f, g \rangle_{\mathcal{L}} := \int f \mathcal{L}g.$$

The space \mathcal{K} is usually called the reproducing kernel Hilbert space. Note that $f \mapsto \langle f, f \rangle_{\mathcal{L}}^{1/2}$ is within multiplicative constants of the homogeneous H^1 norm. By (2.2), the spaces \mathcal{H} and \mathcal{K} are isometric:

Lemma 2.2. *The mapping*

$$\begin{cases} \{\Phi(f), f \in C_c^\infty\} & \rightarrow \mathcal{K} \\ \Phi(f) & \mapsto \mathcal{L}^{-1}f \end{cases}$$

is well-defined and extends to an isometry from \mathcal{H} onto \mathcal{K} .

For every open $U \subseteq \mathbb{R}^d$, we let $\mathcal{H}(U)$ be the closure of $\{\Phi(f), f \in C_c^\infty(U)\}$ in \mathcal{H} , and $\mathcal{K}(U)$ be the closure of $\{\mathcal{L}^{-1}f, f \in C_c^\infty(U)\}$ in \mathcal{K} . The isometry of Lemma 2.2 induces an isometry between $\mathcal{H}(U)$ and $\mathcal{K}(U)$. If $A \subseteq \mathbb{R}^d$ is a closed set, we define

$$\mathcal{H}(A) := \bigcap_{\text{open } U \supseteq A} \mathcal{H}(U), \quad \mathcal{K}(A) := \bigcap_{\text{open } U \supseteq A} \mathcal{K}(U).$$

The following lemma is classical.

Lemma 2.3. *Let U be an open set.*

- (1) *The set $\mathcal{H}(U)$ is the set of $\mathcal{F}(U)$ -measurable elements of \mathcal{H} .*
- (2) *The map*

$$\begin{cases} \mathcal{H} & \rightarrow \mathcal{H} \\ X & \mapsto \mathbb{E}[X \mid \mathcal{F}(U)] \end{cases}$$

is the orthogonal projection onto $\mathcal{H}(U)$.

¹there, the set $\mathfrak{M}(D_-)$ should be closed in order for [9, Lemma 1] to hold; but then the property $F \in \mathfrak{M}(D_-) \implies \text{Supp } F \subseteq D_-$ used in the proof of [9, Lemma 2] is false in general.

Proof. Every element of $\mathcal{H}(U)$ is $\mathcal{F}(U)$ -measurable. Conversely, if $X \in \mathcal{H}(U)^\perp$, then for every $f_1, \dots, f_k \in C_c^\infty(U)$, we have $\mathbb{E}[X \Phi(f_i)] = 0$. Since \mathcal{H} is a Gaussian space, this implies that X is independent of $(\Phi(f_1), \dots, \Phi(f_k))$, and thus X is independent of $\mathcal{F}(U)$. We have shown

$$\mathcal{H}(U)^\perp \subseteq \{Y \in \mathcal{H} : Y \text{ is } \mathcal{F}(U)\text{-measurable}\}^\perp.$$

Since both spaces are closed, this completes the proof of (1). Part (2) follows at once, since for every $X \in \mathcal{H}$, we have $(X - \mathbb{E}[X | \mathcal{F}(U)]) \in \mathcal{H}(U)^\perp$. \square

Lemma 2.4. *Let $\mathcal{H}_0(\bar{U})$ denote the orthogonal complement of $\mathcal{H}(\partial U)$ in $\mathcal{H}(\bar{U})$. If Φ is Markovian with respect to the open set U , then*

$$\mathcal{H} = \mathcal{H}_0(\bar{U}) \overset{\perp}{\oplus} \mathcal{H}(\partial U) \overset{\perp}{\oplus} \mathcal{H}_0(U^c).$$

Proof. We decompose the proof into three steps.

Step 1. We first show that

$$(2.3) \quad \text{the orthogonal projection of } \mathcal{H}(U^c) \text{ onto } \mathcal{H}(\bar{U}) \text{ is } \mathcal{H}(\partial U).$$

By Lemma 2.3, it suffices to show that for every $\mathcal{F}(U^c)$ -measurable $X \in \mathcal{H}$,

$$\mathbb{E}[X | \mathcal{F}(\partial U)] = \mathbb{E}[X | \mathcal{F}(\bar{U})].$$

We show that

$$(2.4) \quad \mathbb{E} \left[\left(\mathbb{E}[X | \mathcal{F}(\partial U)] - \mathbb{E}[X | \mathcal{F}(\bar{U})] \right)^2 \right] = 0.$$

By the Markov property and the inclusion $\mathcal{F}(\partial U) \subseteq \mathcal{F}(\bar{U})$,

$$(2.5) \quad \mathbb{E} \left[X \mathbb{E}[X | \mathcal{F}(\bar{U})] | \mathcal{F}(\partial U) \right] = \left(\mathbb{E}[X | \mathcal{F}(\partial U)] \right)^2.$$

In particular,

$$\begin{aligned} \mathbb{E} \left[\left(\mathbb{E}[X | \mathcal{F}(\bar{U})] \right)^2 \right] &= \mathbb{E} \left[X \mathbb{E}[X | \mathcal{F}(\bar{U})] \right] \\ &\stackrel{(2.5)}{=} \mathbb{E} \left[\left(\mathbb{E}[X | \mathcal{F}(\partial U)] \right)^2 \right], \end{aligned}$$

and this proves (2.4).

Step 2. We show that

$$\mathcal{H}_0(\bar{U}) \overset{\perp}{\oplus} \mathcal{H}(\partial U) \overset{\perp}{\oplus} \mathcal{H}_0(U^c) = \mathcal{H}(\bar{U}) + \mathcal{H}(U^c).$$

The equality

$$\mathcal{H}_0(\bar{U}) + \mathcal{H}(\partial U) + \mathcal{H}_0(U^c) = \mathcal{H}(\bar{U}) + \mathcal{H}(U^c)$$

is clear. The orthogonality of the sets on the left-hand side follows from the definition of \mathcal{H}_0 and the previous step. In particular, the set $\mathcal{H}(\bar{U}) + \mathcal{H}(U^c)$ is closed.

Step 3. Let $f \in C_c^\infty$. In order to complete the proof, it suffices to see that

$$(2.6) \quad \Phi(f) \in \mathcal{H}(\bar{U}) + \mathcal{H}(U^c).$$

This is the step of the proof where we need to use the fact that Φ is “at least as regular as white noise”. In more precise words, we show that we can control the quadratic form $g \mapsto \int g \mathcal{L}^{-1} g$ over local functions g by the L^2 norm of g squared, and that this is sufficient to conclude. Without loss of generality, we assume that the support of f is contained in the unit ball $B(0, 1)$. Let $\chi : \mathbb{R} \rightarrow [0, 1]$ be a C^∞ function such that $\chi = 1$ on $(-\infty, 0]$ and $\chi = 0$ on $[1, +\infty)$, and let

$$\chi_n := \begin{cases} \mathbb{R}^d & \rightarrow [0, 1] \\ x & \mapsto \chi(n \operatorname{dist}(x, U)), \end{cases}$$

where $\text{dist}(x, U)$ denotes the distance between x and the open set U . Note that

$$\Phi(f) = \Phi(f\chi_n) + \Phi(f[1 - \chi_n]),$$

and that $\Phi(f[1 - \chi_n]) \in \mathcal{H}(U^c)$. We now argue that $\Phi(f\chi_n)$ converges to an element of $\mathcal{H}(\bar{U})$ as n tends to infinity. This would imply that $\Phi(f[1 - \chi_n])$ converges to an element of $\mathcal{H}(U^c)$ and therefore complete the proof of (2.6). If the sequence $(\Phi(f\chi_n))$ converges in \mathcal{H} , then the limit is necessarily in $\mathcal{H}(\bar{U})$; so what needs to be argued is simply the convergence of $\Phi(f[1 - \chi_n])$ in \mathcal{H} . By (2.2), for every $g \in C_c^\infty$,

$$\mathbb{E}[\Phi(g)^2] = \int g\mathcal{L}^{-1}g,$$

so we need to check that $f\chi_n$ is a Cauchy sequence with respect to the seminorm

$$g \mapsto \left(\int g\mathcal{L}^{-1}g \right)^{\frac{1}{2}}.$$

Letting Λ be the largest eigenvalue of \mathbf{Q} and $h := (-\nabla \cdot \mathbf{a}\nabla)^{-1}g$, we have

$$\int g\mathcal{L}^{-1}g \leq \Lambda \int |\nabla h|^2.$$

Since $d \geq 3$, using the Green function representation, we obtain that there exists a constant $C < \infty$ (depending only on \mathbf{a}) such that

$$|\nabla h|(x) \leq C \int_{\mathbb{R}^d} \frac{1}{|x-y|^{d-1}} |g|(y) dy.$$

By Young's convolution inequality, the L^2 norm of the function

$$x \mapsto \int_{\mathbb{R}^d} \frac{\mathbf{1}_{|x-y| \leq 1}}{|x-y|^{d-1}} |g|(y) dy$$

is bounded by a constant times $\|g\|_{L^2}$. For functions g with support in $B(0, 1)$, we also have

$$\int_{\mathbb{R}^d} \frac{\mathbf{1}_{|x-y| > 1}}{|x-y|^{d-1}} |g|(y) dy \leq C \frac{\|g\|_{L^1}}{1+|x|^{d-1}} \leq C \frac{\|g\|_{L^2}}{1+|x|^{d-1}}.$$

To sum up, we have shown that there exists a constant $C < \infty$ such that for every $g \in C_c^\infty$ with support in $B(0, 1)$,

$$\left(\int g\mathcal{L}^{-1}g \right)^{\frac{1}{2}} \leq C\|g\|_{L^2}.$$

Since $f\chi_n \rightarrow f$ in L^2 , this completes the proof. \square

Lemma 2.5. *If Φ is Markovian with respect to the open set U , and if $f_1 \in C_c^\infty(U)$ and $f_2 \in C_c^\infty(\bar{U}^c)$, then $\langle f_1, f_2 \rangle_{\mathcal{L}} = 0$.*

Proof. The isometry of Lemma 2.2 transports the decomposition of \mathcal{H} in Lemma 2.4 into a decomposition of \mathcal{K} . In particular,

$$\mathcal{K}(U^c)^\perp \subseteq \mathcal{K}(\bar{U}).$$

Let $f_1 \in C_c^\infty(U)$, and let $S := \text{Supp } f_1$. Note that

$$\bigcap_{\substack{\text{open } O \supseteq U^c \\ O \cap S = \emptyset}} \mathcal{K}(O) = \mathcal{K}(U^c).$$

Let $g \in C_c^\infty$ with support disjoint from S . We have

$$0 = \int f_1 g = \langle f_1, \mathcal{L}^{-1}g \rangle_{\mathcal{L}}.$$

By density, we deduce that $f_1 \in \mathcal{K}(U^c)^\perp \subseteq \mathcal{K}(\bar{U})$. By symmetry, we also have $f_2 \in \mathcal{K}(\bar{U})^\perp$, which completes the proof. \square

Lemma 2.6. *Let U be an open subset of \mathbb{R}^d , $U \notin \{\emptyset, \mathbb{R}^d\}$. If $\bar{\mathbf{a}}$ and \mathbf{Q} are not proportional, then there exist $f_1 \in C_c^\infty(U)$, $f_2 \in C_c^\infty(\bar{U}^c)$ such that $\langle f_1, f_2 \rangle_{\mathcal{L}} \neq 0$.*

Proof. In the Fourier domain, $\langle f_1, f_2 \rangle_{\mathcal{L}}$ can be written explicitly as

$$\begin{aligned} \langle f_1, f_2 \rangle_{\mathcal{L}} &= \int_{\mathbb{R}^d} f_1(x) (-\nabla \cdot \bar{\mathbf{a}} \nabla) (-\nabla \cdot \mathbf{Q} \nabla)^{-1} (-\nabla \cdot \bar{\mathbf{a}} \nabla) f_2(x) dx \\ &= \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} \widehat{f_1}(\xi) \overline{\widehat{f_2}(\xi)} \frac{(\xi \cdot \bar{\mathbf{a}} \xi)^2}{\xi \cdot \mathbf{Q} \xi} d\xi. \end{aligned}$$

We change ξ to $\mathbf{Q}^{-\frac{1}{2}} \xi$, and then go back to physical domain, to obtain that

$$\begin{aligned} &\langle f_1, f_2 \rangle_{\mathcal{L}} \\ &= |\mathbf{Q}|^{\frac{1}{2}} \int_{\mathbb{R}^d} (-\nabla \cdot \mathbf{Q}^{-\frac{1}{2}} \bar{\mathbf{a}} \mathbf{Q}^{-\frac{1}{2}} \nabla) f_1(\mathbf{Q}^{\frac{1}{2}} x) (-\Delta)^{-1} (-\nabla \cdot \mathbf{Q}^{-\frac{1}{2}} \bar{\mathbf{a}} \mathbf{Q}^{-\frac{1}{2}} \nabla) f_2(\mathbf{Q}^{\frac{1}{2}} x) dx. \end{aligned}$$

Since $\bar{\mathbf{a}}$ and \mathbf{Q} are not proportional, $\mathbf{Q}^{-\frac{1}{2}} \bar{\mathbf{a}} \mathbf{Q}^{-\frac{1}{2}}$ is not a multiple of identity. By denoting $\mathbf{A} = \mathbf{Q}^{-\frac{1}{2}} \bar{\mathbf{a}} \mathbf{Q}^{-\frac{1}{2}}$, we only need to show that there exists g_1, g_2 whose supports are in $\mathbf{Q}^{\frac{1}{2}} U$ and $\mathbf{Q}^{\frac{1}{2}} \bar{U}^c$ respectively and such that

$$\int_{\mathbb{R}^d} (-\nabla \cdot \mathbf{A} \nabla) g_1(x) (-\Delta)^{-1} (-\nabla \cdot \mathbf{A} \nabla) g_2(x) dx \neq 0.$$

For x away from the support of g_2 , we can write

$$\begin{aligned} (-\Delta)^{-1} (-\nabla \cdot \mathbf{A} \nabla) g_2(x) &= \int_{\mathbb{R}^d} \mathcal{G}(x-y) (-\nabla \cdot \mathbf{A} \nabla) g_2(y) dy \\ &= \int_{\mathbb{R}^d} (-\nabla \cdot \mathbf{A} \nabla \mathcal{G})(x-y) g_2(y) dy \end{aligned}$$

with \mathcal{G} the Green function of $-\Delta$. Since g_1, g_2 have disjoint supports, we can further integrate by parts to obtain

$$\begin{aligned} &\int_{\mathbb{R}^d} (-\nabla \cdot \mathbf{A} \nabla) g_1(x) (-\Delta)^{-1} (-\nabla \cdot \mathbf{A} \nabla) g_2(x) dx \\ &= \int_{\mathbb{R}^{2d}} g_1(x) K(x-y) g_2(y) dy dx \end{aligned}$$

with $K = \nabla \cdot \mathbf{A} \nabla (\nabla \cdot \mathbf{A} \nabla \mathcal{G})$. By an explicit calculation, we have, for $x \neq 0$,

$$\begin{aligned} (2.7) \quad K(x)/c &= \frac{1}{|x|^{d+2}} [-d \operatorname{Tr}(\mathbf{A})^2 - 2d \operatorname{Tr}(\mathbf{A}^2)] + \frac{x^t \mathbf{A} x \operatorname{Tr}(\mathbf{A})}{|x|^{d+4}} 2d(d+2) \\ &\quad + \frac{x^t \mathbf{A}^2 x}{|x|^{d+4}} 4d(d+2) - \frac{(x^t \mathbf{A} x)^2}{|x|^{d+6}} d(d+2)(d+4) \end{aligned}$$

for some constant $c \neq 0$. We show in Appendix A that if \mathbf{A} is not a multiple of the identity, then $(x^t \mathbf{A} x)^2 / |x|^2$ cannot be a quadratic form. As a consequence, the function $x \mapsto |x|^{d+6} K(x)$ is a non-zero polynomial over $\mathbb{R}^d \setminus \{0\}$. This implies that we can find g_1, g_2 whose supports are in $\mathbf{Q}^{\frac{1}{2}} U$ and $\mathbf{Q}^{\frac{1}{2}} \bar{U}^c$ respectively and satisfy

$$\int_{\mathbb{R}^{2d}} g_1(x) K(x-y) g_2(y) dy dx \neq 0.$$

The proof is complete. \square

Proof of Theorem 2.1. The result follows from Lemmas 2.5 and 2.6. \square

2.2. A counter-example with less regularity. For the reader's convenience, we briefly recall the counter-example given in [23]. Let X be the random distribution over \mathbb{R} such that for every $f \in C_c^\infty$, $X(f)$ is a centered Gaussian with variance $\int_{\mathbb{R}} [f^2 + (f')^2]$, where f' is the derivative of f . We let $\mathcal{L}^{-1}f := f - f''$, so that \mathcal{L} is a convolution operator with kernel $\frac{1}{2} \exp(-|x - y|)$, and hence is not local.

The set of admissible test functions for X can be extended to every element of the Sobolev space $W^{1,2}$ of functions in L^2 with weak derivative in L^2 . We recall that by Morrey's inequality, elements of $W^{1,2}$ are continuous functions. For every open or closed $A \subseteq \mathbb{R}$, we write

$$W^{1,2}(A) := \{f \in W^{1,2} : f = 0 \text{ on } A^c\}.$$

Note that if A is closed, then

$$W^{1,2}(A) = \bigcap_{\text{open } U \supseteq A} W^{1,2}(U).$$

The space $W^{1,2}$ has a Hilbert space structure. We write $W_{\text{perp}}^{1,2}(A)$ for the orthogonal complement of $W^{1,2}(A)$ in $W^{1,2}$. For every $V \subseteq W^{1,2}$, we write

$$X(V) := \{X(f) : f \in V\},$$

and check that for every open or closed $A \subseteq \mathbb{R}^d$,

$$\mathcal{H}(A) = X(W^{1,2}(A)).$$

For A open or closed, denote by $\mathcal{F}_H(A)$ the σ -algebra generated by the random variables in $\mathcal{H}(A)$. We claim that for every such A , we have $\mathcal{F}(A) = \mathcal{F}_H(A)$. The identity is clear if A is open. For A closed, we clearly have $\mathcal{F}_H(A) \subseteq \mathcal{F}(A)$. Conversely, observe that

$$W^{1,2}(A) + \bigcup_{\text{open } U \supseteq A} W_{\text{perp}}^{1,2}(U)$$

is dense in $W^{1,2}$ (the closure of the union above is $W_{\text{perp}}^{1,2}(A)$). Hence, by the martingale convergence theorem, if Z is a bounded \mathcal{F} -measurable random variable, then

$$(2.8) \quad Z = \lim_{\text{open } U \downarrow A} \mathbb{E}[Z | X(W^{1,2}(A) + W_{\text{perp}}^{1,2}(U))].$$

If Z is $\mathcal{F}(A)$ -measurable, then it is independent of $X(W_{\text{perp}}^{1,2}(U))$. By (2.8), it is therefore $\mathcal{F}_H(A)$ -measurable.

Let $f \in W^{1,2}((-\infty, 0])$ and $g \in W^{1,2}([0, +\infty))$. We have

$$\mathbb{E}[X(f)X(g)] = \int_{\mathbb{R}} [fg + f'g'],$$

where $f', g' \in L^2(\mathbb{R})$ are the weak derivatives of $f, g \in L^2(\mathbb{R})$. The functions f and f' are supported in $(-\infty, 0]$, while g and g' are supported in $[0, +\infty)$, so $\mathbb{E}[X(f)X(g)] = 0$. Since the spaces $\mathcal{H}((-\infty, 0])$ and $\mathcal{H}([0, +\infty))$ are Gaussian, we infer that $\mathcal{F}((-\infty, 0])$ and $\mathcal{F}([0, +\infty))$ are independent. Moreover, $\mathcal{F}(\{0\})$ is trivial, and therefore the field X is Markovian with respect to $(-\infty, 0)$.

Note that the field X has the regularity of the derivative of white noise. The proof of Step 3 of Lemma 2.4 breaks down, and

$$\mathcal{H}((-\infty, 0]) + \mathcal{H}([0, +\infty)) \neq \mathcal{H},$$

since $W^{1,2}((-\infty, 0]) + W^{1,2}([0, +\infty))$ only contains functions that vanish at the origin, and therefore

$$W^{1,2}((-\infty, 0]) + W^{1,2}([0, +\infty)) \neq W^{1,2}.$$

3. HOMOGENIZATION AND EXPANSION OF EFFECTIVE FLUCTUATION TENSOR

We now turn to stochastic homogenization. We focus on discrete elliptic equations in divergence form:

$$\nabla^* \mathbf{a}(x) \nabla u(x) = f(x), \quad x \in \mathbb{Z}^d,$$

with $d \geq 3$, where $\mathbf{a} : \mathbb{Z}^d \rightarrow \text{Sym}_d^+$ is a field of diagonal matrices. The entries $\mathbf{a}_{ii}(x)$, $x \in \mathbb{Z}^d$, $i = 1, \dots, d$ are i.i.d. random variables defined on the probability space $(\Omega, \mathbf{F}, \mathbb{P})$. We assume that $\mathbf{a}_{ii}(x) \in [1 - \tau_0, 1 + \tau_0]$ for some fixed $\tau_0 \in (0, 1)$. Let \mathbb{B} be the set of nearest neighbor edges in \mathbb{Z}^d , $\{e_i, i = 1, \dots, d\}$ be the canonical basis, and for $e = (x, x + e_i) \in \mathbb{B}$, we view

$$\mathbf{a}(e) := \mathbf{a}_{ii}(x)$$

as the random conductance on the edge linking x to $x + e_i$. The discrete gradient and divergence for $f : \mathbb{Z}^d \rightarrow \mathbb{R}$ and $F : \mathbb{Z}^d \rightarrow \mathbb{R}^d$ are defined respectively by

$$\nabla f = (\nabla_1 f, \dots, \nabla_d f), \quad \nabla^* F = \sum_{i=1}^d \nabla_i^* F_i,$$

with

$$\nabla_i f(x) = f(x + e_i) - f(x), \quad \nabla_i^* F_i = F_i(x - e_i) - F_i(x).$$

For any $e = (\underline{e}, \bar{e}) \in \mathbb{B}$ and $f : \mathbb{Z}^d \rightarrow \mathbb{R}$, we also write $\nabla f(e) = f(\bar{e}) - f(\underline{e})$. For any $\xi \in \mathbb{R}^d$ and $i = 1, \dots, d$, we define $\xi(e_i) = \xi_i$.

There exists a deterministic matrix $\bar{\mathbf{a}} \in \text{Sym}_d^+$ such that if $\alpha > 0$ and $u^{(\varepsilon)}$ solves

$$(\varepsilon^2 \alpha + \nabla^* \mathbf{a}(x) \nabla) u^{(\varepsilon)}(x) = \varepsilon^2 f(\varepsilon x), \quad x \in \mathbb{Z}^d,$$

then it was shown in [12, Theorem 4] that $u_\varepsilon(x) := u^{(\varepsilon)}(x/\varepsilon)$ ($x \in \varepsilon \mathbb{Z}^d$) converges in $L^2(\mathbb{R}^d \times \Omega)$ to the deterministic solution of

$$(\alpha - \nabla \cdot \bar{\mathbf{a}} \nabla) u_h(x) = f(x), \quad x \in \mathbb{R}^d,$$

where u_ε is extended to \mathbb{R}^d as a piecewise constant function. In other words, on large scales, the random coefficients $\mathbf{a}(x)$ behave like the homogeneous, deterministic coefficients $\bar{\mathbf{a}}$, and the discrete heterogeneous operator $\nabla^* \mathbf{a}(x) \nabla$ is averaged as the continuous homogeneous operator $-\nabla \cdot \bar{\mathbf{a}} \nabla$.

As was pointed out in the introduction, the corrector plays an important role in proving the convergence of $u_\varepsilon \rightarrow u_h$ and calculating the homogenized matrix $\bar{\mathbf{a}}$. The equation of the corrector in the direction of $\xi \in \mathbb{R}^d$ says

$$(3.1) \quad \nabla^* \mathbf{a}(x) (\nabla \phi_\xi(x) + \xi) = 0, \quad x \in \mathbb{Z}^d.$$

Since we assume $d \geq 3$, there exists a stationary zero-mean random field solving (3.1), as was shown in [7].

From now on, we assume furthermore that

$$\mathbf{a}(e) = a(\zeta_e),$$

where $a : \mathbb{R} \rightarrow \mathbb{R}$ is a fixed twice differentiable function with bounded first and second derivatives (and taking values in $[1 - \tau_0, 1 + \tau_0]$), and $\{\zeta_e : e \in \mathbb{B}\}$ is a sequence of i.i.d. standard Gaussian random variables. Under this technically convenient condition, the large-scale behavior of the corrector is approximately a generalized GFF. More precisely, it was shown in [19, 18] that

$$(3.2) \quad \varepsilon^{-(\frac{d}{2}-1)} \phi_\xi(\cdot/\varepsilon) \xrightarrow[\varepsilon \rightarrow 0]{(\text{law})} \Phi_\xi,$$

where Φ_ξ solves

$$(3.3) \quad -\nabla \cdot \bar{\mathbf{a}} \nabla \Phi_\xi = \nabla \cdot W_\xi$$

for a certain white noise vector field W_ξ . A heuristic derivation of (3.3) can be found in [8, Section 1].

The Gaussian white noise vector field W_ξ appearing in (3.3) has a covariance matrix Q_ξ given explicitly by

$$(3.4) \quad [Q_\xi]_{ij} = \sum_{k=1}^d \langle (e_i + \nabla\phi_i)(e_k)(\xi + \nabla\phi_\xi)(e_k)\mathbf{a}'(e_k) \\ (1 + \mathcal{L})^{-1}\mathbf{a}'(e_k)(e_j + \nabla\phi_j)(e_k)(\xi + \nabla\phi_\xi)(e_k) \rangle.$$

Here $\langle \cdot \rangle$ denotes the expectation in Ω and $\{\phi_i = \phi_{e_i}, i = 1, \dots, d\}$ are the correctors in the canonical directions. For sufficiently smooth functions on Ω , the weak derivative with respect to ζ_e is denoted by ∂_e , and its adjoint is $\partial_e^* = \zeta_e - \partial_e$. In (3.4), we have set

$$\mathcal{L} := \sum_{e \in \mathbb{B}} \partial_e^* \partial_e,$$

and

$$\mathbf{a}'(e) := \partial_e \mathbf{a}(e) = a'(\zeta_e).$$

Note that the map $\xi \mapsto Q_\xi$ is quadratic. This object should thus be viewed as a four-fold tensor, which we propose to call the *effective fluctuation tensor*.

The convergence of the random distribution in (3.1) to Φ_ξ motivates our study of generalized GFFs. Since we assume the coefficients to be i.i.d., the homogenized matrix inherits the symmetries of the lattice, and $\bar{\mathbf{a}}$ is thus a multiple of the identity.

The goal of this section is to obtain an expansion of Q_ξ in the regime of small ellipticity contrast. This is interesting per se, e.g. as a means to compute the effective fluctuation tensor when we expect only a small amount of random fluctuation in the underlying medium, in which case Q_ξ may be replaced by the series expansion up to certain order depending on the desired accuracy. It also allows us to give examples of environments such that Q_ξ is not a multiple of the identity, which implies, in view of Theorem 2.1, that the limiting field Φ_ξ does *not* satisfy the Markov property.

Without loss of generality, we may assume that \mathbf{a} takes the form

$$(3.5) \quad \mathbf{a}(x) = \mathbf{I}_d + \tau \mathbf{b}(x)$$

where \mathbf{b} is a field of diagonal matrices with $\langle \mathbf{b} \rangle = 0$ and $\|\mathbf{b}\|_{L^\infty} \leq 1$, and $\tau \in (0, 1)$ is a free parameter that we will take sufficiently small. We let δ_{ij} be the Kronecker symbol, that is, $\delta_{ij} = 1$ if $i = j$, and $\delta_{ij} = 0$ otherwise.

Theorem 3.1. *There exists $\tau_0 > 0$ such that for every $\tau \in [0, \tau_0)$, $\xi \in \mathbb{R}^d$ and $i, j = 1, \dots, d$, we have the convergent expansion*

$$[Q_\xi]_{ij} = \tau^2 \sum_{l=0}^{\infty} c_{\xi, l, i, j} \tau^l,$$

where $c_{\xi, l, i, j}$ can be computed explicitly as explained below, and

$$(3.6) \quad c_{\xi, 0, i, j} = \delta_{ij} \xi_i^2 \langle \mathbf{b}(e)^2 \rangle,$$

$$(3.7) \quad i \neq j \quad \implies \quad c_{\xi, 1, i, j} = 0,$$

$$(3.8) \quad i \neq j \quad \text{and} \quad \xi_i \xi_j \neq 0 \quad \implies \quad c_{\xi, 2, i, j} \neq 0.$$

In particular, if $\tau > 0$ is sufficiently small and $\xi \neq 0$, then Q_ξ is not a multiple of identity.

We first perform a formal expansion of Q_ξ and observe that (3.6)–(3.8) should hold. We then justify the full expansion rigorously.

3.1. A formal expansion. In view of (3.5), the corrector equation (3.1) in the direction $\eta \in \mathbb{R}^d$ can be rewritten as

$$-\Delta\phi_\eta = -\tau\nabla^*\mathbf{b}(\nabla\phi_\eta + \eta),$$

where $\Delta := -\nabla^*\nabla$ is the discrete Laplacian. Formally, we can write

$$\nabla\phi_\eta = -\tau\nabla(-\Delta)^{-1}\nabla^*\mathbf{b}(\nabla\phi_\eta + \eta).$$

If the operator $-\tau\nabla(-\Delta)^{-1}\nabla^*\mathbf{b}$ is a contraction (note that $\tau \ll 1$), then

$$(3.9) \quad \eta + \nabla\phi_\eta = \sum_{k=0}^{\infty} X_{k,\eta},$$

with

$$(3.10) \quad X_{k,\eta} = [-\tau\nabla(-\Delta)^{-1}\nabla^*\mathbf{b}]^k\eta.$$

There are four factors of $\eta + \nabla\phi_\eta$ appearing in (3.4) with $\eta = e_i, e_j, \xi$, which we will replace by the expansion in (3.9). Since $\mathbf{a}'(e) = \tau\mathbf{b}'(e)$ for any $e \in \mathbb{B}$, we obtain an expansion of $[\mathbf{Q}_\xi]_{ij}$ in terms of τ written as:

$$(3.11) \quad [\mathbf{Q}_\xi]_{ij} = \tau^2 \sum_{l=0}^{\infty} c_{\xi,l} \tau^l,$$

where we suppressed the dependence of $c_{\xi,l}$ on i, j in the notation. We emphasize that $\sum_{l=0}^{\infty} c_{\xi,l} \tau^l$ is an expansion of

$$(3.12) \quad \frac{1}{\tau^2} [\mathbf{Q}_\xi]_{ij} = \sum_{k=1}^d \langle (e_i + \nabla\phi_i)(e_k)(\xi + \nabla\phi_\xi)(e_k)\mathbf{b}'(e_k) \\ (1 + \mathcal{L})^{-1}\mathbf{b}'(e_k)(e_j + \nabla\phi_j)(e_k)(\xi + \nabla\phi_\xi)(e_k) \rangle,$$

where the only τ -dependent terms are those factors of $\eta + \nabla\phi_\eta$ with $\eta = e_i, e_j, \xi$. The following are simple calculations using the i.i.d. structure of the random coefficients.

We introduce some notation. Let $G(x, y)$ be the Green function of the discrete Laplacian $-\Delta$, and $\nabla\nabla G(x, y)$ be the Hessian matrix such that

$$[\nabla\nabla G(x, y)]_{ij} = \nabla_i \nabla_j G(x, y),$$

where ∇_i, ∇_j are with respect to the x, y variable respectively. For any edge $e = (\underline{e}, \bar{e}) \in \mathbb{B}$, we also write

$$\nabla\nabla G(e, y) = \nabla G(\bar{e}, y) - \nabla G(\underline{e}, y).$$

Second order. To get the second order term, we replace all factors in (3.4) of the form $\eta + \nabla\phi_\eta$ by $X_{0,\eta} = \eta$ to obtain

$$c_{\xi,0} = \sum_{k=1}^d \langle \delta_{ik} \xi_k \mathbf{b}'(e_k) (1 + \mathcal{L})^{-1} \mathbf{b}'(e_k) \delta_{jk} \xi_k \rangle \\ = \langle \mathbf{b}'(e) (1 + \mathcal{L})^{-1} \mathbf{b}'(e) \rangle \sum_{k=1}^d \delta_{ik} \delta_{jk} \xi_k^2 = \delta_{ij} \xi_i^2 \langle \mathbf{b}(e)^2 \rangle,$$

where we used [19, Proposition 3.1] in the last step.

Third order. From now on, we focus on the case $i \neq j$. To get the third order term, we replace one of the four factors of $\eta + \nabla\phi_\eta$ in (3.4) by $X_{1,\eta}$ and all other three by $X_{0,\eta} = \eta$. We write

$$X_{1,\eta}(x) = [-\tau\nabla(-\Delta)^{-1}\nabla^*\mathbf{b}\eta](x) \\ = -\tau \sum_{y \in \mathbb{Z}^d} \nabla\nabla G(x, y) \mathbf{b}(y) \eta,$$

which is a vector for any $x \in \mathbb{Z}^d$. We also write

$$X_{1,\eta}(e) = -\tau \sum_{y \in \mathbb{Z}^d} \nabla \nabla G(e, y) \cdot \mathbf{b}(y) \eta,$$

which is a scalar for any $e \in \mathbb{B}$. We claim that we only need to consider $e_i + \nabla \phi_i$ or $e_j + \nabla \phi_j$. Otherwise if we replace $\xi + \nabla \phi_\xi$ by $X_{1,\xi}$, i.e., in (3.4)

$$e_i(e_k) + \nabla \phi_i(e_k) \mapsto e_i(e_k), \quad e_j(e_k) + \nabla \phi_j(e_k) \mapsto e_j(e_k),$$

and

$$\xi(e_k) + \nabla \phi_\xi(e_k) \mapsto -\tau \sum_{y \in \mathbb{Z}^d} \nabla \nabla G(e_k, y) \cdot \mathbf{b}(y) \xi,$$

we observe that there is a factor of $e_i(e_k)e_j(e_k) = \delta_{ik}\delta_{jk} = 0$ since $i \neq j$. Therefore, we have

$$\begin{aligned} c_{\xi,1} &= -\xi_k^2 \sum_{k=1}^d \langle \sum_{y \in \mathbb{Z}^d} \nabla \nabla G(e_k, y) \cdot \mathbf{b}(y) e_i \mathbf{b}'(e_k) (1 + \mathcal{L})^{-1} \mathbf{b}'(e_k) \delta_{jk} \rangle \\ &\quad - \xi_k^2 \sum_{k=1}^d \langle \mathbf{b}'(e_k) \delta_{ik} (1 + \mathcal{L})^{-1} \mathbf{b}'(e_k) \sum_{y \in \mathbb{Z}^d} \nabla \nabla G(e_k, y) \cdot \mathbf{b}(y) e_j \rangle. \end{aligned}$$

For the first term on the r.h.s. of the above expression, we can write

$$\begin{aligned} &\langle \nabla \nabla G(e_k, y) \cdot \mathbf{b}(y) e_i \mathbf{b}'(e_k) (1 + \mathcal{L})^{-1} \mathbf{b}'(e_k) \delta_{jk} \rangle \\ &= \langle \nabla \nabla G(e_j, y) \cdot \mathbf{b}(y) e_i \mathbf{b}'(e_j) (1 + \mathcal{L})^{-1} \mathbf{b}'(e_j) \rangle \delta_{jk}. \end{aligned}$$

Since $i \neq j$, it is clear that $\nabla \nabla G(e_j, y) \cdot \mathbf{b}(y) e_i$ and $\mathbf{b}'(e_j) (1 + \mathcal{L})^{-1} \mathbf{b}'(e_j)$ are independent for all $y \in \mathbb{Z}^d$, so we have the expectation equals zero because \mathbf{b} has mean zero. The same discussion applies to the second term, thus

$$c_{\xi,1} = 0.$$

Fourth order. To get the fourth order term, we have multiple options. First we consider the case when we have one factor of $X_{2,\eta}$ coming from $\eta + \nabla \phi_\eta$. By the same reason as before, we can not choose $\eta = \xi$, otherwise we have a factor of $\delta_{ik}\delta_{jk} = 0$. If instead we consider $\eta = e_i$ (the same discussion applies to e_j) and write:

$$X_{2,e_i}(x) = \tau^2 \sum_{y,z \in \mathbb{Z}^d} \nabla \nabla G(x, y) \mathbf{b}(y) \nabla \nabla G(y, z) \mathbf{b}(z) e_i$$

for $x \in \mathbb{Z}^d$ or

$$X_{2,e_i}(e) = \tau^2 \sum_{y,z \in \mathbb{Z}^d} \nabla \nabla G(e, y) \cdot \mathbf{b}(y) \nabla \nabla G(y, z) \mathbf{b}(z) e_i$$

for $e \in \mathbb{B}$, then the contribution to $[\mathbf{Q}_\xi]_{ij}$ is

$$\xi_j^2 \tau^4 \langle \sum_{y,z \in \mathbb{Z}^d} \nabla \nabla G(e_j, y) \cdot \mathbf{b}(y) \nabla \nabla G(y, z) \mathbf{b}(z) e_i \mathbf{b}'(e_j) (1 + \mathcal{L})^{-1} \mathbf{b}'(e_j) \rangle.$$

By the same discussion as before, $\mathbf{b}(z) e_i$ is independent of $\mathbf{b}'(e_j) (1 + \mathcal{L})^{-1} \mathbf{b}'(e_j)$ for all $z \in \mathbb{Z}^d$ because $i \neq j$. Since $\nabla \nabla G(e_j, y) \cdot \mathbf{b}(y) \nabla \nabla G(y, z) \mathbf{b}(z) e_i$ is a linear combination of $\mathbf{b}_{kk}(y)$ and $\mathbf{b}_{ii}(z)$, the only nonzero contribution after taking the

expectation is when $y = z$ and $k = i$:

$$\begin{aligned} & \xi_j^2 \tau^4 \left\langle \sum_{y, z \in \mathbb{Z}^d} \nabla \nabla G(e_j, y) \cdot \mathbf{b}(y) \nabla \nabla G(y, z) \mathbf{b}(z) e_i \mathbf{b}'(e_j) (1 + \mathcal{L})^{-1} \mathbf{b}'(e_j) \right\rangle \\ &= \xi_j^2 \tau^4 \nabla_i \nabla_i G(0, 0) \sum_{y \in \mathbb{Z}^d} \nabla \nabla_i G(e_j, y) \langle \mathbf{b}_i^2(y) \mathbf{b}'(e_j) (1 + \mathcal{L})^{-1} \mathbf{b}'(e_j) \rangle \\ &= \xi_j^2 \tau^4 \nabla_i \nabla_i G(0, 0) \sum_{y \in \mathbb{Z}^d} \nabla \nabla_i G(e_j, y) \langle \mathbf{b}^2(e_i) \rangle \langle \mathbf{b}'(e_j) (1 + \mathcal{L})^{-1} \mathbf{b}'(e_j) \rangle \end{aligned}$$

Since $\nabla \nabla_i G(e_j, y)$ is a gradient, it follows that $\sum_{y \in \mathbb{Z}^d} \nabla \nabla_i G(e_j, y) = 0$.

Now we consider the case when we have two factors of $X_{1, \eta}$. Recall that

$$\begin{aligned} [\mathbf{Q}_\xi]_{ij} &= \sum_{k=1}^d \langle (e_i + \nabla \phi_i)(e_k) (\xi + \nabla \phi_\xi)(e_k) \mathbf{a}'(e_k) \\ & \quad (1 + \mathcal{L})^{-1} \mathbf{a}'(e_k) (e_j + \nabla \phi_j)(e_k) (\xi + \nabla \phi_\xi)(e_k) \rangle. \end{aligned}$$

By symmetry, we only need to consider the following cases:

$$\begin{aligned} (i) &= \sum_{k=1}^d \langle X_{1, e_i}(e_k) X_{1, \xi}(e_k) \mathbf{a}'(e_k) (1 + \mathcal{L})^{-1} \mathbf{a}'(e_k) \delta_{jk} \xi_k \rangle, \\ (ii) &= \sum_{k=1}^d \langle X_{1, e_i}(e_k) \mathbf{a}'(e_k) (1 + \mathcal{L})^{-1} \mathbf{a}'(e_k) X_{1, e_j}(e_k) \rangle \xi_k^2, \\ (iii) &= \sum_{k=1}^d \langle X_{1, e_i}(e_k) \mathbf{a}'(e_k) (1 + \mathcal{L})^{-1} \mathbf{a}'(e_k) X_{1, \xi}(e_k) \rangle \delta_{jk} \xi_k. \end{aligned}$$

Note that we do not consider replacing both factors of $\xi + \nabla \phi_\xi$ since it leads to a factor of $\delta_{ik} \delta_{jk} = 0$.

For (i), we write

$$X_{1, e_i}(e_k) X_{1, \xi}(e_k) = \tau^2 \sum_{y \in \mathbb{Z}^d} \sum_{z \in \mathbb{Z}^d} (\nabla \nabla G(e_k, y) \cdot \mathbf{b}(y) e_i) (\nabla \nabla G(e_k, z) \cdot \mathbf{b}(z) \xi),$$

and in order to get a nonzero contribution in (i), we only need the terms with $y = z$ and the factor ξ_i , i.e.,

$$\tau^2 \sum_{y \in \mathbb{Z}^d} |\nabla \nabla_i G(e_k, y)|^2 |\mathbf{b}_{ii}(y)|^2 \xi_i,$$

which implies

$$\begin{aligned} (i) &= \tau^4 \xi_i \xi_j \sum_{y \in \mathbb{Z}^d} |\nabla \nabla_i G(e_j, y)|^2 |\mathbf{b}_{ii}(y)|^2 \mathbf{b}'(e_j) (1 + \mathcal{L})^{-1} \mathbf{b}'(e_j) \\ &= \tau^4 \xi_i \xi_j \sum_{y \in \mathbb{Z}^d} |\nabla \nabla_i G(e_j, y)|^2 |\mathbf{b}(e_i)|^2 \langle \mathbf{b}'(e_j) (1 + \mathcal{L})^{-1} \mathbf{b}'(e_j) \rangle. \end{aligned}$$

In the last step, we used the fact that $\mathbf{a}'(e_j) (1 + \mathcal{L})^{-1} \mathbf{a}'(e_j)$ is independent of $\mathbf{b}_{ii}(y)$ for all $y \in \mathbb{Z}^d$.

For (ii), we write

$$\begin{aligned} & \langle X_{1, e_i}(e_k) \mathbf{a}'(e_k) (1 + \mathcal{L})^{-1} \mathbf{a}'(e_k) X_{1, e_j}(e_k) \rangle \\ &= \tau^4 \sum_{y, z \in \mathbb{Z}^d} \langle \nabla \nabla G(e_k, y) \cdot \mathbf{b}(y) e_i \mathbf{b}'(e_k) (1 + \mathcal{L})^{-1} \mathbf{b}'(e_k) \nabla \nabla G(e_k, z) \cdot \mathbf{b}(z) e_j \rangle, \end{aligned}$$

and it is clear that for any $k = 1, \dots, d$, we have an independent factor of $\mathbf{b}(y) e_i$ or $\mathbf{b}(z) e_j$ since $i \neq j$, thus the above expression equals to zero.

For (iii), we write

$$\begin{aligned} & \langle X_{1,e_i}(e_j) \mathbf{a}'(e_j) (1 + \mathcal{L})^{-1} \mathbf{a}'(e_j) X_{1,\xi}(e_j) \rangle \\ &= \tau^4 \sum_{y,z \in \mathbb{Z}^d} \langle \nabla \nabla G(e_j, y) \cdot \mathbf{b}(y) e_i \mathbf{b}'(e_j) (1 + \mathcal{L})^{-1} \mathbf{b}'(e_j) \nabla \nabla G(e_j, z) \cdot \mathbf{b}(z) \xi \rangle. \end{aligned}$$

Similarly, we need the terms with $y = z$ and the factor ξ_i , which gives

$$\begin{aligned} & \langle X_{1,e_i}(e_j) \mathbf{a}'(e_j) (1 + \mathcal{L})^{-1} \mathbf{a}'(e_j) X_{1,\xi}(e_j) \rangle \\ &= \xi_i \tau^4 \sum_{y \in \mathbb{Z}^d} \langle \nabla \nabla G(e_j, y) \cdot \mathbf{b}(y) e_i \mathbf{b}'(e_j) (1 + \mathcal{L})^{-1} \mathbf{b}'(e_j) \nabla \nabla G(e_j, y) \cdot \mathbf{b}(y) e_i \rangle, \end{aligned}$$

so

$$\begin{aligned} (iii) &= \xi_i \xi_j \tau^4 \sum_{y \in \mathbb{Z}^d} \langle \nabla \nabla G(e_j, y) \cdot \mathbf{b}(y) e_i \mathbf{b}'(e_j) (1 + \mathcal{L})^{-1} \mathbf{b}'(e_j) \nabla \nabla G(e_j, y) \cdot \mathbf{b}(y) e_i \rangle \\ &= \xi_i \xi_j \tau^4 \sum_{y \in \mathbb{Z}^d} |\nabla \nabla_i G(e_j, y)|^2 \langle \mathbf{b}_{ii}(y) \mathbf{b}'(e_j) (1 + \mathcal{L})^{-1} \mathbf{b}'(e_j) \mathbf{b}_{ii}(y) \rangle \\ &= \xi_i \xi_j \tau^4 \sum_{y \in \mathbb{Z}^d} |\nabla \nabla_i G(e_j, y)|^2 \langle \mathbf{b}(e_i) \mathbf{b}'(e_j) (1 + \mathcal{L})^{-1} \mathbf{b}'(e_j) \mathbf{b}(e_i) \rangle. \end{aligned}$$

To summarize, the above formal calculation shows that for $i \neq j$,

$$(3.13) \quad c_{\xi,0} = c_{\xi,1} = 0,$$

and

$$(3.14) \quad \begin{aligned} c_{\xi,2} &= \xi_i \xi_j \sum_{y \in \mathbb{Z}^d} |\nabla \nabla_i G(e_j, y)|^2 \langle \mathbf{b}(e_i) \mathbf{b}'(e_j) (1 + \mathcal{L})^{-1} \mathbf{b}'(e_j) \mathbf{b}(e_i) \rangle \\ &\quad + \xi_i \xi_j \sum_{y \in \mathbb{Z}^d} |\nabla \nabla_i G(e_j, y)|^2 \langle \mathbf{b}(e_i) \mathbf{b}'(e_j) (1 + \mathcal{L})^{-1} \mathbf{b}'(e_j) \mathbf{b}(e_i) \rangle, \end{aligned}$$

which is non-zero if $\xi_i \xi_j \neq 0$.

3.2. Proof of the expansion. Before proving the expansion rigorously, we introduce more notation. Let $\zeta = (\zeta_e)_{e \in \mathbb{B}} \in \Omega$ denote the sample point, and for $x \in \mathbb{Z}^d$, we define the shift operator τ_x on Ω by $(\tau_x \zeta)_e = \zeta_{x+e}$, where $x + e := (x + \underline{e}, x + \bar{e})$ is the edge obtained by shifting e by x . Since $\{\zeta_e\}_{e \in \mathbb{B}}$ are i.i.d., $\{\tau_x\}_{x \in \mathbb{Z}^d}$ is a group of measure-preserving transformations. We can define the operator

$$T_x f(\zeta) = f(\tau_x \zeta)$$

for any measurable function f on Ω , and the generators of T_x , denoted by $\{D_i\}_{i=1}^d$, are defined by $D_i f := T_{e_i} f - f$. The adjoint D_i^* is defined by $D_i^* f := T_{-e_i} f - f$. We denote the gradient on Ω by $D = (D_1, \dots, D_d)$ and the divergence $D^* F := \sum_{i=1}^d D_i^* F_i$ for $F : \Omega \rightarrow \mathbb{R}^d$. The norm in $L^p(\Omega)$ is denoted by $\|\cdot\|_p$.

Now we can formulate the corrector equation

$$(3.15) \quad \nabla^* \mathbf{a}(x) (\nabla \phi_\eta(x) + \eta) = 0$$

in the probability space as

$$(3.16) \quad D^* \mathbf{a}(D \phi_\eta + \eta) = 0.$$

We note that (3.16) holds almost surely in Ω and (3.15) holds on \mathbb{Z}^d for \mathbb{P} -a.e. ζ . From now on, with an abuse of notation, we regard \mathbf{a}, ϕ_η as functions on Ω with

$$\mathbf{a}(\zeta) = \text{diag}(\{a(\zeta_{e_i})\}_{i=1, \dots, d}).$$

It is clear that if $\zeta \mapsto \phi_\eta(\zeta)$ solves (3.16), then $x \mapsto \phi_\eta(\tau_x \zeta)$ solves (3.15).

Since $\mathbf{a} = \mathbf{I}_d + \tau \mathbf{b}$, (3.16) can be written as

$$-\Delta \phi_\eta = -\tau D^* \mathbf{b}(D \phi_\eta + \eta)$$

with $-\Delta = D^*D$ the Laplacian on the probability space, so formally

$$D\phi_\eta = -\tau D(-\Delta)^{-1}D^*\mathbf{b}(D\phi_\eta + \eta).$$

The operator $D(-\Delta)^{-1}D^*$ is defined by the following lemma.

Lemma 3.2. *Let $F = (F_i)_{i=1,\dots,d} \in L^2(\Omega, \mathbb{R}^d)$, $\phi_\lambda : \Omega \rightarrow \mathbb{R}$ solve*

$$(\lambda - \Delta)\phi_\lambda = D^*F,$$

*and $\Psi = (\Psi_i)_{i=1,\dots,d} = D(-\Delta)^{-1}D^*F$ be a weak limit of $D\phi_\lambda$ in $L^2(\Omega)$, then Ψ is the unique function in $L^2(\Omega, \mathbb{R}^d)$ that satisfies the following three properties:*

- (i) $\langle \Psi \rangle = 0$;
- (ii) $D_i\Psi_j = D_j\Psi_i$ for all $i, j = 1, \dots, d$;
- (iii) $D^*\Psi = D^*F$.

Furthermore, $\|\Psi\|_2 \leq \|F\|_2$, and $D\phi_\lambda \rightarrow \Psi$ in $L^2(\Omega)$ as $\lambda \rightarrow 0$.

Proof. It is a special case of [12, Theorem 3]. The uniqueness follows from [21, Theorem 2] with minor modifications. For the strong $L^2(\Omega)$ convergence, we only need to show the convergence of $\langle D\phi_\lambda \cdot D\phi_\lambda \rangle$. On one hand, we have

$$\lambda \langle \phi_\lambda^2 \rangle + \langle D\phi_\lambda \cdot D\phi_\lambda \rangle = \langle F \cdot D\phi_\lambda \rangle \rightarrow \langle F \cdot \Psi \rangle = \langle \Psi \cdot \Psi \rangle,$$

where the last step comes from $\langle \Psi \cdot DG \rangle = \langle F \cdot DG \rangle$ for any test function G . This implies

$$\limsup_{\lambda \rightarrow 0} \langle D\phi_\lambda \cdot D\phi_\lambda \rangle \leq \langle \Psi \cdot \Psi \rangle.$$

On the other hand,

$$\langle \Psi \cdot \Psi \rangle = \lim_{\lambda \rightarrow 0} \langle \Psi \cdot D\phi_\lambda \rangle \leq \liminf_{\lambda \rightarrow 0} \|\Psi\|_2 \|D\phi_\lambda\|_2.$$

Thus $\langle D\phi_\lambda \cdot D\phi_\lambda \rangle \rightarrow \langle \Psi \cdot \Psi \rangle$ as $\lambda \rightarrow 0$. The proof is complete. \square

To justify the expansion, we need L^4 rather than L^2 boundedness of the operator $D(-\Delta)^{-1}D^*$. The following lemma, whose proof is postponed to Appendix B, is the key to justify the expansion.

Lemma 3.3. *For any $p \in (1, \infty)$, there exists $C_p > 0$ such that*

$$\|D(\lambda - \Delta)^{-1}D^*F\|_p \leq C_p \|F\|_p$$

uniformly in $F \in L^p(\Omega, \mathbb{R}^d)$ and $\lambda \in [0, 1]$.

Proof of Theorem 3.1. We first rewrite (3.4) as

$$(3.17) \quad [\mathbb{Q}_\xi]_{ij} = \tau^2 \sum_{k=1}^d \langle (\delta_{ik} + D_k\phi_i)(\xi_k + D_k\phi_\xi)\mathbf{b}'(e_k) \rangle \\ (1 + \mathcal{L})^{-1}\mathbf{b}'(e_k)(\delta_{jk} + D_k\phi_j)(\xi_k + D_k\phi_\xi) \rangle.$$

Recall that $D\phi_\eta$ solves

$$(\mathbf{I}_d - \tau\mathcal{P})D\phi_\eta = \tau\mathcal{P}\eta,$$

with

$$\mathcal{P} := -D(-\Delta)^{-1}D^*\mathbf{b}.$$

By Lemma 3.3 and the fact that $\mathbf{b} \in L^\infty(\Omega, \mathbb{R}^{d \times d})$, for any $p \in (1, \infty)$ and $n \in \mathbb{N}_+$ we have

$$\|\mathcal{P}^n \eta\|_p \leq C_p^n \|\eta\|_\infty$$

for some constant $C_p > 0$, so

$$D\phi_\eta = (\tau\mathcal{P} + \tau^2\mathcal{P}^2 + \dots)\eta,$$

where the convergence is in $L^p(\Omega, \mathbb{R}^d)$ for any $p \in (1, \infty)$, provided that $\tau < 1/C_p$. Therefore, using the fact that $(1 + \mathcal{L})^{-1}$ is bounded from $L^q(\Omega)$ to itself for any

$q \geq 2$ [19, Proposition 3.2] and $\mathbf{b}' \in L^\infty(\Omega, \mathbb{R}^{d \times d})$, we can replace the factors $\eta_k + D_k \phi_\eta$ with $\eta = e_i, e_j, \xi$ in (3.17) by the series

$$\eta_k + D_k \phi_\eta = \sum_{n=0}^{\infty} \tau^n (\mathcal{P}^n \eta)_k,$$

where $(\mathcal{P}^n \eta)_k$ denotes the k -th component of $\mathcal{P}^n \eta$ with $\mathcal{P}^0 = \mathbf{I}_d$. Notice that we used Lemma 3.3 for $p \geq 4$ and $\lambda = 0$. In other words, we obtain an expansion of \mathbf{Q}_ξ in terms of τ :

$$[\mathbf{Q}_\xi]_{ij} = \sum_{n_1, n_2, n_3, n_4=0}^{\infty} \tau^{2+n_1+n_2+n_3+n_4} Q_{n_1, n_2, n_3, n_4}(i, j, \xi)$$

with

$$Q_{n_1, n_2, n_3, n_4}(i, j, \xi) := \sum_{k=1}^d \langle (\mathcal{P}^{n_1} e_i)_k (\mathcal{P}^{n_2} \xi)_k \mathbf{b}'(e_k) (1 + \mathcal{L})^{-1} \mathbf{b}'(e_k) (\mathcal{P}^{n_3} e_j)_k (\mathcal{P}^{n_4} \xi)_k \rangle.$$

Thus, $[\mathbf{Q}_\xi]_{ij} = \tau^2 \sum_{l=0}^{\infty} c_{\xi, l} \tau^l$ with

$$c_{\xi, l} = \sum_{n_1 + \dots + n_4 = l} Q_{n_1, n_2, n_3, n_4}(i, j, \xi).$$

To compute $c_{\xi, l}$ explicitly, we define

$$\mathcal{P}_\lambda := -D(\lambda - \Delta)^{-1} D^* \mathbf{b}$$

and

(3.18)

$$Q_{n_1, n_2, n_3, n_4}^\lambda(i, j, \xi) := \sum_{k=1}^d \langle (\mathcal{P}_\lambda^{n_1} e_i)_k (\mathcal{P}_\lambda^{n_2} \xi)_k \mathbf{b}'(e_k) (1 + \mathcal{L})^{-1} \mathbf{b}'(e_k) (\mathcal{P}_\lambda^{n_3} e_j)_k (\mathcal{P}_\lambda^{n_4} \xi)_k \rangle.$$

By Lemmas 3.2 and 3.3, it follows that

$$Q_{n_1, n_2, n_3, n_4}^\lambda(i, j, \xi) \rightarrow Q_{n_1, n_2, n_3, n_4}(i, j, \xi)$$

as $\lambda \rightarrow 0$, and we have

$$(3.19) \quad c_{\xi, l} = \sum_{n_1 + \dots + n_4 = l} \lim_{\lambda \rightarrow 0} Q_{n_1, n_2, n_3, n_4}^\lambda(i, j, \xi).$$

With the mass regularization, we can write \mathcal{P}_λ^n in the physical domain

$$(3.20) \quad \begin{aligned} \mathcal{P}_\lambda^n \eta &= (-D(\lambda - \Delta)^{-1} D^* \mathbf{b})^n \xi \\ &= (-1)^n \sum_{y_1, \dots, y_n \in \mathbb{Z}^d} \left(\prod_{k=1}^n \nabla \nabla G_\lambda(y_{k-1}, y_k) \mathbf{b}(\tau_{y_k} \zeta) \right) \eta, \end{aligned}$$

with $y_0 = 0$ and G_λ the Green function of $\lambda - \Delta$ on \mathbb{Z}^d . By plugging (3.20) into (3.18) with $n = n_1, \dots, n_4$ and $\eta = e_i, e_j, \xi$, the expectation can be computed explicitly using the i.i.d. structure of $\{\zeta_e\}_{e \in \mathbb{B}}$.

To show that \mathbf{Q}_ξ may not be a multiple of identity, we only need to repeat the formal calculation in Section 3.1 verbatim with G replaced by G_λ to derive that

$$[\mathbf{Q}_\xi]_{ij} = c_{\xi, 2} \tau^4 + O(\tau^5)$$

with $c_{\xi, 2}$ given by (3.14). The proof is complete. \square

APPENDIX A. A LINEAR ALGEBRA LEMMA

Lemma A.1. *Let A be a symmetric, positive-definite matrix. The following statements are equivalent:*

(i) *there exists a symmetric matrix B such that*

$$\text{for all } x \in \mathbb{R}^n, (x^t A x)^2 = |x|^2 x^t B x$$

(ii) *the matrix A is a multiple of identity.*

Proof. The implication (ii) \implies (i) is obvious. We prove the converse implication by induction on the size n of the matrix. Letting A be an $n \times n$ matrix and assuming that the result holds for matrices of size $n - 1$, we can choose $x = (x_1, \dots, x_{n-1}, 0)$ to obtain $A_{ij} = c\delta_{ij}$ and $B_{ij} = c^2\delta_{ij}$ for some constant $c > 0$ with $i, j = 1, \dots, n - 1$. Now, for any $i < n$, by considering the coefficients of $x_i^3 x_n$ and $x_i x_n^3$, we have

$$2cA_{in} = B_{in}, \quad 2A_{nn}A_{in} = B_{in},$$

which implies

$$(A.1) \quad A_{in}(A_{nn} - c) = 0.$$

By considering the coefficients of $x_i^2 x_n^2$ and x_n^4 , we have

$$2cA_{nn} + A_{in}^2 = c^2 + B_{nn}, \quad A_{nn}^2 = B_{nn},$$

which implies

$$(A.2) \quad (A_{nn} - c)^2 = A_{in}^2.$$

Combining (A.1) and (A.2), we have $A_{nn} = c, A_{in} = 0$. The proof is complete. \square

APPENDIX B. $D(\lambda - \Delta)^{-1}D^*$ IS BOUNDED FROM $L^p(\Omega)$ TO ITSELF

The goal here is to prove the boundedness of $D(\lambda - \Delta)^{-1}D^*$ from $L^p(\Omega)$ to itself, and we will borrow a deterministic estimate from [2].

For $L \in \mathbb{Z}_+$, let Γ_L be a square box

$$\Lambda_L = [-L, L]^d \cap \mathbb{Z}^d.$$

For $\tilde{F} \in L^1(\Lambda_L, \mathbb{R}^d)$, we define the following integral operator for $\lambda > 0$:

$$(B.1) \quad \mathcal{K}_{\lambda, L} \tilde{F} = \nabla(\lambda - \Delta)^{-1} \nabla^* \tilde{F} = \sum_{y \in \Lambda_L} \nabla \nabla G_\lambda(\cdot, y) \tilde{F}(y),$$

where G_λ is the Green function of $\lambda - \Delta$ on \mathbb{Z}^d . We claim that the following weak type-(1, 1) estimate holds:

$$(B.2) \quad |\{x \in \Lambda_L : |\mathcal{K}_{\lambda, L} \tilde{F}(x)| > \alpha\}| \leq \frac{C}{\alpha} \sum_{x \in \Lambda_L} |\tilde{F}(x)|$$

for some $C > 0$ independent of $\alpha, \lambda, L > 0$. In [2, Lemma 4.6], (B.2) was shown with G_λ in (B.1) replaced by the Green function of Laplacian with zero boundary condition in Λ_L . The same proof works in our case since the only ingredient we need to change in their proof is the following bound on the triple gradient of G_λ , which was given by [2, Lemma 4.9]:

$$(B.3) \quad |\nabla_{y,i} \nabla_{x,j} \nabla_{y,k} G_\lambda(x, y)| \leq C|x - y|^{-d-1}$$

for some $C > 0$ independent of $\lambda > 0, i, j, k = 1, \dots, d$.

The estimate (B.2) on the physical space can be lifted up to the probability space:

Lemma B.1. *For any $F \in L^1(\Omega, \mathbb{R}^d)$, we have*

$$\mathbb{P}(|D(\lambda - \Delta)^{-1}D^*F| > \alpha) \leq \frac{C}{\alpha} \langle |F| \rangle$$

for some constant $C > 0$ independent of $\lambda, \alpha > 0$.

Proof. We fix $\lambda, \alpha > 0$. First, let $\tilde{F}(x) = F(\tau_x\omega)$ for $x \in \mathbb{Z}^d$. Since $F \in L^1(\Omega, \mathbb{R}^d)$, for almost every $\omega \in \Omega$, we have $\tilde{F} \in L^1(\Lambda_L, \mathbb{R}^d)$ for any $L \in \mathbb{Z}_+$, so by (B.2)

$$\sum_{x \in \Lambda_L} 1_{|\mathcal{K}_{\lambda, L}\tilde{F}(x)| > \alpha} \leq \frac{C}{\alpha} \sum_{x \in \Lambda_L} |\tilde{F}(x)|.$$

Taking expectation on both sides, we derive

$$\frac{1}{|\Lambda_L|} \sum_{x \in \Lambda_L} \mathbb{P}(|\mathcal{K}_{\lambda, L}\tilde{F}(x)| > \alpha) \leq \frac{C}{\alpha} \langle |F| \rangle.$$

We can write

$$\begin{aligned} \frac{1}{|\Lambda_L|} \sum_{x \in \Lambda_L} \mathbb{P}(|\mathcal{K}_{\lambda, L}\tilde{F}(x)| > \alpha) &= \frac{1}{|\Lambda_L|} \sum_{x \in \Lambda_{L-\sqrt{L}}} \mathbb{P}(|\mathcal{K}_{\lambda, L}\tilde{F}(x)| > \alpha) \\ &\quad + \frac{1}{|\Lambda_L|} \sum_{x \in \Lambda_L \setminus \Lambda_{L-\sqrt{L}}} \mathbb{P}(|\mathcal{K}_{\lambda, L}\tilde{F}(x)| > \alpha) \\ &:= (i) + (ii). \end{aligned}$$

For (ii), it is clear that

$$(ii) \leq \frac{|\Lambda_L \setminus \Lambda_{L-\sqrt{L}}|}{|\Lambda_L|} \rightarrow 0$$

as $L \rightarrow \infty$. For (i), we have

$$\sum_{y \in \mathbb{Z}^d} \nabla \nabla G_\lambda(x, y) \tilde{F}(y) = \mathcal{K}_{\lambda, L}\tilde{F}(x) + \sum_{y \in \mathbb{Z}^d \setminus \Lambda_L} \nabla \nabla G_\lambda(x, y) \tilde{F}(y),$$

then

$$\begin{aligned} \mathbb{P}\left(\left|\sum_{y \in \mathbb{Z}^d} \nabla \nabla G_\lambda(x, y) \tilde{F}(y)\right| > 2\alpha\right) &\leq \mathbb{P}(|\mathcal{K}_{\lambda, L}\tilde{F}(x)| > \alpha) \\ &\quad + \mathbb{P}\left(\left|\sum_{y \in \mathbb{Z}^d \setminus \Lambda_L} \nabla \nabla G_\lambda(x, y) \tilde{F}(y)\right| > \alpha\right). \end{aligned}$$

By stationarity,

$$\mathbb{P}\left(\left|\sum_{y \in \mathbb{Z}^d} \nabla \nabla G_\lambda(x, y) \tilde{F}(y)\right| > 2\alpha\right) = \mathbb{P}(|D(\lambda - \Delta)^{-1}D^*F| > 2\alpha)$$

is independent of $x \in \mathbb{Z}^d$. For the summation outside Λ_L , we have

$$\begin{aligned} \mathbb{P}\left(\left|\sum_{y \in \mathbb{Z}^d \setminus \Lambda_L} \nabla \nabla G_\lambda(x, y) \tilde{F}(y)\right| > \alpha\right) &\leq \frac{1}{\alpha} \left\langle \sum_{y \in \mathbb{Z}^d \setminus \Lambda_L} |\nabla \nabla G_\lambda(x, y) \tilde{F}(y)| \right\rangle \\ &\leq \frac{1}{\alpha} \sum_{y \in \mathbb{Z}^d \setminus \Lambda_L} |\nabla \nabla G_\lambda(x, y)| \rightarrow 0 \end{aligned}$$

as $L \rightarrow \infty$, uniformly in $x \in \Lambda_{L-\sqrt{L}}$.

Now we have

$$\begin{aligned} &\frac{|\Lambda_{L-\sqrt{L}}|}{|\Lambda_L|} \mathbb{P}(|D(\lambda - \Delta)^{-1}D^*F| > 2\alpha) - \frac{1}{\alpha |\Lambda_L|} \sum_{x \in \Lambda_{L-\sqrt{L}}} \sum_{y \in \mathbb{Z}^d \setminus \Lambda_L} |\nabla \nabla G_\lambda(x, y)| \\ &\leq \frac{C}{\alpha} \langle |F| \rangle - (ii). \end{aligned}$$

By sending $L \rightarrow \infty$, we obtain

$$\mathbb{P}(|D(\lambda - \Delta)^{-1}D^*F| > 2\alpha) \leq \frac{C}{\alpha}\langle |F| \rangle,$$

which completes the proof. \square

Using Lemma B.1 and the fact that $D(\lambda - \Delta)^{-1}D^*$ is bounded from $L^2(\Omega, \mathbb{R}^d)$ to itself, we can apply the standard interpolation argument, e.g., [2, Theorem 4.4], to conclude that $D(\lambda - \Delta)^{-1}D^*$ is bounded from $L^p(\Omega, \mathbb{R}^d)$ to itself for any $p \in (1, \infty)$.

For the case $\lambda = 0$, we only need to note that for any $F \in L^2(\Omega, \mathbb{R}^d)$,

$$D(\lambda - \Delta)^{-1}D^*F \rightarrow D(-\Delta)^{-1}D^*F$$

in $L^2(\Omega, \mathbb{R}^d)$ as $\lambda \rightarrow 0$, so by applying Fatou's Lemma, we conclude $D(-\Delta)^{-1}D^*$ is bounded from $L^p(\Omega, \mathbb{R}^d)$ to itself. The proof of Lemma 3.3 is complete.

ACKNOWLEDGMENTS

We would like to thank the Mathematics Research Center of Stanford University for the hospitality during JCM's visit. Y.G. was partially supported by the NSF through DMS-1613301. We would like to thank the anonymous referee for his careful reading of the paper and helpful suggestions.

REFERENCES

- [1] N. Berger, M. Biskup. Quenched invariance principle for simple random walk on percolation clusters. *Probab. Theory Related Fields* **137** (1-2), 83-120 (2007).
- [2] M. Biskup, M. Salvi, and T. Wolff. A central limit theorem for the effective conductance: linear boundary data and small ellipticity contrasts. *Comm. Math. Phys.*, 328(2):701-731, 2014.
- [3] P. Courrège. Sur la forme intégrable-différentielle des opérateurs de C_k^∞ dans C satisfaisant au principe du maximum. In *Séminaire de Théorie du Potentiel, Dirigé par M. Brelot, G. Choquet et J. Deny, 1965/66*, tome 10, exposé 2, 1-38. Secrétariat mathématique, Paris, 1966.
- [4] E. B. Dynkin. Markov processes and random fields. *Bull. Amer. Math. Soc. (N.S.)*, 3(3):975-999, 1980.
- [5] M. Fukushima, Y. Ōshima, and M. Takeda. *Dirichlet forms and symmetric Markov processes*, volume 19 of *de Gruyter Studies in Mathematics*. Walter de Gruyter & Co., Berlin, 1994.
- [6] G. Giacomini, S. Olla, H. Spohn. Equilibrium fluctuations for $\nabla\varphi$ interface model. *Ann. Probab.* **29** (3), 1138-1172 (2001).
- [7] A. Gloria and F. Otto. An optimal variance estimate in stochastic homogenization of discrete elliptic equations. *Ann. Probab.*, 39(3):779-856, 2011.
- [8] Y. Gu and J.-C. Mourrat. Scaling limit of fluctuations in stochastic homogenization. *Multiscale Model. Simul.*, 14(1):452-481, 2016.
- [9] G. Kallianpur and V. Mandrekar. The Markov property for generalized Gaussian random fields. *Ann. Inst. Fourier (Grenoble)*, 24(2):vi, 143-167, 1974. Colloque International sur les Processus Gaussiens et les Distributions Aléatoires (Colloque Internat. du CNRS, No. 222, Strasbourg, 1973).
- [10] D. Koller and N. Friedman. *Probabilistic graphical models*. Adaptive Computation and Machine Learning. MIT Press, Cambridge, MA, 2009.
- [11] T. Kolsrud. On the Markov property for certain Gaussian random fields. *Probab. Theory Related Fields*, 74(3):393-402, 1987.
- [12] R. Künneemann. The diffusion limit for reversible jump processes on \mathbb{Z}^d with ergodic random bond conductivities. *Comm. Math. Phys.*, 90(1):27-68, 1983.
- [13] H. Künsch. Gaussian Markov random fields. *J. Fac. Sci. Univ. Tokyo Sect. IA Math.*, 26(1):53-73, 1979.
- [14] P. Lévy. *Processus Stochastiques et Mouvement Brownien. Suivi d'une note de M. Loève*. Gauthier-Villars, Paris, 1948.
- [15] H. P. McKean, Jr. Brownian motion with a several-dimensional time. *Teor. Veroyatnost. i Primenen.*, 8:357-378, 1963. Engl. transl. *Theor. Probability Appl.* 8:335-354, 1963.
- [16] J. Miller. Fluctuations for the Ginzburg-Landau $\nabla\phi$ interface model on a bounded domain. *Comm. Math. Phys.*, 308(3):591-639, 2011.
- [17] G. M. Molčan. Some problems connected with the Brownian motion of Lévy. *Teor. Veroyatnost. i Primenen.*, 12:747-755, 1967. Engl. transl. *Theor. Probability Appl.* 12:682-690, 1967.

- [18] J.-C. Mourrat and J. Nolen. Scaling limit of the corrector in stochastic homogenization. Preprint, arXiv:1502.07440 (2015).
- [19] J.-C. Mourrat and F. Otto. Correlation structure of the corrector in stochastic homogenization. *Ann. Probab.*, 44(5):3207–3233, 2016.
- [20] A. Naddaf, T. Spencer. On homogenization and scaling limit of some gradient perturbations of a massless free field. *Comm. Math. Phys.* **183** (1), 55–84 (1997).
- [21] G. C. Papanicolaou and S. R. S. Varadhan. Boundary value problems with rapidly oscillating random coefficients. In *Random fields, Vol. I, II (Esztergom, 1979)*, volume 27 of *Colloq. Math. Soc. János Bolyai*, pages 835–873. North-Holland, Amsterdam, 1981.
- [22] L. D. Pitt. A Markov property for Gaussian processes with a multidimensional parameter. *Arch. Rational Mech. Anal.*, 43:367–391, 1971.
- [23] D. Preiss and R. Kotecký. Markoff property of generalized random fields. In *Seventh Winter School on Abstract Analysis*, pages 61–66. Czechoslovak Academy of Sciences, Mathematical Institute, Prague, 1979.
- [24] M. Röckner. Generalized Markov fields and Dirichlet forms. *Acta Appl. Math.*, 3(3):285–311, 1985.
- [25] S. Sheffield. Gaussian free fields for mathematicians. *Probab. Theory Related Fields* **139** (3-4), 521-541 (2007).

(Yu Gu) DEPARTMENT OF MATHEMATICS, BUILDING 380, STANFORD UNIVERSITY, STANFORD, CA, 94305, USA

(Jean-Christophe Mourrat) ENS LYON, CNRS, 46 ALLÉE D'ITALIE, 69007 LYON, FRANCE