

ASYMPTOTICS OF FAST ROTATING DENSITY-DEPENDENT INCOMPRESSIBLE FLUIDS IN TWO SPACE DIMENSIONS

Francesco Fanelli¹ and Isabelle Gallagher²

¹ UNIVERSITÉ DE LYON, UNIVERSITÉ CLAUDE BERNARD LYON 1
Institut Camille Jordan – UMR 5208
fanelli@math.univ-lyon1.fr

² UNIVERSITÉ PARIS DIDEROT, SORBONNE PARIS CITÉ
Institut de Mathématiques de Jussieu-Paris Rive Gauche – UMR 7586
gallagher@math.univ-paris-diderot.fr

July 2, 2017

Abstract

In the present paper we study the fast rotation limit for viscous incompressible fluids with variable density, whose motion is influenced by the Coriolis force. We restrict our analysis to two dimensional flows. In the case when the initial density is a small perturbation of a constant state, we recover in the limit the convergence to the homogeneous incompressible Navier-Stokes equations (up to an additional term, due to density fluctuations). For general non-homogeneous fluids, the limit equations are instead linear, and the limit dynamics is described in terms of the vorticity and the density oscillation function: we lack enough regularity on the latter to prove convergence on the momentum equation itself. The proof of both results relies on a compensated compactness argument, which enables one to treat also the possible presence of vacuum.

2010 Mathematics Subject Classification: 35Q35 (primary); 35B25, 76U05, 35Q86, 35B40, 76M45 (secondary).

Keywords: incompressible fluids; Navier-Stokes equations; variable density; vacuum; Coriolis force; singular perturbation problem; low Rossby number.

1 Introduction

Given a small parameter $\varepsilon \in]0, 1]$, let us consider the non-homogeneous incompressible Navier-Stokes-Coriolis system

$$(1) \quad \begin{cases} \partial_t \rho + \operatorname{div}(\rho u) = 0 \\ \partial_t(\rho u) + \operatorname{div}(\rho u \otimes u) + \frac{1}{\varepsilon} \nabla \Pi + \frac{1}{\varepsilon} \mathfrak{C}(\rho, u) - \nu \Delta u = 0 \\ \operatorname{div} u = 0 \end{cases}$$

in a domain $\Omega \subset \mathbb{R}^d$, with $d = 2$ or 3 . This system describes the dynamics of a viscous incompressible fluid with variable density, and whose dynamics is influenced by the rotation of the ambient physical system: our main motivation concerns ocean currents on the Earth surface, but the system could also relate to flows on stars or other rotating celestial bodies. The scalar function $\rho \geq 0$ represents the density of the fluid, $u \in \mathbb{R}^d$ its velocity field and Π its pressure;

the term $\nabla\Pi$ can be interpreted as a Lagrangian multiplier associated with the incompressibility constraint $\operatorname{div} u = 0$. The parameter $\nu > 0$ is the viscosity coefficient, and we have chosen the very simple form Δu for the viscous stress tensor of the fluid. Finally, we have denoted by $\mathfrak{C}(\rho, u)$ the Coriolis operator, which represents the influence of the rotation on the fluid motion.

Operator \mathfrak{C} can assume various forms, depending on the precise assumptions and features one is interested in. In this paper we place ourselves at mid-latitudes, that is to say in a not too extended region of the Earth surface which is far enough from the poles and the equator, and we suppose the rotation axis to be constant. These are perhaps quite restrictive assumptions from the physical viewpoint, but they are usually assumed in mathematical studies (see e.g. [9]): indeed, the model one obtains is already able to describe several important features of ocean dynamics (for instance the Taylor-Proudman theorem, the formation of boundary layers...). Of course, more precise models can be considered, see e.g. [12] and [24], where inhomogeneities of the Coriolis force are taken into account, or [14], [15], [16], [23] and [25] for an analysis of the β -plane model for equatorial flows (where the Coriolis force vanishes). In the present paper, we restrict our attention to the case $d = 2$, for which the Coriolis operator takes the form

$$\mathfrak{C}(\rho, u) = \rho u^\perp,$$

where, for any vector $v = (v^1, v^2) \in \mathbb{R}^2$, we have defined $v^\perp := (-v^2, v^1)$. After adimensionalization, this term is multiplied by a factor $1/\operatorname{Ro}$, where the *Rossby number* Ro represents the inverse of the speed of the Earth rotation. Hence, the scaling introduced in (1) corresponds to taking $\operatorname{Ro} = \varepsilon$: our main goal here is to study the asymptotic behaviour of a family $(\rho_\varepsilon, u_\varepsilon)_\varepsilon$ of solutions to this system in the limit $\varepsilon \rightarrow 0$ and to characterize the limit dynamics. The presence of the factor $1/\varepsilon$ in front of the pressure term comes from the remark that, in the limit of fast rotation (i.e. for ε going to 0), the Coriolis term can be balanced only by $\nabla\Pi$, and so they must be of the same order (see [9] and Subsection 3.2 for more details about this point).

The incompressibility constraint is well justified in a first approximation for oceanic flows, see e.g. books [10], [26] and [32]. In the case of constant density, i.e. when (say) $\rho \equiv 1$ in (1), the fast rotation limit has been deeply investigated in the last two decades. We refer e.g. to [1], [2], [3], [8], [22], [24], among various contributions (the present list is far from being exhaustive). We refer to book [9] for an overview of the results and further references. It goes without saying that, both for modeling and application purposes, it is important to include in the model also other features of the fluid, like e.g. stratification, temperature variations, salinity... For systems allowing density fluctuations, mathematical studies are more recent, and to the best of our knowledge they concern only *compressible flows*: see e.g. [6], [23], [25] for 2-D viscous shallow water models, where the depth function plays the same role as the density (we refer to [14], [15], [16] for the inviscid case), see [19], [20] for 3-D barotropic Navier-Stokes equations (see also [17], [18] for the case when strong surface tension is taken into consideration). As is well-known, in the compressible (and barotropic) case the pressure is a given function of the density: then suitable scalings (roughly speaking, taking the Mach number of order higher than or equal to that of the Rossby number, with respect to the small parameter ε) enable one to find:

- on the one hand, good uniform bounds (by energy estimates) for the density fluctuation functions;
- on the other hand, a stream-function relation (by the analysis of the singular perturbation operator) linking the limit velocity and density profiles.

The combination of these two features is fundamental in the mathematical study, in order to handle the more difficult issues of the analysis (above all, at the convergence level). In the end, in the limit one obtains a quasi-geostrophic type equation for the limit of the density fluctuations, and this is enough to describe the whole asymptotic dynamics.

In the present paper we tackle instead the case of density-dependent fluids which are *incompressible*, which is (to the best of our knowledge) completely new in the mathematical literature. As a first approach, we restrict ourselves to the case $\Omega = \mathbb{R}^2$ or \mathbb{T}^2 : indeed, on the one hand these simple geometries allow us to neglect boundary layers effects; on the other hand, already in space dimension $d = 2$, the analysis presents several difficulties. We expect this study to be a first step in the understanding of the problem in its whole generality for three-dimensional domains. We consider two different frameworks: the former is the *slightly non-homogeneous case*, i.e. when the initial density is a small variation, of order ε , of a constant reference state; the latter is referred to as the *fully non-homogeneous case*, because the initial density is a perturbation of an arbitrary positive state. Let us remark that, in the latter instance, we are able to handle also the presence of possible vacuum.

At this point, let us make a physical *intermezzo* (see Chapter 3 of [10] or Appendix 3 of [26] for details). In the ocean, the water density is in general a complicated function of the pressure, temperature and salinity. Nonetheless, in many applications one usually neglects dependence on pressure and assumes linear dependence on the other two quantities. Moreover, in a first approximation it is usual to suppose that temperature and salinity evolve through a pure diffusion process (parabolic equations for each of these quantities). A final simplification, commonly adopted in physical models (see e.g. Sections 2.6 and 2.9 of [27] or Chapter 1 of [31]), is the so-called *Boussinesq approximation*: namely, the non-homogeneity of the fluid is supposed to be a small variation of a constant state $\bar{\rho}$ (without loss of generality, we can set $\bar{\rho} = 1$). In other words, one can write $\rho = 1 + \rho'$, where $|\rho'| \ll 1$: then the incompressibility constraint is directly derived from the mass conservation equation at the highest order. In addition, combining the equations for temperature (or energy) and salinity supplies an explicit evolution equation, of parabolic type, for the density variation function ρ' . We refer to Section 3.7 of [10] for more details.

In the light of the previous discussion, we notice here that the framework we adopt in this paper is “critical”, in the following sense:

- (i) first of all, the fluid is assumed to be incompressible, so that the pressure term is just a Lagrangian multiplier and does not provide any information on the density, unlike the case of compressible fluids;
- (ii) moreover our model, although being simplified since it does not take into account other quantities like energy and salinity, lacks information about the density fluctuations (i.e. the function ρ' mentioned above); even more, we do not want to restrict our attention only to small variations of constant states.

Thus our study of the density must rely on the mass conservation relation only. In addition, this is a hyperbolic type equation (pure transport by the velocity field), for which we can expect no gain of regularity: this fact is another source of difficulty in the analysis.

Nonetheless, the previous physical considerations suggest that the slightly non-homogeneous situation is somehow easier to be handled, and this turns out to be indeed the case. As a matter of fact, since the constant state is simply transported by a divergence-free velocity field, the density can be written as $\rho_\varepsilon = 1 + \rho'_\varepsilon = 1 + \varepsilon r_\varepsilon$, where, by pure transport again, good bounds on the variations r_ε can be derived. Then, an adaptation of the arguments of [24] enables us to pass to the limit directly in the mass and momentum equations: the limit dynamics is characterized by a coupling of a two-dimensional incompressible homogeneous Navier-Stokes system for the limit velocity field u , with a transport equation for the limit fluctuation function r by u itself. We point out that, in the momentum equation, an additional term ru^\perp appears: it can be interpreted as a remainder of the oscillations of the density. This fact can be seen by inserting the ansatz $\rho_\varepsilon = 1 + \varepsilon r_\varepsilon$ into the term $\mathfrak{C}(\rho_\varepsilon, u_\varepsilon)/\varepsilon$ and formally passing to the limit $\varepsilon \rightarrow 0$ (in particular, if $\rho_\varepsilon \equiv 1$ for all ε , then $r \equiv 0$ and this term disappears). The result about the slightly non-homogeneous case is given in Theorem 2.5, and is proved in Sections 3 and 4.

The fully non-homogeneous case is more involved; the corresponding result is given in Theorem 2.7, proved in Sections 3 to 5. Keep in mind points (i)-(ii) mentioned above: since now the reference density is no longer constant, we lack information on the density fluctuations (which, for the sake of clarity, will be called σ_ε in the fully non-homogeneous setting). This lack of information represents a major difficulty in our study; in order to overcome it, we resort to the vorticity formulation of the momentum equation, and combine it with the mass equation. This strategy provides a link between the vorticity η_ε of $v_\varepsilon := \rho_\varepsilon u_\varepsilon$ and σ_ε , which nonetheless reveals to be of partial help only, since we derive from it bounds for $(\sigma_\varepsilon)_\varepsilon$ in very rough spaces. Interpolating this information with another one coming from the mass equation allows nevertheless to grasp a key property in order to prove convergence in the general case (see Subsection 3.3). Strictly connected with this difficulty, another important issue in the analysis of the fully non-homogeneous framework is the control of oscillations, which is achieved by studying the corresponding system of waves, which we call *Rossby waves*. This terminology is typically used for fluctuations due to latitude (i.e., from the mathematical viewpoint, due to variations of the rotation axis), but in our context having a truly non-constant reference density produces analogous effects. Since, roughly speaking, in this case the singular perturbation operator has non-constant coefficients, in order to prove convergence we resort to a compensated compactness argument, introduced by P.-L. Lions and N. Masmoudi (see e.g. [30]) in the study of incompressible limit and first employed by the second author and L. Saint-Raymond in [24] in the context of rotating fluids with variations of the rotation axis. Let us mention that our compensated compactness argument is slightly different from the one used in e.g. [19] (for barotropic compressible Navier-Stokes equations), and it allows us to treat also the possible presence of vacuum regions.

This method enables us to prove the convergence of the convective term: similarly to [24] and [19], we deduce that the transport term vanishes in the limit of fast rotation, up to some remainders associated with an additional constraint which has to be satisfied by the limit density and velocity field. The explanation is that, as already remarked in the above mentioned papers, having a variable limit density imposes a strong constraint on the motion, which translates into the fact that the kernel of the penalized operator is smaller. As a consequence, the limit equations become linear: this remarkable fact can be interpreted as a sort of turbulent behaviour of the fluid, where all the scales are mixed and one can identify only an averaged dynamics. In our setting, this will be even more apparent: let us briefly justify this claim. Essentially due to the very weak information on the density variations σ_ε , we are not able to pass to the limit directly in the original formulation of our system: the major problems reside in the convergence of the Coriolis term and in the fact that we do not dispose of an explicit equation for the σ_ε 's. As a consequence, exploiting the above mentioned special relation linking σ_ε and η_ε , we are only able to prove convergence on the vorticity formulation of our system, which exactly mixes these two quantities. This is similar to what happens in the compressible case, but in our context we dispose of no relations linking the velocity field u_ε and the density variations: so this equation provides information only on the special combination of the limit quantities σ and η , but not on the dynamics of the limit velocity field u and σ separately. See also Remark 2.10 below for further comments about this point.

As a last comment, let us mention that, in Subsection 5.4, we will show a convergence result to the full system, where the dynamics of the limit density variation σ and of the limit velocity field u are decoupled. However, this statement is only a conditional convergence result, since it requires strong *a priori* bounds on the family of velocity fields. These bounds involve higher regularity than the one we can obtain by classical energy estimates, and is not propagated uniformly in ε in general, due to the singular behaviour of the Coriolis term.

Let us conclude this introduction with a short overview of the paper.

In the next section we formulate our assumptions and main results. In Section 3 we study the singular perturbation operator: namely we establish uniform bounds for the family $(\rho_\varepsilon, u_\varepsilon)_\varepsilon$ and the constraints their limit points have to satisfy. Section 4 is devoted to proving convergence

in the slightly non-homogeneous case, while in Section 5 we pass to the limit in the fully non-homogeneous framework. We postpone to Appendix A some tools from Littlewood-Paley theory and paradifferential calculus, which are needed in our analysis.

Acknowledgements

The work of the first author has been partially supported by the LABEX MILYON (ANR-10-LABX-0070) of Université de Lyon, within the program “Investissement d’Avenir” (ANR-11-IDEX-0007), and by the project BORDS, both operated by the French National Research Agency (ANR).

2 Assumptions and results

In the present section we introduce our working assumptions, and after discussing briefly the existence of weak solutions to our system, we formulate our main results.

2.1 Main assumptions

We state here the main assumptions on the initial data. For this and the next subsection (where we discuss the existence of weak solutions), we refer to Subsection 2.1 of [29] for more details.

Given a small parameter $\varepsilon \in]0, 1]$, let us consider, in the domain

$$\Omega := \mathbb{R}^2 \quad \text{or} \quad \mathbb{T}^2,$$

the non-homogeneous incompressible Navier-Stokes equations with Coriolis force

$$(2) \quad \begin{cases} \partial_t \rho + \operatorname{div}(\rho u) = 0 \\ \partial_t(\rho u) + \operatorname{div}(\rho u \otimes u) + \frac{1}{\varepsilon} \nabla \Pi + \frac{1}{\varepsilon} \rho u^\perp - \nu \Delta u = 0 \\ \operatorname{div} u = 0. \end{cases}$$

The scalar function $\rho \geq 0$ represents the density of the fluid and u is its velocity field; the scalar function Π is the pressure of the fluid; finally, the term ρu^\perp is due to the action of the Coriolis force on the fluid. We recall that the term $\nabla \Pi$ can be interpreted as the Lagrangian multiplier related to the incompressibility constraint $\operatorname{div} u = 0$.

For any fixed value of ε , we consider a weak solution $(\rho_\varepsilon, u_\varepsilon)$ to the previous system, in the sense specified by Definition 2.3 below. For this, we supplement system (2) with general *ill-prepared* initial data: for the density functions, we take

$$(3) \quad \rho_{0,\varepsilon} = \rho_0 + \varepsilon r_{0,\varepsilon}, \quad \text{with} \quad \rho_0 \in \mathcal{C}_b^2 \quad \text{and} \quad (r_{0,\varepsilon})_\varepsilon \subset L^2(\Omega) \cap L^\infty(\Omega).$$

Here and in the following, the notation $(a_\varepsilon)_\varepsilon \subset X$ (for some Banach space X) is to be understood as the fact that the sequence $(a_\varepsilon)_\varepsilon$ is uniformly bounded in X . The symbol \mathcal{C}_b^2 denotes the subspace of \mathcal{C}^2 functions which are bounded as well as their first and second order derivatives; this condition on ρ_0 can be somehow relaxed in the spirit of [18], but we assume it for simplicity.

Furthermore, we assume also that there exists a positive constant $\rho^* > 0$ such that

$$0 \leq \rho_0 \leq \rho^* \quad \text{and} \quad 0 \leq \rho_{0,\varepsilon} \leq 2\rho^*,$$

where ρ^* is a positive real number. The previous assumption means in particular that we are considering the possible presence of vacuum. Notice that the positivity requirement on ρ_0 is a sort of compatibility condition, due to the fact that $\rho_{0,\varepsilon} \rightarrow \rho_0$ when ε goes to 0. Of course, this imposes some conditions also on the sign of the perturbation functions $r_{0,\varepsilon}$ when ρ_0 vanishes, but this is not important at this stage.

Remark 2.1. Observe that, in the case when $\rho_0 \geq \rho_* > 0$ (and especially when $\rho_0 \equiv \rho_*$), with no loss of generality we can suppose that

$$0 < \rho_*/2 \leq \rho_{0,\varepsilon} \leq 2\rho^* \quad \text{for all } \varepsilon \in]0, 1].$$

For all ε , we also take an initial momentum $m_{0,\varepsilon}$ such that $m_{0,\varepsilon} = 0$ almost everywhere on the set $\{\rho_{0,\varepsilon} = 0\}$, and

$$(m_{0,\varepsilon})_\varepsilon \subset L^2(\Omega) \quad \text{and} \quad \left(|m_{0,\varepsilon}|^2/\rho_{0,\varepsilon}\right)_\varepsilon \subset L^1(\Omega),$$

where we agree that $|m_{0,\varepsilon}|^2/\rho_{0,\varepsilon} = 0$ on the set $\{\rho_{0,\varepsilon} = 0\}$.

Remark 2.2. In general, one would like to say that $m_{0,\varepsilon} = \rho_{0,\varepsilon} u_{0,\varepsilon}$. Nonetheless, conditions are imposed on $m_{0,\varepsilon}$ rather than on the velocity fields $u_{0,\varepsilon}$, since these fields may be not defined when the densities $\rho_{0,\varepsilon}$ vanish. For the same reason, we do not impose any divergence-free condition on the initial velocities. We refer to the discussion of [29] (see page 25 therein) for more details about this issue.

Of course, in the absence of vacuum (keep in mind Remark 2.1 above), the previous assumptions on $m_{0,\varepsilon}$ can be reformulated in an easier way, as conditions on the “true” velocity fields: we suppose then

$$(u_{0,\varepsilon})_\varepsilon \subset L^2(\Omega), \quad \text{with} \quad \operatorname{div} u_{0,\varepsilon} = 0.$$

Finally, in the case when $\Omega = \mathbb{R}^2$, we additionally require that there exists $\delta > 0$ such that, for all $\varepsilon \in]0, 1]$, one has the uniform embedding

$$(4) \quad \left(\frac{1}{\rho_{0,\varepsilon}} \mathbb{1}_{\{\rho_{0,\varepsilon} < \delta\}}\right)_\varepsilon \subset L^1(\Omega),$$

where $\mathbb{1}_A$ denotes the characteristic function of a set $A \subset \mathbb{R}^2$. Alternatively, we suppose that there exist a $\bar{\rho} > 0$ and a $p_0 \in]1, +\infty[$ such that

$$(5) \quad \left((\bar{\rho} - \rho_{0,\varepsilon})^+\right)_\varepsilon \subset L^{p_0}(\Omega),$$

where we have denoted by f^+ the positive part of a function f . We point out that other assumptions can be considered on the initial densities: we refer once again to Chapter 2 of [29].

Up to passing to subsequences, we can assume that

$$(6) \quad r_{0,\varepsilon} \xrightarrow{*} r_0 \quad \text{in } L^2(\Omega) \cap L^\infty(\Omega) \quad \text{and} \quad m_{0,\varepsilon} \rightharpoonup m_0 \quad \text{in } L^2(\Omega).$$

Of course, in absence of vacuum, we can suppose that

$$(7) \quad u_{0,\varepsilon} \rightharpoonup u_0 \quad \text{in } L^2(\Omega).$$

Before stating our main results, let discuss briefly energy estimates, and the existence of weak solutions for our equations.

2.2 Energy inequality, finite energy weak solutions

Suppose for a while that (ρ, u) are smooth solutions to system (2), related to smooth initial data (ρ_0, m_0) . First of all, we notice that the density is simply transported by a divergence-free velocity field: so all its L^q norms are preserved in time. On the other hand, system (2) has a conserved energy: for all $t > 0$

$$(8) \quad \left\| \sqrt{\rho(t)} u(t) \right\|_{L^2(\Omega)}^2 + 2\nu \int_0^t \|\nabla u(\tau)\|_{L^2(\Omega)}^2 d\tau \leq \| |m_0|^2 / \rho_0 \|_{L^1}.$$

Notice that, for smooth enough solutions, inequality (8) becomes actually an equality. We refer to Subsection 3.1 below for more details. However, the previous considerations motivate the following definition of *weak solution* to our system.

Definition 2.3. Fix initial data (ρ_0, m_0) such that the conditions listed in Subsection 2.1 above are fulfilled.

We say that (ρ, u) is a *weak solution* to system (2) in $]0, T[\times \Omega$ (for some time $T > 0$) with initial datum (ρ_0, m_0) if the following conditions are satisfied:

- (i) $\rho \in L^\infty([0, T[\times \Omega)$ and $\rho \in \mathcal{C}([0, T[; L^q_{\text{loc}}(\Omega))$ for all $1 \leq q < +\infty$;
- (ii) $\rho |u|^2 \in L^\infty([0, T[; L^1(\Omega))$, with $\nabla u \in L^2([0, T[\times \Omega)$ and $u \in L^2([0, T[\times \Omega)$;
- (iii) the mass equation is satisfied in a weak sense: for any $\phi \in \mathcal{D}([0, T[\times \Omega)$ one has

$$(9) \quad - \int_0^T \int_\Omega \left(\rho \partial_t \phi + \rho u \cdot \nabla \phi \right) dx dt = \int_\Omega \rho_0 \phi(0) dx;$$

- (iv) the divergence-free condition on u is satisfied in $\mathcal{D}'(]0, T[\times \Omega)$;
- (v) the momentum equation is satisfied in a weak sense: for any $\psi \in \mathcal{D}([0, T[\times \Omega)$ such that $\text{div } \psi = 0$, one has

$$(10) \quad \int_0^T \int_\Omega \left(-\rho u \cdot \partial_t \psi - \rho u \otimes u : \nabla \psi + \frac{1}{\varepsilon} \rho u^\perp \cdot \psi + \nu \nabla u : \nabla \psi \right) dx dt = \int_\Omega m_0 \cdot \psi(0) dx;$$

- (vi) for almost every $t \in]0, T[$, the energy inequality (8) holds true.

The solution is *global* if the previous conditions are satisfied for all $T > 0$.

Notice that point (vi) of the previous definition prescribes a finite energy condition on (ρ, u) , which are then solutions *à la Leray* (see [28]). Such a condition is the basis for the investigation of the singular perturbation problem, since most of the *a priori* bounds on our family of weak solutions will be derived from it.

The next result guarantees the existence of a global in time weak solution to the non-homogeneous Navier-Stokes equations with Coriolis force, for any fixed value of the parameter $\varepsilon \in]0, 1]$.

Theorem 2.4. Fix $\varepsilon \in]0, 1]$ and consider an initial datum (ρ_0, m_0) as in Definition 2.3 above. Then there exists a global in time weak solution (ρ, u) to equations (2).

For a proof of the previous statement, we refer to Chapter 2 of [29]. We remark that the presence of the Coriolis term can easily be handled: indeed, it vanishes identically in the energy estimates (since it is orthogonal to u), and passing to the limit in it in the compactness argument asks for a similar effort as in treating the time derivative, for instance.

2.3 Statement of the results

We now consider a family of initial data $(\rho_{0,\varepsilon}, m_{0,\varepsilon})_{\varepsilon \in]0, 1]}$ satisfying all the assumptions stated in Subsection 2.1 above. For any fixed $\varepsilon \in]0, 1]$, Theorem 2.4 provides a global in time weak solution $(\rho_\varepsilon, u_\varepsilon)$ to system (2). Our main purpose is to characterize the dynamics in the limit for ε going to 0, namely the system of equations verified by the limit points of the family $(\rho_\varepsilon, u_\varepsilon)_\varepsilon$.

It turns out that two different scenarios may occur:

- (i) the initial density is a small perturbation of a constant state, namely $\rho_0 \equiv 1$ (or another positive constant);
- (ii) the reference state of density is truly non-constant, meaning (roughly) that $\nabla \rho_0 \neq 0$: the precise assumption which is needed is given in condition (12) below.

In the former case, since the constant density configuration is transported by the velocity field, we are led to studying the incompressible and homogeneous limits simultaneously. Although the non-homogeneity $\rho_{0,\varepsilon}$ requires some effort in order to be handled, a not so difficult adaptation of the arguments used in [24] allows us to prove convergence to 2-D homogeneous incompressible Navier-Stokes equations. More precisely, we have the following statement, which will be proved in Section 4. Remark that we can formulate assumptions directly on the initial velocities $u_{0,\varepsilon}$ (recall Remark 2.2 above), since the density does not vanish.

Theorem 2.5. *Let us set $\rho_0 = 1$ and consider a family of initial data $(\rho_{0,\varepsilon}, u_{0,\varepsilon})_\varepsilon$ satisfying the assumptions fixed in Subsection 2.1. Let $(\rho_\varepsilon, u_\varepsilon)_\varepsilon$ be a family of corresponding weak solutions to system (2) in $\mathbb{R}_+ \times \Omega$, as given by Theorem 2.4. Let us define u_0 and r_0 as done in (6)-(7), and set $r_\varepsilon := \varepsilon^{-1}(\rho_\varepsilon - 1)$.*

Then, there exist $r \in L^\infty(\mathbb{R}_+; L^2(\Omega) \cap L^\infty(\Omega))$ and $u \in L^\infty(\mathbb{R}_+; L^2(\Omega)) \cap L^2_{\text{loc}}(\mathbb{R}_+; H^1(\Omega))$, with $\text{div } u = 0$ in the distributional sense, such that, up to the extraction of a subsequence, for any time $T > 0$ fixed, one has the following convergence properties:

- (a) $r_\varepsilon \xrightarrow{*} r$ in $L^\infty([0, T]; L^2(\Omega) \cap L^\infty(\Omega))$;
- (b) $u_\varepsilon \xrightarrow{*} u$ in $L^\infty([0, T]; L^2(\Omega)) \cap L^2([0, T]; H^1(\Omega))$.

Moreover, u is a global in time weak solution of the homogeneous Navier-Stokes equations with initial datum u_0 , while r satisfies (still in the weak sense) a pure transport equation by u , with initial datum r_0 . Namely, the following equations hold true in the weak sense:

$$(11) \quad \begin{cases} \partial_t r + \text{div}(r u) = 0 \\ \partial_t u + \text{div}(u \otimes u) + \nabla \Pi + r u^\perp - \nu \Delta u = 0 \\ \text{div } u = 0, \end{cases}$$

with initial data $r|_{t=0} = r_0$ and $u|_{t=0} = u_0$, for a suitable pressure function Π .

If, in addition, $r_0 \in H^{1+\beta}(\Omega)$, for some $\beta > 0$, and $u_0 \in H^1(\Omega)$, then the solution (r, u, Π) to system (11), related to the initial datum (r_0, u_0) , is unique. Hence, the convergence holds for the whole sequence $(r_\varepsilon, u_\varepsilon)_\varepsilon$.

The previous result is not really surprising, since it allows us to recover for slightly non-homogeneous fluids (almost) the same result as for homogeneous ones (see also Remark 2.6 below). However, its proof is interesting since it will suggest how to argue in the truly non-constant density case, where we work in a very low regularity framework.

Let us comment also on the uniqueness for the limit system. System (11) is studied in Subsection 4.4 below: Theorem 4.3 specifies the functional class where the solution is unique. We also show energy and stability estimates for those equations.

Remark 2.6. Compared to the homogeneous case, in the limit equation (11) we notice the presence of the additional term $r u^\perp$, which is essentially due to variations of the density, as explained in the Introduction. Notice that one can pass to the vorticity formulation of the previous equations: denoting the vorticity of u by $\omega := \text{curl } u := \partial_1 u^2 - \partial_2 u^1$ and using also the first relation in (11), we find

$$\partial_t(\omega - r) + u \cdot \nabla \omega - \nu \Delta \omega = 0.$$

Using the Biot-Savart law $u = -(-\Delta)^{-1} \nabla^\perp \omega$, we can remark the strong analogy of the previous equation with the one usually found for compressible fluids, see e.g. [19], [20] (see also [17] for a modified formulation, due to capillary effects in the limit). There, the vorticity formulation is more convenient in passing to the limit, since one disposes of the additional relation $u = \nabla^\perp r$ (namely r is a stream function for u), which gives $\omega = \Delta r$.

Let us now turn our attention to the case of an effectively variable reference density. Similarly to what happens for compressible fluids (see e.g. [19]), having a non-constant density in the limit dynamics imposes a strong constraint on weak-limit points of the family of solutions: the consequence of this fact is that the convective term vanishes in the limit and the final equations become linear. Analogous effects can be noticed in the case of variations of the rotation axis (see e.g. [24] and [18]).

For technical reasons, we need to assume that the reference density ρ_0 has non-degenerate critical points, in the sense of [24]. Namely, we suppose

$$(12) \quad \lim_{\delta \rightarrow 0} \mu \left(\left\{ x \in \Omega \mid |\nabla \rho_0(x)| \leq \delta \right\} \right) = 0,$$

where $\mu(\mathcal{O})$ denotes the Lebesgue measure of a set $\mathcal{O} \subset \mathbb{R}^2$.

Theorem 2.7. *Assume the reference density ρ_0 satisfies condition (12), and consider a family of initial data $(\rho_{0,\varepsilon}, u_{0,\varepsilon})_\varepsilon$ satisfying the assumptions fixed in Subsection 2.1. Let $(\rho_\varepsilon, u_\varepsilon)_\varepsilon$ be a family of corresponding weak solutions to system (2) in $\mathbb{R}_+ \times \Omega$, as given by Theorem 2.4. Let us define m_0 and r_0 as in (6), and set $\sigma_\varepsilon := \varepsilon^{-1}(\rho_\varepsilon - \rho_0)$.*

Then, there exist $\sigma \in L^\infty(\mathbb{R}_+; H^{-2-\delta}(\Omega))$, for $\delta > 0$ arbitrarily small, and a vector field u in the space $L^2_{\text{loc}}(\mathbb{R}_+; H^1(\Omega))$, with $\text{div } u = \text{div}(\rho_0 u) = 0$, such that, up to the extraction of a subsequence, for any fixed time $T > 0$, one has the following convergence properties:

- (a) $\rho_\varepsilon \xrightarrow{*} \rho_0$ in $L^\infty([0, T]; L^2(\Omega) \cap L^\infty(\Omega))$;
- (b) $u_\varepsilon \rightharpoonup u$ in $L^2([0, T]; H^1(\Omega))$;
- (c) $\sigma_\varepsilon \xrightarrow{*} \sigma$ in $L^\infty([0, T]; H^{-2-\delta}(\Omega))$ for all $\delta > 0$.

Moreover, set $\omega := \text{curl } u$ and $\eta := \text{curl}(\rho_0 u)$: then there exists a distribution $\Gamma \in \mathcal{D}'(\mathbb{R}_+ \times \Omega)$ such that the equation

$$(13) \quad \partial_t(\eta - \sigma) - \nu \Delta \omega + \text{curl}(\rho_0 \nabla \Gamma) = 0,$$

supplemented with the initial condition $(\eta - \sigma)|_{t=0} = \text{curl } m_0 - r_0$, is verified in the weak sense.

Remark 2.8. Let us stress the fact that we are able to prove a true weak convergence of system (2) to equation (13). More precisely, the convergence holds true whenever the mass and momentum equations in (2) are tested respectively on scalar test functions ϕ and on vector-valued divergence free test functions $\psi = \nabla^\perp \phi$ (without requiring any other constraint).

Remark 2.9. Let us point out that equation (13) can be formulated in terms of ω and σ only, using the following relation:

$$\eta = -\text{div}(\rho_0 \nabla(-\Delta)^{-1} \omega),$$

which easily follows from writing $\rho_0 u = \nabla^\perp \lambda_1$ and $u = \nabla^\perp \lambda_2$ (which both hold true thanks to the divergence-free conditions on the limit). Indeed, on the one hand, taking the curl of both equations yields $\eta = \Delta \lambda_1$ and $\omega = \Delta \lambda_2$; on the other hand, the relation $\rho_0 \nabla \lambda_2 = \nabla \lambda_1$ implies that $\text{div}(\rho_0 \nabla \lambda_2) = -\text{div}(\rho_0 \nabla(-\Delta)^{-1} \omega) = \Delta \lambda_1 = \eta$.

Remark 2.10. In terms of the limit density fluctuation σ and the limit velocity field u , the final equations (13) can be written formally as

$$(14) \quad \begin{cases} \partial_t \sigma + u \cdot \nabla \sigma = 0 \\ \rho_0 \partial_t u + \nabla \Pi + \rho_0 \nabla \Gamma + \sigma u^\perp - \nu \Delta u = 0 \\ \text{div } u = \text{div}(\rho_0 u) = 0. \end{cases}$$

We remark that the term $\rho_0 \nabla \Gamma$ in the second relation can be understood as the Lagrangian multiplier associated with the constraint $\operatorname{div}(\rho_0 u) = 0$. Roughly speaking, this term comes from the Coriolis operator when considered at the highest order, in the limit of fast rotation: see also Remark 3.5 about this point.

Nonetheless, contrary to the case of a constant reference density, here we do not have enough regularity on the density oscillations $(\sigma_\varepsilon)_\varepsilon$ to prove the convergence on the velocity equations and to find an equation for the σ_ε 's themselves. So the derivation of the previous system is just formal and, in order to rigorously pass to the limit, we are forced to resort to the vorticity formulation.

We postpone to Subsection 5.4 (see Theorem 5.8 therein) the statement of a conditional result where, under suitable assumptions, we are able to show convergence to (a modified version of) the full system (14).

3 Study of the singular perturbation

In the present section we establish important properties on the family of weak solutions we are considering. First we derive uniform bounds, and then we look for conditions their limit points have to satisfy; finally, we return to $(\rho_\varepsilon, u_\varepsilon)_\varepsilon$ and we infer additional properties, especially in the case of non-constant ρ_0 .

3.1 Uniform bounds

We establish here first uniform bounds on the family of weak solutions $(\rho_\varepsilon, u_\varepsilon)_\varepsilon$ to system (2). The formal arguments we use are fully justified for smooth solutions, and the bounds obtained pass to weak solutions thanks to a standard approximation procedure (see for instance Chapter 2 of [29]).

3.1.1 Bounds on the density functions

To begin with, let us focus on the density functions ρ_ε . By Theorem 2.1 of [29], their L^∞ bounds are preserved since they satisfy a pure transport equation by a divergence-free vector field. Therefore we get, for any $\varepsilon \in]0, 1]$ and for almost every $(t, x) \in \mathbb{R}_+ \times \Omega$,

$$(15) \quad 0 \leq \rho_\varepsilon(t, x) \leq 2\rho^*.$$

In particular, $(\rho_\varepsilon)_\varepsilon$ is uniformly bounded in the space $L^\infty(\mathbb{R}_+ \times \Omega)$, and hence one gathers the existence of a positive function $\rho \in L^\infty(\mathbb{R}_+ \times \Omega)$, which satisfies the same bounds as (15), and such that (up to the extraction of a subsequence)

$$(16) \quad \rho_\varepsilon \xrightarrow{*} \rho \quad \text{in} \quad L^\infty(\mathbb{R}_+; L^1_{\text{loc}}(\Omega) \cap L^\infty(\Omega)).$$

Let us focus for a while on the slightly non-homogeneous case, i.e. when $\rho_0 \equiv 1$. Defining the quantity $r_\varepsilon := \varepsilon^{-1}(\rho_\varepsilon - 1)$ as in the statement of Theorem 2.5 above, the mass equation can be rewritten as

$$(17) \quad \partial_t r_\varepsilon + \operatorname{div}(r_\varepsilon u_\varepsilon) = 0, \quad (r_\varepsilon)|_{t=0} = r_{0,\varepsilon}.$$

From this equation and the assumption on the initial data (3), we deduce the uniform bounds

$$(18) \quad (r_\varepsilon)_\varepsilon \subset L^\infty(\mathbb{R}_+; L^2(\Omega) \cap L^\infty(\Omega)),$$

and therefore there exists a $r \in L^\infty(\mathbb{R}_+; L^2(\Omega) \cap L^\infty(\Omega))$ such that, up to an extraction, $r_\varepsilon \xrightarrow{*} r$ in this space.

Let us now consider the general non-homogeneous case. Suppose that u_ε is smooth. Then we can introduce its flow $\psi_\varepsilon(t, x) \equiv \psi_{\varepsilon, t}(x)$, defined by the formula

$$\psi_{\varepsilon, t}(x) := x + \int_0^t u_\varepsilon(\tau, \psi_{\varepsilon, \tau}(x)) d\tau$$

for all $(t, x) \in \mathbb{R}_+ \times \Omega$. Then, by the transport equation for the density, we deduce that

$$\rho_\varepsilon(t, x) = \rho_{0, \varepsilon}(\psi_{\varepsilon, t}^{-1}(x)) = \tilde{\rho}_\varepsilon(t, x) + \varepsilon r_\varepsilon(t, x),$$

where, in analogy to what was done above, we have defined

$$\tilde{\rho}_\varepsilon(t, x) = \rho_0(\psi_{\varepsilon, t}^{-1}(x)) \quad \text{and} \quad r_\varepsilon(t, x) = r_{0, \varepsilon}(\psi_{\varepsilon, t}^{-1}(x)).$$

Namely both ρ_0 and $r_{0, \varepsilon}$ are transported by u_ε ; correspondingly, $\tilde{\rho}_\varepsilon$ and r_ε verify a pure transport equation $\partial_t a + u_\varepsilon \cdot \nabla a = 0$, from which we deduce

$$0 \leq \tilde{\rho}_\varepsilon \leq \rho^* \quad \text{and} \quad (r_\varepsilon)_\varepsilon \subset L^\infty(\mathbb{R}_+; L^2(\Omega) \cap L^\infty(\Omega)).$$

The previous uniform bounds are inherited by weak solutions, even if u_ε is no longer smooth. However, in light of Proposition 3.3 below, it turns out that, apart from the case $\rho_0 \equiv 1$ (and thus $\tilde{\rho}_\varepsilon(t) \equiv 1$ for any time $t \geq 0$), the decomposition $\rho_\varepsilon = \tilde{\rho}_\varepsilon + \varepsilon r_\varepsilon$ is not suitable for the analysis of the singular perturbation problem. We refer to Subsection 3.3 for more comments about this point.

3.1.2 Estimates for the velocity fields, and consequences

We turn now our attention to the momentum equation. Thanks to (8) we have the bounds

$$\left\| \sqrt{\rho_\varepsilon(t)} u_\varepsilon(t) \right\|_{L^2(\Omega)}^2 + 2\nu \int_0^t \|\nabla u_\varepsilon(\tau)\|_{L^2(\Omega)}^2 d\tau \leq C \| |m_{0, \varepsilon}|^2 / \rho_{0, \varepsilon} \|_{L^1}$$

for all $t > 0$. Notice that, by assumption on the initial data, the right-hand side of the previous inequality is uniformly bounded. Hence, keeping in mind property (15) on the density, we easily deduce that

$$(19) \quad (\sqrt{\rho_\varepsilon} u_\varepsilon)_\varepsilon \subset L^\infty(\mathbb{R}_+; L^2(\Omega)) \quad \text{and} \quad (\nabla u_\varepsilon)_\varepsilon \subset L^2(\mathbb{R}_+; L^2(\Omega)),$$

uniformly in $\varepsilon \in]0, 1]$.

Let us focus on the case $\Omega = \mathbb{R}^2$ for a while. Conditions (4) and (5) are preserved for ρ_ε , as a consequence of the pure transport equation by a divergence-free velocity field. So, referring to point 8 in Remark 2.1 of [29], we also infer the inclusion

$$(20) \quad (u_\varepsilon)_\varepsilon \subset L^2([0, T[\times \Omega)$$

for any fixed time $T > 0$. For notational convenience, in the following we denote, for any p and any Banach space X ,

$$L_T^p(X) := L^p([0, T[; X(\Omega)).$$

Then we deduce that $(u_\varepsilon)_\varepsilon$ is a bounded family in $L_T^2(H^1)$ for any $T > 0$. On the other hand, in the case $\Omega = \mathbb{T}^2$ this property follows by combining (19), Sobolev embeddings and Gagliardo-Nirenberg inequality (see Proposition A.8 in the Appendix).

Therefore, there exists $u \in L_{\text{loc}}^2(\mathbb{R}_+; H^1(\Omega))$ such that, up to extraction of a subsequence, there holds $u_\varepsilon \rightharpoonup u$ in this space, when $\varepsilon \rightarrow 0$.

Thanks to the previous properties, we can also establish strong convergence of the densities. More precisely, we write the mass equation under the form $\partial_t \rho_\varepsilon = -\operatorname{div}(\rho_\varepsilon u_\varepsilon)$: this implies that, for all fixed $T > 0$, $(\partial_t \rho_\varepsilon)_\varepsilon$ is uniformly bounded in $L^\infty([0, T]; H^{-1}(\Omega))$, and so $(\rho_\varepsilon)_\varepsilon$ is bounded in $W^{1, \infty}([0, T]; H_{\text{loc}}^{-1}(\Omega))$ uniformly in ε (the local condition in $H_{\text{loc}}^{-1}(\Omega)$ comes from the fact that $\rho_{0, \varepsilon}$ is just $L_{\text{loc}}^2(\Omega)$). Then, by the Ascoli-Arzelà Theorem we gather that the family $(\rho_\varepsilon)_\varepsilon$ is compact in $L^\infty([0, T]; H_{\text{loc}}^{-1}(\Omega))$, and hence, keeping in mind (16), by interpolation we deduce the strong convergence property (up to passing to a suitable subsequence)

$$(21) \quad \rho_\varepsilon \longrightarrow \rho \quad \text{in} \quad \mathcal{C}^{0, 1-\eta}([0, T]; H_{\text{loc}}^{-1+\eta}(\Omega))$$

for all $0 \leq \eta < 1$, where we have denoted by $\mathcal{C}^{0, \gamma}$ the space of γ -Hölder continuous functions, with $\mathcal{C}^{0, \gamma}$ replaced by $W^{1, \infty}$ when $\gamma = 1$.

Remark 3.1. We notice that a feature analogous to (21) can be deduced also on the family $(r_\varepsilon)_\varepsilon$, in the slightly non-homogeneous case.

Finally, let us establish a very simple convergence property. Since it will be repeatedly used in what follows, we choose to devote a statement to it.

Lemma 3.2. *Let $(\rho_\varepsilon, u_\varepsilon)_\varepsilon$ be a family of weak solutions to system (2), associated with initial data $(\rho_{0, \varepsilon}, u_{0, \varepsilon})$ satisfying the assumptions fixed in Subsection 2.1. Let (ρ, u) be a limit point of the sequence $(\rho_\varepsilon, u_\varepsilon)_\varepsilon$, as identified in Subsection 3.1.*

Then the product $(\rho_\varepsilon u_\varepsilon)_\varepsilon$ converges to ρu in the distributional sense, and more precisely in the weak topology of $L^2([0, T]; H_{\text{loc}}^{-\kappa}(\Omega))$, for all $0 < \kappa < 1$.

Proof. We have proved in (21) that the family $(\rho_\varepsilon)_\varepsilon$ is compact in e.g. $\mathcal{C}^{0, \eta}([0, T]; H_{\text{loc}}^{-\eta}(\Omega))$ for some $0 < \eta < 1$, and it strongly converges to ρ in the previous space. On the other hand, $(u_\varepsilon)_\varepsilon$ is weakly convergent in $L_T^2(H^1)$ to u . Therefore, by Corollary A.6 (ii) we get $\rho_\varepsilon u_\varepsilon \rightharpoonup \rho u$ in the space $L_T^2(H^{-\eta-\delta})$ and the result follows. \square

3.2 Constraints on the limit

In the previous part we have proved uniform bounds on the family of weak solutions $(\rho_\varepsilon, u_\varepsilon)_\varepsilon$, which allow us to identify (up to extraction) weak limits (ρ, u) . In the present subsection, we collect some properties these limit points have to satisfy. We point out that these conditions do not fully characterize the limit dynamics.

Proposition 3.3. *Let $(\rho_\varepsilon, u_\varepsilon)_\varepsilon$ be a family of weak solutions to system (2), associated with initial data $(\rho_{0, \varepsilon}, u_{0, \varepsilon})$ satisfying the assumptions fixed in Subsection 2.1. Let (ρ, u) be a limit point of the sequence $(\rho_\varepsilon, u_\varepsilon)_\varepsilon$, as identified in Subsection 3.1.*

Then one deduces the relations $\rho(t) \equiv \rho_0$ for all $t \geq 0$ and $\operatorname{div} u = \operatorname{div}(\rho_0 u) = 0$ in the sense of $\mathcal{D}'(\mathbb{R}_+ \times \Omega)$. In particular, we have that $u(t) \cdot \nabla \rho_0 = 0$ for almost every $t \geq 0$.

Proof. For $T > 0$ fixed, let us consider a smooth function $\psi \in \mathcal{D}([0, T[\times \Omega)$ such that $\operatorname{div} \psi = 0$, and let us test the momentum equation on $\varepsilon \psi$. We get

$$\begin{aligned} \int_0^T \int_\Omega \left(-\varepsilon \rho_\varepsilon u_\varepsilon \cdot \partial_t \psi - \varepsilon \rho_\varepsilon u_\varepsilon \otimes u_\varepsilon : \nabla \psi + \right. \\ \left. + \rho_\varepsilon u_\varepsilon^\perp \cdot \psi + \varepsilon \nu \nabla u_\varepsilon : \nabla \psi \right) dx dt = \varepsilon \int_\Omega \rho_{0, \varepsilon} u_{0, \varepsilon} \cdot \psi(0) dx. \end{aligned}$$

By inequalities established in Subsection 3.1, we know that $(\rho_\varepsilon u_\varepsilon)_\varepsilon$ is uniformly bounded in e.g. $L_T^\infty(L^2)$, and so are $(\rho_\varepsilon u_\varepsilon \otimes u_\varepsilon)_\varepsilon$ and $(\nabla u_\varepsilon)_\varepsilon$ respectively in $L_T^\infty(L^1)$ and in $L_T^2(L^2)$. Then, keeping in mind the assumptions on the initial data it is easy to see that, apart from the Coriolis operator, all the other terms in the previous relation converge to 0 in the limit $\varepsilon \rightarrow 0$.

Therefore in light of Lemma 3.2 above, we infer that

$$\lim_{\varepsilon \rightarrow 0} \int_0^T \int_{\Omega} \rho_{\varepsilon} u_{\varepsilon}^{\perp} \cdot \psi \, dx \, dt = \int_0^T \int_{\Omega} \rho u^{\perp} \cdot \psi \, dx \, dt = 0$$

for all test functions ψ such that $\operatorname{div} \psi = 0$. This property tells us that

$$(22) \quad \rho u^{\perp} = \nabla \pi,$$

for some suitable function π . Taking the curl of the previous relation, we immediately deduce that $\operatorname{div}(\rho u) = 0$. Obviously, the divergence-free condition on u , namely $\operatorname{div} u = 0$, also has to be satisfied in the weak sense.

Let us now consider the mass equation. Thanks to the previous bounds and Lemma 3.2, we can pass to the limit in its weak formulation with no difficulty: we thus find (still in the weak sense) the relation

$$\partial_t \rho + \operatorname{div}(\rho u) = 0,$$

with initial condition $\rho|_{t=0} = \rho_0$. But the constraint $\operatorname{div}(\rho u) = 0$ in \mathcal{D}' forces us to have $\partial_t \rho \equiv 0$, and therefore

$$\rho(t, x) \equiv \rho_0(x) \quad \text{for all } (t, x) \in \mathbb{R}_+ \times \Omega.$$

To conclude, since now we have enough regularity, we can unravel the property $\operatorname{div}(\rho_0 u) = 0$ and write it as $u \cdot \nabla \rho_0 = 0$. Hence, the last sentence of the statement follows by noting that $u \cdot \nabla \rho_0$ belongs to $L_T^2(H^1)$. \square

Before continuing our study, some remarks are in order.

Remark 3.4. For $\rho_0 = 1$, the only constraint on the limit velocity field is the relation $u^{\perp} = \nabla \pi$, which follows simply from the divergence-free condition. On the other hand, it is easy to pass to the limit in the equation for the r_{ε} 's (see also Section 4).

Remark 3.5. In the fully non-homogeneous case, in the limit $\varepsilon \rightarrow 0$, the Coriolis operator can be interpreted as a Lagrangian multiplier associated with the constraint $\operatorname{div}(\rho_0 u) = 0$.

3.3 Further properties and remarks

Before proving the convergence results, let us focus on the fully non-homogeneous case for a while, and establish further important properties for the density function.

3.3.1 The “good unknown” for the density

As already pointed out at the end of Paragraph 3.1.1, the decomposition $\rho_{\varepsilon} = \tilde{\rho}_{\varepsilon} + \varepsilon r_{\varepsilon}$, where $\tilde{\rho}_{\varepsilon}$ and r_{ε} are obtained transporting respectively ρ_0 and $r_{0,\varepsilon}$ by u_{ε} , is not suitable for our analysis.

In view of Proposition 3.3, the right *ansatz* to formulate is rather

$$\rho_{\varepsilon}(t, x) = \rho_0(x) + s_{\varepsilon}(t, x).$$

First of all, the family $(s_{\varepsilon})_{\varepsilon}$ is uniformly bounded in $L^{\infty}(\mathbb{R}_+ \times \Omega)$, because both $(\rho_{\varepsilon})_{\varepsilon}$ and ρ_0 are. Moreover, it satisfies (in the weak sense) the equation

$$(23) \quad \partial_t s_{\varepsilon} + \operatorname{div}(s_{\varepsilon} u_{\varepsilon}) = -\operatorname{div}(\rho_0 u_{\varepsilon}) = -u_{\varepsilon} \cdot \nabla \rho_0,$$

with initial datum $(s_{\varepsilon})|_{t=0} = \varepsilon r_{0,\varepsilon} \in L^2 \cap L^{\infty}$. From (19) and (20) we infer that $u_{\varepsilon} \cdot \nabla \rho_0$ is uniformly bounded, for any $T > 0$, in the space $L_T^2(L^2)$ and, by Sobolev embeddings, also in $L_T^2(L^q)$ for all $2 \leq q < +\infty$. Therefore, since $\operatorname{div} u_{\varepsilon} = 0$, we deduce the uniform bounds

$$(24) \quad (s_{\varepsilon})_{\varepsilon} \subset L^{\infty}([0, T]; L^2(\Omega) \cap L^{\infty}(\Omega))$$

for all $T > 0$.

Remark 3.6. Remark that $s_\varepsilon \equiv \varepsilon r_\varepsilon$ in the case $\rho_0 \equiv 1$; similar uniform bounds have already been established above for the r_ε 's, see (18) and Remark 3.1.

Furthermore, writing $\partial_t s_\varepsilon = -\operatorname{div}(\rho_\varepsilon u_\varepsilon)$, we deduce that $(s_\varepsilon)_\varepsilon \subset W_T^{1,\infty}(H^{-1}(\Omega))$. Therefore, on the one hand, arguing exactly as in the last part of Paragraph 3.1.2, we can establish strong convergence properties for $s_\varepsilon \rightarrow 0$ analogous to (21) (keep in mind also Proposition 3.3). On the other hand, we gather in particular the uniform embeddings

$$(25) \quad (s_\varepsilon)_\varepsilon \subset \mathcal{C}^{0,\gamma}([0, T]; H^{-\gamma}(\Omega))$$

for all $\gamma \in [0, 1]$.

Let us now define

$$(26) \quad \sigma_\varepsilon := \frac{1}{\varepsilon} s_\varepsilon, \quad V_\varepsilon := \rho_\varepsilon u_\varepsilon \quad \text{and} \quad f_\varepsilon := -\operatorname{div}(\rho_\varepsilon u_\varepsilon \otimes u_\varepsilon) + \nu \Delta u_\varepsilon.$$

We recall that, by uniform bounds, we have $(V_\varepsilon)_\varepsilon \subset L_T^\infty(L^2) \cap L_T^2(L^q)$ for all $2 \leq q < +\infty$. In the same way, we have $\rho_\varepsilon u_\varepsilon \otimes u_\varepsilon$ uniformly bounded in $L_T^2(L^a)$, with $1/a = 1/2 + 1/q$ and hence $a \in [1, 2[$. By duality of Sobolev embeddings, we have that $L^a \hookrightarrow H^{-\alpha}$, where $\alpha = 1 - 2/a'$ has to belong to $[0, 1[$, and $1/a + 1/a' = 1$. Hence we find $\alpha = 2(1/a - 1/2)$ for all $a \in]1, 2[$, that is to say we can write $\alpha = 2/q$ with $2 < q < +\infty$. From this we deduce that

$$(27) \quad (f_\varepsilon)_\varepsilon \subset L^2([0, T]; H^{-1-\alpha}(\Omega))$$

uniformly in $\varepsilon \in]0, 1[$, for any $\alpha \in]0, 1[$.

Remark 3.7. By (24), we know that, for each ε , σ_ε belongs to $L_T^\infty(L^2 \cap L^\infty)$, but *a priori* we have *no uniform bounds* at our disposal for these quantities.

With the previous notation, system (2) can be rewritten as

$$(28) \quad \begin{cases} \varepsilon \partial_t \sigma_\varepsilon + \operatorname{div} V_\varepsilon = 0 \\ \varepsilon \partial_t V_\varepsilon + \nabla \Pi_\varepsilon + V_\varepsilon^\perp = \varepsilon f_\varepsilon, \end{cases}$$

with initial data $(\sigma_\varepsilon)|_{t=0} = r_{0,\varepsilon} \in L^2 \cap L^\infty$ and $(V_\varepsilon)|_{t=0} = m_{0,\varepsilon} \in L^2$. Taking the curl of the second equation in the sense of distributions, and then computing the difference with the first one, we arrive at the relation

$$(29) \quad \partial_t(\eta_\varepsilon - \sigma_\varepsilon) = \operatorname{curl} f_\varepsilon,$$

where we have set $\eta_\varepsilon := \operatorname{curl}(V_\varepsilon) = \partial_1 V_\varepsilon^2 - \partial_2 V_\varepsilon^1$. In view of (27) and the properties on the initial data, we discover that $(\eta_\varepsilon - \sigma_\varepsilon)_\varepsilon$ is uniformly bounded in $\mathcal{C}_T^{0,1/2}(H^{-2-\alpha})$, and therefore, since by definition $(\eta_\varepsilon)_\varepsilon \subset L_T^\infty(H^{-1})$,

$$(30) \quad (\sigma_\varepsilon)_\varepsilon \subset L_T^\infty(H^{-2-\delta})$$

uniformly in ε , for all $\delta \in]0, 1[$. In particular, there exists a $\sigma \in L_{\text{loc}}^\infty(\mathbb{R}_+; H^{-2-\delta}(\Omega))$ such that $\sigma_\varepsilon \xrightarrow{*} \sigma$ in this space.

Remark 3.8. Property (30) is remarkable, because it establishes uniform bounds (even though in very negative spaces) for $(\rho_\varepsilon - \rho_0)/\varepsilon$. This uniform control is not obvious in the fully non-homogeneous case, if one just looks at the mass equation (pure transport); to find it, we have used rather deeply the structure of our system.

We are now going to exploit further this feature, and to establish a geometric property for the solutions to (2).

3.3.2 A geometric property for the velocity fields

In this section we want to derive another remarkable feature of the dynamics. Namely, dividing equation (23) by ε we can formally write

$$\partial_t \sigma_\varepsilon + \operatorname{div}(\sigma_\varepsilon u_\varepsilon) = -\frac{1}{\varepsilon} u_\varepsilon \cdot \nabla \rho_0.$$

Thus, we somehow expect that $u_\varepsilon \cdot \nabla \rho_0 \rightarrow 0$ (in some sense) for $\varepsilon \rightarrow 0$, with rate $O(\varepsilon)$. This is a strong geometric property for the family of u_ε 's, which asymptotically align along the direction given by $\nabla^\perp \rho_0$ (keep also in mind Proposition 3.3), with some rate of convergence. Unluckily, our bounds are not good enough to deduce that $\operatorname{div}(\sigma_\varepsilon u_\varepsilon)$ is uniformly controlled in a suitable space (keep in mind Remark 3.7 and property (30) above). Nonetheless, we are able to establish the following proposition, which will make this result a little more quantitative.

Proposition 3.9. *Given $\gamma \in]0, 1[$, there exist*

$$0 < \beta < \gamma, \quad \gamma < k < 1 \quad \text{and} \quad \theta \in]0, 1[$$

such that the uniform embeddings

$$\left(\frac{1}{\varepsilon^\theta} s_\varepsilon \right)_\varepsilon \subset \mathcal{C}^{0,\beta}([0, T]; H^{-k}(\Omega)) \quad \text{and} \quad \left(\frac{1}{\varepsilon^\theta} s_\varepsilon u_\varepsilon \right)_\varepsilon \subset L^2([0, T]; H^{-k-\delta}(\Omega))$$

hold true for all $\delta > 0$ arbitrarily small.

In particular, $(\varepsilon^{-\theta} s_\varepsilon)_\varepsilon \rightarrow 0$ (strong convergence) in $L^\infty([0, T]; H_{\text{loc}}^{-k-\delta}(\Omega))$ and $\varepsilon^{-\theta} s_\varepsilon u_\varepsilon \rightharpoonup 0$ (weak convergence) in $L^2([0, T]; H_{\text{loc}}^{-k-\delta}(\Omega))$, for all $\delta > 0$.

Proof. We start the proof by recalling that, thanks to (25), $(s_\varepsilon)_\varepsilon \subset \mathcal{C}_T^{0,\gamma}(H^{-\gamma}(\Omega))$ for all $\gamma \in [0, 1]$, while, by (30), $(\sigma_\varepsilon)_\varepsilon \subset L_T^\infty(H^{-2-\delta})$ for $\delta > 0$ arbitrarily small, where $\sigma_\varepsilon = s_\varepsilon/\varepsilon$.

Then, by interpolation of the previous uniform bounds, it is easy to find (just estimate the difference $\|s_\varepsilon(t) - s_\varepsilon(\tau)\|_{H^{-k}}$ for $0 \leq \tau < t \leq T$)

$$\|s_\varepsilon\|_{\mathcal{C}_T^{0,\beta}(H^{-k})} \leq C \|s_\varepsilon\|_{L_T^\infty(H^{-2-\lambda})}^\theta \|s_\varepsilon\|_{\mathcal{C}_T^{0,\gamma}(H^{-\gamma})}^{1-\theta},$$

under the conditions $\beta = (1-\theta)\gamma$ and $k = \theta(2+\lambda) + (1-\theta)\gamma$, for some $\theta \in]0, 1[$. For instance, one can make the explicit choices $\gamma = 1/2$, $\lambda = 1/n$ for some $n \in \mathbb{N}$ and $k = 3/4$, and determine consequently also the values of θ and β . The important point here is that there exist $0 < \beta < \gamma$ and $\gamma < k < 1$ and a suitable corresponding $\theta \in]0, 1[$, for which one has

$$\varepsilon^{1-\theta} \sigma_\varepsilon = \varepsilon^{-\theta} s_\varepsilon \in \mathcal{C}^{0,\beta}([0, T]; H^{-k}(\Omega)),$$

and this property holds uniformly for $0 < \varepsilon \leq 1$. In particular, by the Ascoli-Arzelà theorem, we deduce that this family is compact in e.g. $L^\infty([0, T]; H_{\text{loc}}^{-k-\lambda}(\Omega))$, for all $\lambda > 0$, and hence it converges strongly to 0 in this space.

Next, let us remark that, by Corollary A.6 (iii) of the Appendix, the product is continuous on $H^{-k} \times H^1$, with values in $H^{-k-\delta}$, for arbitrarily small $\delta > 0$. By these considerations, we immediately see that the product $\varepsilon^{-\theta} u_\varepsilon s_\varepsilon$ is well-defined and uniformly bounded in $L_T^2(H^{-k-\delta})$ with respect to ε , for all $\delta > 0$:

$$(\varepsilon^{-\theta} s_\varepsilon u_\varepsilon)_\varepsilon \subset L^2([0, T]; H^{-k-\delta}(\Omega)).$$

Furthermore, $\varepsilon^{-\theta} s_\varepsilon u_\varepsilon \rightharpoonup 0$ in the space $L_T^2(H_{\text{loc}}^{-k-\delta}(\Omega))$, by the strong convergence property established above on $(\varepsilon^{-\theta} s_\varepsilon)_\varepsilon$. \square

Let us end the discussion of this paragraph by pointing out further interesting properties, which however reveal to be not strong enough to be used in the analysis of the following sections.

Thanks to the properties established above, it makes sense (for instance in \mathcal{D}') to write the equation

$$\partial_t \left(\varepsilon^{-\theta} s_\varepsilon \right) + \operatorname{div} \left(\varepsilon^{-\theta} s_\varepsilon u_\varepsilon \right) = -\varepsilon^{-\theta} u_\varepsilon \cdot \nabla \rho_0,$$

which has to be interpreted in a weak sense: for all test functions $\phi \in \mathcal{D}([0, T[\times \Omega)$, one has

$$(31) \quad - \int_0^T \int_\Omega \varepsilon^{-\theta} s_\varepsilon \partial_t \phi - \int_0^T \int_\Omega \varepsilon^{-\theta} s_\varepsilon u_\varepsilon \cdot \nabla \phi = \int_\Omega \varepsilon^{1-\theta} r_{0,\varepsilon} \phi(0) - \int_0^T \int_\Omega \varepsilon^{-\theta} \rho_0 u_\varepsilon \cdot \nabla \phi.$$

By density and the uniform bounds on $\varepsilon^{-\theta} s_\varepsilon$ established above, we notice that this expression makes sense and is continuous actually for all $\phi \in W_T^{1,1}(H^{k+\delta+1})$, for any $\delta > 0$: regularity in time comes from the first term (notice that we are losing the gain of β derivatives in time for $\varepsilon^{-\theta} s_\varepsilon$), regularity in space (i.e. $k + \delta + 1$) from the second one in the left-hand side. Therefore we deduce the uniform bound

$$(32) \quad \left(\varepsilon^{-\theta} u_\varepsilon \cdot \nabla \rho_0 \right)_\varepsilon \subset W_T^{-1,\infty}(H^{-k}) + L_T^2(H^{-k-\delta-1}) \hookrightarrow W^{-1,\infty}([0, T]; H^{-k-\delta-1}(\Omega)),$$

for all $\delta > 0$ arbitrarily small. Actually, this term has to converge weakly to 0 in the previous space, because all the other terms in (31) go to 0 in \mathcal{D}' . So, we have gained a “dispersion” of order ε^θ (for some $\theta > 0$) for the quantity $(u_\varepsilon \cdot \nabla \rho_0)_\varepsilon$.

We will not use directly this property in the convergence proof (for which we refer to Section 5), but rather the uniform controls provided by Proposition 3.9.

As it may appear clear from the discussion of this subsection, the main difficulty in the analysis when ρ_0 is non-constant relies on the fact that we do not have at hand an explicit equation for the density oscillations σ_ε , nor do we have good enough uniform estimates for them: we are forced to use the vorticity associated with the velocity fields, and we get bounds in very rough spaces.

We will see in Section 5 how to handle these obstructions. For the time being, let us show the convergence in the slightly non-homogeneous case.

4 Combining homogeneous and fast rotation limits

We consider here the case $\rho_0 \equiv 1$, and we complete the proof of Theorem 2.5: namely, we prove a convergence result in the weak formulation of equations (2) and we identify the target system. We point out that, in this case, our fast rotation limit combines with a homogeneous limit, because $\rho_\varepsilon \rightarrow 1$ in the limit $\varepsilon \rightarrow 0$.

4.1 Preliminary convergence properties

When $\rho_0 \equiv 1$, we have an explicit equation (17) for the density oscillations $r_\varepsilon := (\rho_\varepsilon - 1)/\varepsilon$, from which we have deduced also suitable uniform bounds (18). Moreover, in light of Remark 3.1, one can argue exactly as in Lemma 3.2 and easily pass to the limit in the weak formulation of equation (17). Thus, for ε going to 0, we find

$$\partial_t r + \operatorname{div}(r u) = 0, \quad \text{with} \quad r|_{t=0} = r_0,$$

where r_0 is the limit point of the sequence $(r_{0,\varepsilon})_\varepsilon$ identified in (6).

Let us now consider the momentum equation. For $T > 0$ fixed, let us consider a smooth function $\psi \in \mathcal{D}([0, T[\times \Omega)$ such that $\operatorname{div} \psi = 0$. Using it in (10) we find

$$\int_0^T \int_\Omega \left(-\rho_\varepsilon u_\varepsilon \cdot \partial_t \psi - \rho_\varepsilon u_\varepsilon \otimes u_\varepsilon : \nabla \psi + \frac{1}{\varepsilon} \rho_\varepsilon u_\varepsilon^\perp \cdot \psi + \nu \nabla u_\varepsilon : \nabla \psi \right) dx dt = \int_\Omega \rho_{0,\varepsilon} u_{0,\varepsilon} \cdot \psi(0) dx.$$

Decomposing $\rho_\varepsilon = 1 + \varepsilon r_\varepsilon$ and making use of the uniform bounds established in Subsection 3.1, it is an easy matter to pass to the limit in the ∂_t term and in the viscosity term. Obviously, thanks to the assumptions on the initial data and especially properties (6), we have that $\rho_{0,\varepsilon} u_{0,\varepsilon} \rightharpoonup u_0$ in L^2 .

As for the Coriolis term, we can write

$$\frac{1}{\varepsilon} \int_0^T \int_\Omega \rho_\varepsilon u_\varepsilon^\perp \cdot \psi = \frac{1}{\varepsilon} \int_0^T \int_\Omega u_\varepsilon^\perp \cdot \psi + \int_0^T \int_\Omega r_\varepsilon u_\varepsilon^\perp \cdot \psi.$$

Now, keeping in mind Remark 3.4, we have that $\psi = \nabla^\perp \phi$: hence, the first term on the right-hand side identically vanishes (since u_ε is divergence-free), while the second converges, in view of Lemma 3.2 again. In the end, we find

$$\frac{1}{\varepsilon} \int_0^T \int_\Omega \rho_\varepsilon u_\varepsilon^\perp \cdot \psi \longrightarrow \int_0^T \int_\Omega r u^\perp \cdot \psi.$$

Notice that, alternatively, we could have used equation (17) tested against ϕ , to arrive at the relation $\varepsilon^{-1} \int_0^T \int_\Omega \rho_\varepsilon u_\varepsilon^\perp \cdot \psi = - \int_0^T \int_\Omega r_\varepsilon \partial_t \phi$. This would have led us to exploit the vorticity formulation of the momentum equation, see also Remark 2.6.

It remains to study the convergence of the convective term $\rho_\varepsilon u_\varepsilon \otimes u_\varepsilon$.

We proceed in three steps: first of all, since we are in the slightly non-homogeneous regime, we reduce our study to the constant density case. Then we approximate the velocities by smooth vector fields, which verify the same system up to small perturbations. These two steps are carried out in the next paragraph. Finally, in the third step we use a compensated compactness argument, as in [24]: we perform integrations by parts (which are possible because we are dealing now with smooth vector fields) and, using the structure of the equations, we are finally able to pass to the limit.

4.2 Some approximation lemmas

The following result is an obvious consequence of the fact that $\rho_\varepsilon = 1 + \varepsilon r_\varepsilon$, along with the uniform bounds for $(r_\varepsilon)_\varepsilon$ in $L_T^\infty(L^\infty)$, given by (18), combined with uniform bounds for $(u_\varepsilon)_\varepsilon$ in e.g. $L_T^\infty(L^2)$, see relation (19) above.

Lemma 4.1. *For any test function $\psi \in \mathcal{D}([0, T[\times \Omega)$, one has*

$$\lim_{\varepsilon \rightarrow 0^+} \left| \int_0^T \int_\Omega \rho_\varepsilon u_\varepsilon \otimes u_\varepsilon : \nabla \psi \, dx \, dt - \int_0^T \int_\Omega u_\varepsilon \otimes u_\varepsilon : \nabla \psi \, dx \, dt \right| = 0.$$

To perform the second step, for any $M \in \mathbb{N}$ we introduce the vector fields $u_{\varepsilon, M} := S_M u_\varepsilon$, where S_M is the low frequency cut-off operator defined by relation (62) in the Appendix. Notice that, for any fixed M , we have

$$(33) \quad \|u_{\varepsilon, M}\|_{L_T^\infty(H^k) \cap L_T^2(H^{k+1})} \leq C(T, k, M),$$

for any $T > 0$ and any $k > 0$. The constant $C(T, k, M)$ depends on T , k and M , but is uniform in $\varepsilon > 0$.

Lemma 4.2. *For any test function $\psi \in \mathcal{D}([0, T[\times \Omega)$, one has*

$$\lim_{M \rightarrow +\infty} \limsup_{\varepsilon \rightarrow 0^+} \left| \int_0^T \int_\Omega u_\varepsilon \otimes u_\varepsilon : \nabla \psi \, dx \, dt - \int_0^T \int_\Omega u_{\varepsilon, M} \otimes u_{\varepsilon, M} : \nabla \psi \, dx \, dt \right| = 0.$$

Proof. We start by estimating (see also (63) and Lemma A.4 below), for any $\delta > 0$, the difference

$$(34) \quad \|(\text{Id} - S_M)u_\varepsilon\|_{L_T^2(H^{1-\delta})} \leq 2^{-\delta M} \|u_\varepsilon\|_{L_T^2(H^1)} \leq C 2^{-\delta M},$$

for a constant $C > 0$, uniform in ε . Therefore, keeping in mind also bounds (19) for the family of velocity fields $(u_\varepsilon)_\varepsilon$, we gather that both integrals

$$I_1 := \int_0^T \int_\Omega (\text{Id} - S_M)u_\varepsilon \otimes u_\varepsilon : \nabla \psi \quad \text{and} \quad I_2 := \int_0^T \int_\Omega u_{\varepsilon,M} \otimes (\text{Id} - S_M)u_\varepsilon : \nabla \psi$$

converge to 0 for $M \rightarrow +\infty$, uniformly with respect to the parameter $\varepsilon \in]0, 1]$. This completes the proof of the statement. \square

4.3 The compensated compactness argument

From now on, the argument is absolutely analogous to the one used in [24]: let us sketch it for the reader's convenience. Thanks to Lemmas 4.1 and 4.2, we are reduced to studying, for any fixed $M \in \mathbb{N}$, the convergence (with respect to ε) of the integral

$$- \int_0^T \int_\Omega u_{\varepsilon,M} \otimes u_{\varepsilon,M} : \nabla \psi \, dx \, dt = \int_0^T \int_\Omega \text{div} (u_{\varepsilon,M} \otimes u_{\varepsilon,M}) \cdot \psi \, dx \, dt,$$

where we have integrated by parts since now each $u_{\varepsilon,M}$ is smooth in the space variable. Remark that, since partial derivatives commute with operator S_M , we have $\text{div} u_{\varepsilon,M} = 0$: then

$$(35) \quad \text{div} (u_{\varepsilon,M} \otimes u_{\varepsilon,M}) = u_{\varepsilon,M} \cdot \nabla u_{\varepsilon,M} = \frac{1}{2} \nabla |u_{\varepsilon,M}|^2 + \omega_{\varepsilon,M} u_{\varepsilon,M}^\perp,$$

where $\omega_{\varepsilon,M} := \text{curl} u_{\varepsilon,M}$ is the vorticity of $u_{\varepsilon,M}$. Notice that the former term in the last equality vanishes identically when tested against any test function ψ such that $\text{div} \psi = 0$.

As for the vorticity term, we start by reformulating the momentum equation in (2) in a similar way to what was done in Paragraph 3.3.1. More precisely, it is equivalent to write (still in the weak sense)

$$\varepsilon \partial_t V_\varepsilon + \nabla \Pi_\varepsilon + u_\varepsilon^\perp = \varepsilon f_\varepsilon + \varepsilon g_\varepsilon,$$

where V_ε and f_ε are defined in (26) and we have set $g_\varepsilon := -r_\varepsilon u_\varepsilon^\perp$. Remark that the family $(g_\varepsilon)_\varepsilon$ is uniformly bounded in $L_T^\infty(L^2)$. Applying the operator S_M to the previous equation we find

$$\varepsilon \partial_t V_{\varepsilon,M} + \nabla \Pi_{\varepsilon,M} + u_{\varepsilon,M}^\perp = \varepsilon \tilde{f}_{\varepsilon,M}, \quad \text{with} \quad \tilde{f}_\varepsilon := f_\varepsilon + g_\varepsilon.$$

We point out that, by uniform bounds and relation (27), for any M and k fixed and for all $T > 0$, one has

$$\|V_{\varepsilon,M}\|_{L_T^\infty(H^k)} + \|\tilde{f}_{\varepsilon,M}\|_{L_T^2(H^k)} \leq C(T, k, M),$$

where the constant $C(T, k, M)$ does not depend on ε . Furthermore, taking the curl of the equation and denoting $\eta_{\varepsilon,M} = \text{curl} V_{\varepsilon,M}$, we get

$$\partial_t \eta_{\varepsilon,M} = \text{curl} \tilde{f}_{\varepsilon,M}.$$

This last relation tells us that, for any fixed $M \in \mathbb{N}$ and $k \in \mathbb{R}$, the family $(\eta_{\varepsilon,M})_\varepsilon$ is compact (in ε) in e.g. $L_T^\infty(H_{\text{loc}}^k)$, and thus it converges strongly (up to extraction of a subsequence) to a tempered distribution η_M in this space. But since we already know the convergence $V_\varepsilon \rightharpoonup u$ in e.g. $L_T^\infty(L^2)$, it follows that $\eta_\varepsilon \rightharpoonup \omega = \text{curl} u$ in e.g. \mathcal{D}' , hence $\eta_M \equiv \omega_M$.

Finally writing $\rho_\varepsilon = 1 + \varepsilon r_\varepsilon$, we have the identity

$$\eta_{\varepsilon,M} = \text{curl} S_M(V_\varepsilon) = \omega_{\varepsilon,M} + \varepsilon \text{curl} S_M(r_\varepsilon u_\varepsilon),$$

where for any M and k , the family $(\text{curl } S_M(r_\varepsilon u_\varepsilon))_\varepsilon$ is uniformly bounded in $L_T^\infty(H^k)$. From this relation and the above analysis, we deduce the strong convergence (still up to an extraction)

$$\omega_{\varepsilon, M} \longrightarrow \omega_M \quad \text{in} \quad L_T^\infty(H_{\text{loc}}^k)$$

for $\varepsilon \rightarrow 0$. Using this property in equation (35), we see that, up to passing to a suitable subsequence, for any $T > 0$ and any $M \in \mathbb{N}$, we have the convergence

$$\lim_{\varepsilon \rightarrow 0} \int_0^T \int_\Omega u_{\varepsilon, M} \otimes u_{\varepsilon, M} : \nabla \psi \, dx \, dt = \int_0^T \int_\Omega \omega_M u_M^\perp \cdot \psi \, dx \, dt.$$

On the other hand, by uniform bounds and arguing as in (34), we have the strong convergence $u_M \rightarrow u$ in $L_T^2(H^1)$ in the limit for $M \rightarrow +\infty$ (by Lemma A.4 below and Lebesgue dominated convergence theorem). Therefore, performing equalities (35) backwards we can pass to the limit also in M . In the end, putting these properties together with Lemmas 4.1 and 4.2, we have shown that

$$\int_0^T \int_\Omega \rho_\varepsilon u_\varepsilon \otimes u_\varepsilon : \nabla \psi \, dx \, dt \longrightarrow \int_0^T \int_\Omega u \otimes u : \nabla \psi \, dx \, dt$$

for $\varepsilon \rightarrow 0$, for all smooth divergence-free test functions ψ .

In order to complete the proof of Theorem 2.5, it remains us to show that the whole sequence $(\rho_\varepsilon, u_\varepsilon)_\varepsilon$ converge. This fact is a straightforward consequence of a uniqueness property for the target system, which is established in the next subsection.

4.4 Study of the limit system

The analysis we have just carried out provides us with the existence of weak solutions to the target system (11). In this subsection we want to establish a uniqueness result for equations (11). In particular, this result will imply the convergence of the whole sequence in Theorem 2.5.

Let us remark that, in performing stability estimates, a loss of one derivative appears, due to the hyperbolicity of the “density” equation (i.e. the equation for r); now, propagating regularity for the gradient of r asks for additional smoothness on (r_0, u_0) . In this respect, although r and u are coupled via a zero order term, the system looks very much like a 2-D incompressible density-dependent Navier-Stokes equations, for which uniqueness can be established for more regular initial data (we refer for instance to Theorem 3.41 of [4] and to Proposition 5.1 of [11]).

For simplicity of the exposition, we will prove stability estimates in energy spaces, the extension to more general functional frameworks going beyond the scope of the present paper.

The main result of this subsection reads as follows.

Theorem 4.3. *Given $\beta > 0$, let $r_0 \in H^{1+\beta}(\Omega)$ and $u_0 \in H^1(\Omega)$ such that $\text{div } u_0 = 0$.*

Then there exists a unique weak solution (r, u) to system (11) with initial datum (r_0, u_0) , which satisfies the following properties:

- (i) *the density r belongs to $\mathcal{C}(\mathbb{R}_+; H^{1+\gamma}(\Omega))$ for all $0 \leq \gamma < \beta$;*
- (ii) *for all $T > 0$, u belongs to $\mathcal{C}([0, T]; H^1(\Omega)) \cap L^2([0, T]; H^2(\Omega))$.*

The rest of this section is devoted to the proof of the previous statement. More precisely, we will focus on the uniqueness part, the existence part being quite standard (as a byproduct of Theorem 2.5 and propagation of regularity); for the sake of completeness, we nevertheless show a priori estimates in Paragraph 4.4.1 before turning to the uniqueness result in Paragraph 4.4.2.

4.4.1 A priori estimates

In Theorem 4.3 we require additional smoothness on the initial data, compared to what can be expected from the singular limit problem. We do not give details about the existence of solutions at that level of regularity but focus on how to get *a priori* estimates for system (11).

First of all, since by assumption $r_0 \in L^2 \cap L^\infty$ is transported by a divergence-free velocity field, for all $t \geq 0$ we get

$$(36) \quad \|r(t)\|_{L^q} = \|r_0\|_{L^q} \quad \text{for all } q \in [2, +\infty].$$

As for u , we need to get higher regularity estimates: for this, we shall follow the same steps as for the proof of the existence of weak solutions to the non-homogeneous Navier-Stokes equations at H^1 level of regularity. We refer to pages 31-32 of [29] for details and original references of that result (see also comments in Theorem 3.41 of [4], and Proposition 5.1 of [11]). Actually, the situation here is simpler, since our equation for u is a homogeneous Navier-Stokes equation: r appears just on the Coriolis term, which can be treated as a forcing term at this level. As a consequence, we shall not attempt to prove refined estimates but only what is useful to infer the existence of a unique solution.

Before going on, let us point out that we will not keep track of the dependence of the various constants on the viscosity coefficient ν , which is positive and fixed.

First of all, if we multiply the second equation in (11) by u , we integrate over Ω and then in time, we immediately get

$$(37) \quad \|u(t)\|_{L^2}^2 + \nu \int_0^t \|\nabla u(\tau)\|_{L^2}^2 d\tau \leq C \|u_0\|_{L^2}^2.$$

Now let us multiply the equation by $-\Delta u$. We find, thanks to Hölder's inequality,

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|\nabla u\|_{L^2}^2 + \nu \|\Delta u\|_{L^2}^2 &= - \int u \cdot \nabla u \cdot (-\Delta u) dx - \int r u^\perp \cdot (-\Delta u) dx \\ &\leq \|u\|_{L^4} \|\nabla u\|_{L^4} \|\Delta u\|_{L^2} + \|r\|_{L^\infty} \|u\|_{L^2} \|\Delta u\|_{L^2}. \end{aligned}$$

By (36) and (37), along with Gagliardo-Nirenberg's inequality, we infer that

$$\frac{1}{2} \frac{d}{dt} \|\nabla u\|_{L^2}^2 + \frac{\nu}{2} \|\Delta u\|_{L^2}^2 \leq \|u_0\|_{L^2}^{1/2} \|\nabla u\|_{L^2} \|\Delta u\|_{L^2}^{3/2} + C \|r_0\|_{L^\infty}^2 \|u_0\|_{L^2}^2,$$

hence finally

$$\frac{d}{dt} \|\nabla u\|_{L^2}^2 + \nu \|\Delta u\|_{L^2}^2 \leq C \|u_0\|_{L^2}^2 \|\nabla u\|_{L^2}^4 + \frac{\nu}{2} \|\Delta u\|_{L^2}^2 + C \|r_0\|_{L^\infty}^2 \|u_0\|_{L^2}^2.$$

Gronwall's inequality gives, using (37) again,

$$(38) \quad \|\nabla u(t)\|_{L^2}^2 + \int_0^t \|\Delta u(\tau)\|_{L^2}^2 d\tau \leq C \left(\|\nabla u_0\|_{L^2}^2 + T \|r_0\|_{L^\infty}^2 \|u_0\|_{L^2}^2 \right) \exp(C \|u_0\|_{L^2}^4).$$

Now that we have obtained the property $u \in L^2([0, T]; H^2)$, we can apply Proposition 5.2 of [11] (see also Theorem 3.33 of [4]): we deduce that $r \in \mathcal{C}([0, T]; H^{1+\gamma})$ for all $\gamma < \beta$, and it satisfies the estimate

$$(39) \quad \begin{aligned} \|r(t)\|_{H^{1+\gamma}} &\leq C \exp \left(C \left(\int_0^t \|\nabla u\|_{H^1} d\tau \right)^2 \right) \|r_0\|_{H^{1+\beta}} \\ &\leq C \exp \left(C T \left(\|u_0\|_{L^2}^2 + (\|\nabla u_0\|_{L^2}^2 + T \|r_0\|_{L^\infty}^2 \|u_0\|_{L^2}^2) e^{C \|u_0\|_{L^2}^4} \right) \right) \|r_0\|_{H^{1+\beta}} \end{aligned}$$

for all $t \in [0, T]$, for a suitable constant C depending on β and γ .

These properties having been established, let us turn our attention to the *uniqueness* issue.

4.4.2 Stability estimates and uniqueness

Uniqueness of solutions to system (11) is an immediate consequence of the next proposition, which provides a stability estimate. In the proof, we limit ourselves to presenting the formal estimates, omitting a standard regularization procedure to make the computations rigorous.

Proposition 4.4. *Fix $\beta > 0$. Consider two initial densities $r_{0,1}$ and $r_{0,2}$ in $H^{1+\beta}(\Omega)$, and two initial divergence-free velocity fields $u_{0,1}$ and $u_{0,2}$ in $H^1(\Omega)$. Let (r_1, u_1) and (r_2, u_2) be two solutions to system (11) on $[0, T] \times \Omega$, related to the initial data $(r_{0,1}, u_{0,1})$ and $(r_{0,2}, u_{0,2})$ respectively. Define $\delta r := r_1 - r_2$ and $\delta u := u_1 - u_2$, and let δr_0 and δu_0 be the same quantities computed on the initial data.*

Then there exists a constant $C_0(T)$, depending just on the norms of the initial data and on the fixed time $T > 0$, such that the following estimates hold true for all $t \in [0, T]$:

$$\|\delta r(t)\|_{L^2}^2 + \|\delta u(t)\|_{H^1}^2 \leq C_0(T) e^{C_0(T)} (\|\delta r_0\|_{L^2}^2 + \|\delta u_0\|_{H^1}^2).$$

Remark 4.5. The H^1 assumption on the initial velocity field is needed in order to control ∇r . In fact, this term arises from stability estimates for the mass equation; it represents a loss of one derivative due to hyperbolicity of this equation. Imposing higher smoothness on u allows us to avoid the use of the L^∞ norm of ∇r , which has no chance of being bounded at this level of regularity (see inequality (40) below for further details). Note that the statement and its proof are far from being optimal but are enough for our purposes.

Proof of Proposition 4.4. Let us start by considering the transport equation for the density: taking the difference of the equation for r_1 by the equation for r_2 , we find that δr satisfies

$$\partial_t \delta r + u_2 \cdot \nabla \delta r = -\delta u \cdot \nabla r_1.$$

We now take the scalar product in L^2 of this equation with δr and integrate in time: easy computations lead to the estimate

$$\|\delta r(t)\|_{L^2}^2 \leq C \left(\|\delta r_0\|_{L^2}^2 + \int_0^t \|\delta u \cdot \nabla r_1\|_{L^2}^2 d\tau + \int_0^t \|\delta r\|_{L^2}^2 d\tau \right).$$

Thanks to Corollary A.6 (iii) we have

$$\|\delta u \cdot \nabla r_1\|_{L^2} \leq \|\delta u\|_{H^1} \|r_1\|_{H^{1+\beta/2}};$$

so we infer, for all $0 \leq t \leq T$, that

$$\begin{aligned} \|\delta r(t)\|_{L^2}^2 &\leq C \|\delta r_0\|_{L^2}^2 + C \int_0^t \|\delta u\|_{H^1}^2 \|r_1\|_{H^{1+\beta/2}}^2 d\tau + C \int_0^t \|\delta r\|_{L^2}^2 d\tau \\ &\leq C \|\delta r_0\|_{L^2}^2 + C_0(T) \int_0^t \|\delta u\|_{H^1}^2 d\tau + C \int_0^t \|\delta r\|_{L^2}^2 d\tau, \end{aligned}$$

thanks to (39), where $C_0(T)$ is a constant depending on norms of the initial data and on time $T > 0$, and can change from line to line. By Gronwall's inequality we obtain that, for all $0 \leq t \leq T$,

$$(40) \quad \|\delta r(t)\|_{L^2}^2 \leq \left(\|\delta r_0\|_{L^2}^2 + C_0(T) \int_0^t \|\delta u\|_{H^1}^2 d\tau \right) \exp(CT).$$

Let us focus now on the momentum equations: easy computations show that δu solves

$$(41) \quad \partial_t \delta u + u_2 \cdot \nabla \delta u + \delta u \cdot \nabla u_1 + \nabla \delta \Pi + r_2 \delta u^\perp + \delta r u_1^\perp - \nu \Delta \delta u = 0,$$

where $\delta \Pi := \Pi_1 - \Pi_2$. An L^2 energy estimate provides

$$\frac{1}{2} \frac{d}{dt} \|\delta u\|_{L^2}^2 + \nu \|\nabla \delta u\|_{L^2}^2 = - \int \delta u \cdot \nabla u_1 \cdot \delta u dx - \int \delta r u_1^\perp \cdot \delta u dx.$$

The terms on the right-hand side can be bounded in the following way:

$$\begin{aligned} \left| \int \delta r u_1^\perp \cdot \delta u \, dx \right| &\leq C \|\delta r\|_{L^2} \|u_1\|_{L^\infty} \|\delta u\|_{L^2} \leq C \|u_1\|_{L^\infty} (\|\delta r\|_{L^2}^2 + \|\delta u\|_{L^2}^2) \\ \left| \int \delta u \cdot \nabla u_1 \cdot \delta u \, dx \right| &\leq C \|\delta u\|_{L^4} \|u_1\|_{L^4} \|\nabla \delta u\|_{L^2} \leq C \|\delta u\|_{L^2}^{1/2} \|\nabla \delta u\|_{L^2}^{3/2} \|u_1\|_{L^4}, \end{aligned}$$

where, for the latter term, we have performed an integration by parts and we have used also Gagliardo-Nirenberg inequality. By Young's inequality, after an integration in time, it is easy to arrive at the estimate

$$\begin{aligned} \|\delta u(t)\|_{L^2}^2 + \nu \int_0^t \|\nabla \delta u(\tau)\|_{L^2}^2 \, d\tau \\ \leq \|\delta u_0\|_{L^2}^2 + C \int_0^t \|\delta u\|_{L^2}^2 \|u_1\|_{L^4}^4 \, d\tau + C \int_0^t (\|\delta r\|_{L^2}^2 + \|\delta u\|_{L^2}^2) \|u_1\|_{L^\infty} \, d\tau. \end{aligned}$$

Gronwall's lemma provides

$$\begin{aligned} \|\delta u(t)\|_{L^2}^2 + \int_0^t \|\nabla \delta u(\tau)\|_{L^2}^2 \, d\tau \leq C \left(\|\delta u_0\|_{L^2}^2 + \int_0^t \|\delta r\|_{L^2}^2 \|u_1\|_{L^\infty} \, d\tau \right) \\ \times \exp \left(C \int_0^T (\|u_1\|_{L^4}^4 + \|u_1\|_{L^\infty}) \, d\tau \right). \end{aligned}$$

Hence, using the fact that

$$\|u_1\|_{L^\infty} \leq C(\|u_1\|_{L^2} + \|u_1\|_{H^2}),$$

we infer

$$\|\delta u(t)\|_{L^2}^2 + \int_0^t \|\nabla \delta u(\tau)\|_{L^2}^2 \, d\tau \leq C \left(\|\delta u_0\|_{L^2}^2 + \int_0^t \|\delta r\|_{L^2}^2 (\|u_1\|_{L^2} + \|u_1\|_{H^2}) \, d\tau \right) e^{C_0(T)}.$$

It remains to plug that inequality into (40) to find

$$\begin{aligned} \|\delta r(t)\|_{L^2}^2 + \|\delta u(t)\|_{L^2}^2 + \int_0^t \|\nabla \delta u(\tau)\|_{L^2}^2 \, d\tau \\ \leq C_0(T) \left(\|\delta u_0\|_{L^2}^2 + \|\delta r_0\|_{L^2}^2 + \int_0^t \|\delta r\|_{L^2}^2 (\|u_1\|_{L^2} + \|u_1\|_{H^2}) \, d\tau + \int_0^t \|\delta u\|_{L^2}^2 \, d\tau \right) e^{C_0(T)}. \end{aligned}$$

Gronwall's lemma ends the proof of the proposition. \square

5 The fully non-homogeneous case

In this section we tackle the convergence for $\varepsilon \rightarrow 0$ in the case of a non-constant target density. Subsections 5.1 to 5.3 are devoted to the proof of Theorem 2.7. In Subsection 5.4 we present a conditional result, where we are able to show convergence to (a slightly modified version of) the full system (14).

We start by noticing that passing to the limit in the mass equation involves no special difficulty, and can be done as in the proof of Proposition 3.3. So we have to show convergence in the momentum equation, i.e. in the relation

$$\int_0^T \int_\Omega \left(-\rho_\varepsilon u_\varepsilon \cdot \partial_t \psi - \rho_\varepsilon u_\varepsilon \otimes u_\varepsilon : \nabla \psi + \frac{1}{\varepsilon} \rho_\varepsilon u_\varepsilon^\perp \cdot \psi + \nu \nabla u_\varepsilon : \nabla \psi \right) \, dx \, dt = \int_\Omega m_{0,\varepsilon} \cdot \psi(0) \, dx,$$

where ψ is a smooth divergence-free test function, compactly supported in $[0, T] \times \Omega$.

Once again, thanks to uniform bounds and Lemma 3.2, it is easy to perform the limit $\varepsilon \rightarrow 0$ in the time derivative and viscous terms. On the contrary, the Coriolis term involves some complications: let us focus on it for a while.

5.1 Passing to the vorticity formulation

For convenience, let us adopt the same notation as in (26) and write the previous expression under the form

$$(42) \quad \int_0^T \int_{\Omega} \left(-V_{\varepsilon} \cdot \partial_t \psi + \frac{1}{\varepsilon} V_{\varepsilon}^{\perp} \cdot \psi \right) dx dt = \int_0^T \langle f_{\varepsilon}, \psi \rangle dt + \int_{\Omega} V_{0,\varepsilon} \cdot \psi(0) dx,$$

where we denote by $\langle \cdot, \cdot \rangle$ the duality product in e.g. $H^{-1-\alpha} \times H^{1+\alpha}$ (recall relation (27) above) and we have set $V_{0,\varepsilon} := m_{0,\varepsilon}$. As before, ψ is a smooth test function having zero divergence.

Our concern here is to pass to the limit in the rotation term $\varepsilon^{-1} V_{\varepsilon}^{\perp}$. To do so, owing to the fact that $\operatorname{div} \psi = 0$, let us write $\psi = \nabla^{\perp} \phi$, for some smooth scalar function $\phi \in \mathcal{D}([0, T] \times \Omega)$. Using the mass equation tested against ϕ , we get (keep in mind (9) and (26) above)

$$\frac{1}{\varepsilon} \int_0^T \int_{\Omega} V_{\varepsilon}^{\perp} \cdot \psi = \frac{1}{\varepsilon} \int_0^T \int_{\Omega} V_{\varepsilon} \cdot \nabla \phi = - \int_0^T \int_{\Omega} \sigma_{\varepsilon} \partial_t \phi - \int_{\Omega} r_{0,\varepsilon} \phi(0).$$

On the one hand, this trick seems to us the only way to avoid the singularity of the Coriolis term. On the other hand, it forces us to pass to the vorticity formulation; hence, we rewrite (42) in the following form:

$$(43) \quad - \int_0^T \int_{\Omega} V_{\varepsilon} \cdot \partial_t \nabla^{\perp} \phi - \int_0^T \int_{\Omega} \sigma_{\varepsilon} \partial_t \phi = \int_0^T \langle f_{\varepsilon}, \nabla^{\perp} \phi \rangle + \int_{\Omega} V_{0,\varepsilon} \cdot \nabla^{\perp} \phi(0) + \int_{\Omega} r_{0,\varepsilon} \phi(0).$$

Notice that we have done nothing but finding again equation (29).

Our goal becomes then to pass to the limit in equation (43). Recalling the definition of f_{ε} given in (26) and the discussion at the beginning of the present section, the only difficulty relies in proving convergence in the convective term: so let us focus our attention on it.

5.2 Handling the convective term

In light of the previous discussion, it remains to pass to the limit in the convective term, namely in relation

$$\int_0^T \int_{\Omega} \rho_{\varepsilon} u_{\varepsilon} \otimes u_{\varepsilon} : \nabla \nabla^{\perp} \phi dx dt.$$

The strategy is similar to the one adopted above for slightly non-homogeneous fluids, see in particular Subsections 4.2 and 4.3.

First of all, let us write down the system of wave equations, which governs the propagation of oscillations in the dynamics. We could call them *Rossby waves*: this terminology is typically used when there are variations of the rotation axis (see e.g. [10]), but in our context having a non-constant ρ_0 produces very similar effects.

5.2.1 Description of Rossby waves and regularization

With the notation introduced in (26), we can write system (2) in the form of a wave equation (28), which we recall here for convenience:

$$(44) \quad \begin{cases} \varepsilon \partial_t \sigma_{\varepsilon} + \operatorname{div} V_{\varepsilon} = 0 \\ \varepsilon \partial_t V_{\varepsilon} + \nabla \Pi_{\varepsilon} + V_{\varepsilon}^{\perp} = \varepsilon f_{\varepsilon}. \end{cases}$$

Recall also the uniform bounds (27) for $(f_{\varepsilon})_{\varepsilon}$. Applying the curl operator to the second equation, we find

$$(45) \quad \begin{cases} \varepsilon \partial_t \sigma_{\varepsilon} + \operatorname{div} V_{\varepsilon} = 0 \\ \varepsilon \partial_t \eta_{\varepsilon} + \operatorname{div} V_{\varepsilon} = \varepsilon \operatorname{curl} f_{\varepsilon}, \end{cases}$$

where $\eta_\varepsilon = \operatorname{curl} V_\varepsilon$ as in Subsection 3.3 above.

As in Subsection 4.2, we now fix $M \in \mathbb{N}$ and, denoting by S_M the low frequency cut-off operator of a Littlewood-Paley decomposition introduced in (62) in the Appendix, we define

$$\sigma_{\varepsilon,M} := S_M \sigma_\varepsilon, \quad V_{\varepsilon,M} := S_M V_\varepsilon \quad \text{and} \quad \eta_{\varepsilon,M} := \operatorname{curl} V_{\varepsilon,M} = S_M \eta_\varepsilon.$$

Notice that all these quantities are smooth with respect to the x variable. For later use, let us immediately introduce also the smooth functions

$$u_{\varepsilon,M} := S_M u_\varepsilon \quad \text{and} \quad \omega_{\varepsilon,M} := \operatorname{curl} u_{\varepsilon,M} = S_M \omega_\varepsilon.$$

Let us point out that, analogously to (33), by uniform bounds we get

$$\|u_{\varepsilon,M}\|_{L_T^2(H^k)} \leq C(T, k, M)$$

for all $k \geq 1$, for some constant $C(T, k, M)$ just depending on the quantities in the brackets; in particular, for $k = 1$ we have

$$(46) \quad \|u_{\varepsilon,M}\|_{L_T^2(H^1)} \leq C(T).$$

The following regularization result holds true. Its proof is standard, and hence omitted. We limit ourselves to noticing here that the convergence properties (47) are straightforward consequences of the uniform bound (30) for $(\sigma_\varepsilon)_\varepsilon$ and the embedding $(\eta_\varepsilon)_\varepsilon \subset L_T^\infty(H^{-1})$.

Proposition 5.1. *For any fixed time $T > 0$, the following convergence properties hold, in the limit $M \rightarrow +\infty$:*

$$(47) \quad \begin{cases} \sup_{\varepsilon > 0} \|\sigma_\varepsilon - \sigma_{\varepsilon,M}\|_{L_T^\infty(H^{-s})} \rightarrow 0 & \forall s > 2 \\ \sup_{\varepsilon > 0} \|\eta_\varepsilon - \eta_{\varepsilon,M}\|_{L_T^\infty(H^{-s})} \rightarrow 0 & \forall s > 1. \end{cases}$$

Moreover, for any $M > 0$, the couple $(\sigma_{\varepsilon,M}, \eta_{\varepsilon,M})$ satisfies the wave equation

$$(48) \quad \begin{cases} \varepsilon \partial_t \sigma_{\varepsilon,M} + \operatorname{div} V_{\varepsilon,M} = 0 \\ \varepsilon \partial_t \eta_{\varepsilon,M} + \operatorname{div} V_{\varepsilon,M} = \varepsilon \operatorname{curl} f_{\varepsilon,M}, \end{cases}$$

where $(f_{\varepsilon,M})_\varepsilon$ is a family of smooth functions satisfying

$$(49) \quad \sup_{\varepsilon > 0} \|f_{\varepsilon,M}\|_{L_T^2(H^s)} \leq C(s, M, T) \quad \forall s \geq 0,$$

for suitable constants $C(s, M, T)$ depending only on the fixed $s \geq 0$, $M > 0$ and $T > 0$.

Let us conclude this part by proving an important statement, which enables us to compare the two velocity fields $V_{\varepsilon,M}$ and $u_{\varepsilon,M}$.

Proposition 5.2. *For all $M \in \mathbb{N}$ and all $\varepsilon \in]0, 1]$, the following equality holds true:*

$$V_{\varepsilon,M} = \rho_0 u_{\varepsilon,M} + \varepsilon^\theta \zeta_{\varepsilon,M} + h_{\varepsilon,M},$$

where $0 < \theta < 1$ is the exponent fixed in Proposition 3.9. Moreover, the families $(\zeta_{\varepsilon,M})_\varepsilon$ and $(h_{\varepsilon,M})_\varepsilon$ satisfy the uniform bounds

$$\begin{cases} \sup_{\varepsilon > 0} \|\zeta_{\varepsilon,M}\|_{L_T^2(H^s)} \leq C(s, M, T) & \forall s \geq 0 \\ \sup_{\varepsilon > 0} \|h_{\varepsilon,M}\|_{L_T^2(H^1)} \leq C_0(T) 2^{-M}, \end{cases}$$

for any time $T > 0$, where the constants $C(s, M)$ and C_0 depend only on norms of the initial data, on time T and, respectively, on the fixed s and M , and on the norm $\|\rho_0\|_{W^{2,\infty}}$.

Proof. By definition, we can write

$$(50) \quad V_{\varepsilon, M} = S_M(\rho_\varepsilon u_\varepsilon) = \varepsilon^\theta S_M(\varepsilon^{-\theta} s_\varepsilon u_\varepsilon) + S_M(\rho_0 u_\varepsilon),$$

where we recall that $s_\varepsilon = \rho_\varepsilon - \rho_0$ expresses oscillations of the density with respect to the target profile. Thanks to Proposition 3.9, we deduce that, for any fixed $M \in \mathbb{N}$, the family

$$\zeta_{\varepsilon, M} := \frac{1}{\varepsilon^\theta} S_M(s_\varepsilon u_\varepsilon)$$

is bounded in $L_T^2(H^s)$, uniformly with respect to $\varepsilon \in]0, 1]$, for any $s \geq 0$.

On the other hand, denoting by $[\mathcal{P}, \mathcal{Q}]$ the commutator between two operators \mathcal{P} and \mathcal{Q} , the following relation holds true:

$$S_M(\rho_0 u_\varepsilon) = \rho_0 u_{\varepsilon, M} + [S_M, \rho_0] u_\varepsilon.$$

So, let us set $h_{\varepsilon, M} := [S_M, \rho_0] u_\varepsilon$ and estimate its H^1 norm. First of all, by Lemma A.7 and uniform bounds (20), we immediately gather

$$\sup_{\varepsilon > 0} \|h_{\varepsilon, M}\|_{L_T^2(L^2)} \leq C 2^{-M} \|\nabla \rho_0\|_{L^\infty} \|u_\varepsilon\|_{L_T^2(L^2)} \leq C' 2^{-M}.$$

Furthermore, from the equality $\partial_j h_{\varepsilon, M} = [S_M, \rho_0] \partial_j u_\varepsilon + [S_M, \partial_j \rho_0] u_\varepsilon$, the combination of Lemma A.7 with (19) and (20) implies

$$\sup_{\varepsilon > 0} \|\partial_j h_{\varepsilon, M}\|_{L_T^2(L^2)} \leq C 2^{-M} \left(\|\nabla \rho_0\|_{L^\infty} \|\nabla u_\varepsilon\|_{L_T^2(L^2)} + \|\nabla^2 \rho_0\|_{L^\infty} \|u_\varepsilon\|_{L_T^2(L^2)} \right) \leq C 2^{-M}.$$

The proposition is now completely proved. \square

From Proposition 5.2 we immediately infer the next statement. Notice that we propose two (related but different) decompositions for the approximate vorticities $\eta_{\varepsilon, M}$: they will both be useful in our analysis.

Corollary 5.3. *For all $M \in \mathbb{N}$ and all $\varepsilon \in]0, 1]$, the following equalities hold true:*

$$\begin{cases} \eta_{\varepsilon, M} &= \eta_{\varepsilon, M}^{(1)} + \varepsilon^\theta \eta_{\varepsilon, M}^{(2)} \\ &= \rho_0 \omega_{\varepsilon, M} + u_{\varepsilon, M} \cdot \nabla^\perp \rho_0 + \varepsilon^\theta \operatorname{curl} \zeta_{\varepsilon, M} + \operatorname{curl} h_{\varepsilon, M} \\ \operatorname{div} V_{\varepsilon, M} &= u_{\varepsilon, M} \cdot \nabla \rho_0 + \varepsilon^\theta \operatorname{div} \zeta_{\varepsilon, M} + \operatorname{div} h_{\varepsilon, M}, \end{cases}$$

where the families $(\eta_{\varepsilon, M}^{(1)})_\varepsilon$ and $(\eta_{\varepsilon, M}^{(2)})_\varepsilon$ satisfy the following estimates:

$$\sup_{M \in \mathbb{N}} \sup_{\varepsilon > 0} \|\eta_{\varepsilon, M}^{(1)}\|_{L_T^2(L^2)} \leq C \quad \text{and} \quad \sup_{\varepsilon > 0} \|\eta_{\varepsilon, M}^{(2)}\|_{L_T^2(H^s)} \leq C(s, M),$$

for any given $s \geq 0$ fixed and where $(\zeta_{\varepsilon, M})_\varepsilon$ and $(h_{\varepsilon, M})_\varepsilon$ are defined in Proposition 5.2 and, in particular, satisfy the same bounds established there.

Proof. The previous statement is a straightforward consequence of Proposition 5.2, and hence omitted. We just notice here that the former decomposition for $\eta_{\varepsilon, M}$ comes directly from relation (50), by setting

$$\eta_{\varepsilon, M}^{(1)} := \operatorname{curl} S_M(\rho_0 u_\varepsilon) \quad \text{and} \quad \eta_{\varepsilon, M}^{(2)} := \operatorname{curl} S_M(\varepsilon^{-\theta} s_\varepsilon u_\varepsilon).$$

The uniform bounds for these quantities then derive from (19), (20) and Proposition 3.9. \square

5.2.2 Approximation and convergence

In this paragraph we state and prove some approximation results, in the same spirit as the ones given in Subsection 4.2 above, and we take the limit in the convective term by compensated compactness arguments.

Let us start by establishing the counterpart of Lemma 4.1 in the case of a non-constant density profile ρ_0 .

Lemma 5.4. *For any test function $\psi \in \mathcal{D}([0, T[\times \Omega)$, one has*

$$\lim_{\varepsilon \rightarrow 0^+} \left| \int_0^T \int_{\Omega} \rho_{\varepsilon} u_{\varepsilon} \otimes u_{\varepsilon} : \nabla \psi \, dx \, dt - \int_0^T \int_{\Omega} \rho_0 u_{\varepsilon} \otimes u_{\varepsilon} : \nabla \psi \, dx \, dt \right| = 0.$$

Proof. We decompose the density, according to our *ansatz*, as $\rho_{\varepsilon} = \rho_0 + s_{\varepsilon}$. Recall that, by (21) (keep in mind also (25) and Proposition 3.9), we know that, up to passing to subsequences, $(s_{\varepsilon})_{\varepsilon}$ strongly converges to 0 in e.g. $L_T^{\infty}(H_{\text{loc}}^{-\gamma})$ for any $0 < \gamma < 1$.

On the other hand, uniform bounds for $(u_{\varepsilon})_{\varepsilon}$ in $L_T^2(H^1)$ and a paraproduct decomposition immediately imply that $u_{\varepsilon} \otimes u_{\varepsilon}$ is uniformly bounded in $L_T^1(H^{1-\delta})$, for all $\delta > 0$. Here, we have used Corollary A.6 (v) in the Appendix, together with the continuous embedding $H^1 \hookrightarrow B_{\infty, \infty}^0$.

Therefore, by a direct application of Corollary A.6 (i) in the Appendix, we gather that the quantity $(\rho_{\varepsilon} u_{\varepsilon} \otimes u_{\varepsilon})_{\varepsilon}$ weakly converges to 0 (up to a suitable extraction of a subsequence) in the space $L_T^1(H^{-\gamma-\delta})$, for any $\delta > 0$ arbitrarily small. \square

The next step consists in regularizing the velocity fields in the convection term: thanks to the smoothness of ρ_0 , we can establish an analogue of Lemma 4.2, whose proof is exactly the same so is omitted.

Lemma 5.5. *For any test function $\psi \in \mathcal{D}([0, T[\times \Omega)$, one has*

$$\lim_{M \rightarrow +\infty} \limsup_{\varepsilon \rightarrow 0^+} \left| \int_0^T \int_{\Omega} \rho_0 u_{\varepsilon} \otimes u_{\varepsilon} : \nabla \psi \, dx \, dt - \int_0^T \int_{\Omega} \rho_0 u_{\varepsilon, M} \otimes u_{\varepsilon, M} : \nabla \psi \, dx \, dt \right| = 0.$$

Thanks to the previous statements, we just have to study, for any fixed $M \in \mathbb{N}$, the convergence for $\varepsilon \rightarrow 0$ of the integral

$$- \int_0^T \int_{\Omega} \rho_0 u_{\varepsilon, M} \otimes u_{\varepsilon, M} : \nabla \psi \, dx \, dt = \int_0^T \int_{\Omega} \operatorname{div}(\rho_0 u_{\varepsilon, M} \otimes u_{\varepsilon, M}) \cdot \psi \, dx \, dt,$$

where, as previously, we have integrated by parts since now we have enough smoothness in the space variable. We proceed in several steps.

Step 1. By direct differentiation, we obtain (because $\operatorname{div} u_{\varepsilon, M} = 0$)

$$\begin{aligned} (51) \quad \operatorname{div}(\rho_0 u_{\varepsilon, M} \otimes u_{\varepsilon, M}) &= \rho_0 u_{\varepsilon, M} \cdot \nabla u_{\varepsilon, M} + u_{\varepsilon, M} \cdot \nabla \rho_0 u_{\varepsilon, M} \\ &= \frac{1}{2} \rho_0 \nabla |u_{\varepsilon, M}|^2 + \rho_0 \omega_{\varepsilon, M} u_{\varepsilon, M}^{\perp} + u_{\varepsilon, M} \cdot \nabla \rho_0 u_{\varepsilon, M}. \end{aligned}$$

We remark that the first term in the right-hand side of the last equality gives rise to a contribution of the type $\rho_0 \nabla \Gamma$ in the limit. Analogously, the same fact happens for any term of the form $\Lambda_{\varepsilon, M} \nabla \rho_0$ where $\Lambda_{\varepsilon, M}$ is any smooth enough distribution: indeed, if we take a test vector field ψ of zero divergence and we integrate by parts, we get

$$\int_0^T \int_{\Omega} \Lambda_{\varepsilon, M} \nabla \rho_0 \cdot \psi \, dx \, dt = \int_0^T \int_{\Omega} \Lambda_{\varepsilon, M} \operatorname{div}(\rho_0 \psi) \, dx \, dt = - \int_0^T \int_{\Omega} \rho_0 \nabla \Lambda_{\varepsilon, M} \cdot \psi \, dx \, dt.$$

For notational convenience, from now on we will generically denote by $\Gamma_{\varepsilon,M}$ any vector field which can be written under the form

$$(52) \quad \Gamma_{\varepsilon,M} = \rho_0 \nabla \Lambda_{\varepsilon,M}^{(1)} + \Lambda_{\varepsilon,M}^{(2)} \nabla \rho_0,$$

with

$$(53) \quad \lim_{M \rightarrow +\infty} \lim_{\varepsilon \rightarrow 0^+} \int_0^T \int_{\Omega} \Gamma_{\varepsilon,M} \cdot \psi \, dx \, dt = \langle \rho_0 \nabla \Gamma, \psi \rangle_{\mathcal{D}' \times \mathcal{D}}$$

for all $\psi \in \mathcal{D}([0, T[\times \Omega)$ such that $\operatorname{div} \psi = 0$, and where Γ is a distribution over $[0, T[\times \Omega$.

On the other hand, we will generically denote by $\mathcal{R}_{\varepsilon,M}$ any remainder, i.e. any term satisfying the property

$$(54) \quad \lim_{M \rightarrow +\infty} \limsup_{\varepsilon \rightarrow 0^+} \left| \int_0^T \int_{\Omega} \mathcal{R}_{\varepsilon,M} \cdot \psi \, dx \, dt \right| = 0$$

for all divergence free test functions $\psi \in \mathcal{D}([0, T[\times \Omega)$.

With these notations, relation (51) can be written under the form

$$\operatorname{div} (\rho_0 u_{\varepsilon,M} \otimes u_{\varepsilon,M}) = \Gamma_{\varepsilon,M} + \rho_0 \omega_{\varepsilon,M} u_{\varepsilon,M}^{\perp} + u_{\varepsilon,M} \cdot \nabla \rho_0 u_{\varepsilon,M}.$$

Step 2. Let us focus on the vorticity term: by use of Corollary 5.3 and especially of the second decomposition for $\eta_{\varepsilon,M}$, we can write

$$\rho_0 \omega_{\varepsilon,M} u_{\varepsilon,M}^{\perp} = \eta_{\varepsilon,M} u_{\varepsilon,M}^{\perp} - u_{\varepsilon,M} \cdot \nabla^{\perp} \rho_0 u_{\varepsilon,M}^{\perp} - \varepsilon^{\theta} \operatorname{curl} \zeta_{\varepsilon,M} u_{\varepsilon,M}^{\perp} - \operatorname{curl} h_{\varepsilon,M} u_{\varepsilon,M}^{\perp},$$

where, in view of Proposition 5.2 and of estimates (46), the last two terms contribute as remainders, in the sense of relation (54). Plugging this expression into the equation for the convective term, we arrive at

$$(55) \quad \operatorname{div} (\rho_0 u_{\varepsilon,M} \otimes u_{\varepsilon,M}) = \Gamma_{\varepsilon,M} + \mathcal{R}_{\varepsilon,M} + \eta_{\varepsilon,M} u_{\varepsilon,M}^{\perp} - u_{\varepsilon,M} \cdot \nabla^{\perp} \rho_0 u_{\varepsilon,M}^{\perp} + u_{\varepsilon,M} \cdot \nabla \rho_0 u_{\varepsilon,M}.$$

Step 3. Let us concentrate now on the term

$$(56) \quad X_{\varepsilon,M} := - \left(u_{\varepsilon,M} \cdot \nabla^{\perp} \rho_0 \right) u_{\varepsilon,M}^{\perp} + (u_{\varepsilon,M} \cdot \nabla \rho_0) u_{\varepsilon,M}.$$

For reasons which will appear clear in a while (see also [24]), let us introduce a function $b \in \mathcal{C}_0^{\infty}(\mathbb{R}^2)$, with $0 \leq b(y) \leq 1$, such that $b \equiv 1$ on $\{|y| \leq 1\}$ and $b \equiv 0$ on $\{|y| \geq 2\}$. For any $x \in \Omega$, we define

$$(57) \quad b_M(x) := b(2^{M/2} \nabla \rho_0(x)).$$

To begin with, we notice that, once a $2 < q < +\infty$ is fixed, for any compact set $K \subset \Omega$ we can estimate

$$\|b_M X_{\varepsilon,M}\|_{L^1([0,T] \times K)} \leq C \|X_{\varepsilon,M}\|_{L_T^1(L^{q/2})} \left(\mu \left\{ x \in \mathbb{R}^2 \mid |\nabla \rho_0(x)| \leq 2^{1-M/2} \right\} \right)^{(q-2)/q}.$$

Since we can bound $\|X_{\varepsilon,M}\|_{L_T^1(L^{q/2})}$ by the quantity $C \|\nabla \rho_0\|_{L^{\infty}} \|u_{\varepsilon,M}\|_{L_T^2(L^q)}^2$, which is controlled by a uniform constant in view of (19) and Sobolev embeddings, assumption (12) tells us that the term $b_M X_{\varepsilon,M}$ is a remainder in the sense of relation (54).

On the other hand, on the support of $1 - b_M$, the vector $\nabla \rho_0$ is far from 0, so that we can decompose $u_{\varepsilon,M}$ along the basis of \mathbb{R}^2 (conveniently renormalized) $\{\nabla \rho_0, \nabla^{\perp} \rho_0\}$. More precisely, we can write

$$(1 - b_M) u_{\varepsilon,M} = (1 - b_M) \frac{1}{|\nabla \rho_0|^2} \left((u_{\varepsilon,M} \cdot \nabla \rho_0) \nabla \rho_0 + (u_{\varepsilon,M} \cdot \nabla^{\perp} \rho_0) \nabla^{\perp} \rho_0 \right)$$

$$(1 - b_M) u_{\varepsilon, M}^\perp = (1 - b_M) \frac{1}{|\nabla \rho_0|^2} \left((u_{\varepsilon, M}^\perp \cdot \nabla \rho_0) \nabla \rho_0 + (u_{\varepsilon, M} \cdot \nabla \rho_0) \nabla^\perp \rho_0 \right).$$

Notice that, in view of (52), we can write the terms in $\nabla \rho_0$ in the generic form $\Gamma_{\varepsilon, M}$. Therefore, in the end, by definition (56) of $X_{\varepsilon, M}$ we get

$$\begin{aligned} (1 - b_M) X_{\varepsilon, M} &= \Gamma_{\varepsilon, M} - (1 - b_M) \frac{1}{|\nabla \rho_0|^2} (u_{\varepsilon, M} \cdot \nabla^\perp \rho_0) (u_{\varepsilon, M} \cdot \nabla \rho_0) \nabla^\perp \rho_0 + \\ &\quad + (1 - b_M) \frac{1}{|\nabla \rho_0|^2} (u_{\varepsilon, M} \cdot \nabla \rho_0) (u_{\varepsilon, M} \cdot \nabla^\perp \rho_0) \nabla^\perp \rho_0 \\ &= \Gamma_{\varepsilon, M}, \end{aligned}$$

due to the very special cancellations of the two terms in $\nabla^\perp \rho_0$.

Finally, we obtain that $X_{\varepsilon, M}$ contributes as a term of the form (52). Hence, in view of (55), we have

$$\operatorname{div} (\rho_0 u_{\varepsilon, M} \otimes u_{\varepsilon, M}) = \Gamma_{\varepsilon, M} + \mathcal{R}_{\varepsilon, M} + \eta_{\varepsilon, M} u_{\varepsilon, M}^\perp.$$

Step 4. In order to treat the last term in the right-hand side of the previous equation, we resort again to the function b_M defined in (57) above. We also exploit the first decomposition of $\eta_{\varepsilon, M}$ given in Corollary 5.3.

Thanks to the properties stated in Corollary 5.3, for any $T > 0$ and any compact set $K \subset \Omega$, it is an easy matter to estimate

$$\begin{aligned} \left\| b_M \eta_{\varepsilon, M} u_{\varepsilon, M}^\perp \right\|_{L^1([0, T] \times K)} &\leq C \varepsilon^\theta \left\| \eta_{\varepsilon, M}^{(2)} \right\|_{L_T^2(L^2)} \|u_{\varepsilon, M}\|_{L_T^2(L^2)} + \\ &\quad + C \left\| \eta_{\varepsilon, M}^{(1)} \right\|_{L_T^2(L^2)} \|u_{\varepsilon, M}\|_{L_T^2(L^q)} \left(\mu \left\{ x \in \mathbb{R}^2 \mid |\nabla_h \rho_0(x)| \leq 2^{1-M/2} \right\} \right)^{1/m} \\ &\leq C(M, T) \varepsilon^\theta + C \left(\mu \left\{ x \in \mathbb{R}^2 \mid |\nabla_h \rho_0(x)| \leq 2^{1-M/2} \right\} \right)^{1/m}, \end{aligned}$$

where q and m , both larger than 2, are linked by the relation $1/q + 1/m = 1/2$. Therefore, in view of assumption (12), also this term is a remainder, in the sense that it gives a contribution of the form (54).

For the other term $(1 - b_M) \eta_{\varepsilon, M} u_{\varepsilon, M}^\perp$, we decompose the velocity field along $\{\nabla \rho_0, \nabla^\perp \rho_0\}$, as done before. Forgetting about the term along $\nabla \rho_0$, since is a contribution of the form $\Gamma_{\varepsilon, M}$ (recall (52) above), we can write

$$\begin{aligned} (1 - b_M) \eta_{\varepsilon, M} u_{\varepsilon, M}^\perp &= \Gamma_{\varepsilon, M} + (1 - b_M) \frac{1}{|\nabla \rho_0|^2} \eta_{\varepsilon, M} (u_{\varepsilon, M} \cdot \nabla \rho_0) \nabla^\perp \rho_0 \\ &= \Gamma_{\varepsilon, M} + \frac{1 - b_M}{|\nabla \rho_0|^2} \eta_{\varepsilon, M} \left(\operatorname{div} V_{\varepsilon, M} - \varepsilon^\theta \operatorname{div} \zeta_{\varepsilon, M} - \operatorname{div} h_{\varepsilon, M} \right) \nabla^\perp \rho_0, \end{aligned}$$

where we have used also Corollary 5.3. Notice that the terms presenting $\varepsilon^\theta \operatorname{div} \zeta_{\varepsilon, M}$ and $\operatorname{div} h_{\varepsilon, M}$ give rise to remainders, in the sense of (54). For the former item, this fact is obvious thanks to the presence of ε^θ . As for the latter, it is enough to decompose again $\eta_{\varepsilon, M} = \eta_{\varepsilon, M}^{(1)} + \varepsilon^\theta \eta_{\varepsilon, M}^{(2)}$ and to use the bounds for $h_{\varepsilon, M}$ given in Proposition 5.2.

Putting all these facts together, we discover that the convective term can be written as

$$\operatorname{div} (\rho_0 u_{\varepsilon, M} \otimes u_{\varepsilon, M}) = \Gamma_{\varepsilon, M} + \mathcal{R}_{\varepsilon, M} + (1 - b_M) \frac{1}{|\nabla \rho_0|^2} \eta_{\varepsilon, M} \operatorname{div} V_{\varepsilon, M} \nabla^\perp \rho_0.$$

Step 5. As final step, we use the first relation in (48) to write

$$\begin{aligned}\eta_{\varepsilon,M} \operatorname{div} V_{\varepsilon,M} &= -\varepsilon \eta_{\varepsilon,M} \partial_t \sigma_{\varepsilon,M} = -\varepsilon (\eta_{\varepsilon,M} - \sigma_{\varepsilon,M}) \partial_t \sigma_{\varepsilon,M} - \frac{\varepsilon}{2} \frac{d}{dt} |\sigma_{\varepsilon,M}|^2 \\ &= -\varepsilon \frac{d}{dt} \left((\eta_{\varepsilon,M} - \sigma_{\varepsilon,M}) \sigma_{\varepsilon,M} + |\sigma_{\varepsilon,M}|^2 / 2 \right) + \varepsilon \operatorname{curl} f_{\varepsilon,M} \sigma_{\varepsilon,M},\end{aligned}$$

where we also exploited the difference of the two equations in (48), recall (29).

In the end, in light also of (49) and uniform bounds, the previous relations prove that

$$(1 - b_M) \frac{1}{|\nabla \rho_0|^2} \eta_{\varepsilon,M} \operatorname{div} V_{\varepsilon,M} \nabla^\perp \rho_0 = \mathcal{R}_{\varepsilon,M},$$

which finally implies that

$$(58) \quad \operatorname{div} (\rho_0 u_{\varepsilon,M} \otimes u_{\varepsilon,M}) = \Gamma_{\varepsilon,M} + \mathcal{R}_{\varepsilon,M}.$$

Remark 5.6. We notice here that, *in absence of vacuum*, we could have alternatively worked on a different approximation of the convective term, namely on

$$\operatorname{div} \left(\frac{1}{\rho_0} V_{\varepsilon,M} \otimes V_{\varepsilon,M} \right).$$

An analogous compensated compactness argument (see also [19]) would have led us to the same conclusion: namely, the transport term vanishes in the limit, up to a remainder of the form $\rho_0 \nabla \Gamma$, due to the strong constraint which is imposed by having a variable target density.

The advantage of our approach is that here we do not need the bound $\rho_0 \geq \rho_* > 0$.

Remark 5.7. Notice that the terms contributing to $\Gamma_{\varepsilon,M}$ are either quadratic in $u_{\varepsilon,M}$ (see Steps 1 and 3), or they depend on the product $\eta_{\varepsilon,M} u_{\varepsilon,M}$ (see Step 4). In view of Corollary 5.3, we conclude that these terms are uniformly bounded, both with respect to ε and M , in $L_T^1(L^1)$, up to a remainder of order $O(\varepsilon^\theta)$. So we can assume that the distribution Γ in (53) is a Radon measure over $[0, +\infty[\times \Omega$ and that the convergence holds true in the weak-* topology of the space $\mathcal{M}([0, T] \times \Omega)$ of Radon measures over $[0, T] \times \Omega$ (see Comments to Chapter 4 in [7]).

5.3 Conclusion: passing to the limit

We are now in position of passing to the limit in the modified weak formulation (43), completing in this way the proof to Theorem 2.7.

We have already seen that we can pass to the limit with no difficulty in the terms V_ε , σ_ε and in the viscosity term, as well as in the initial data. On the other hand, putting together Lemmas 5.4 and 5.5, relation (58) and Remark 5.7, we gather the convergence

$$(59) \quad \int_0^T \int_\Omega \rho_\varepsilon u_\varepsilon \otimes u_\varepsilon : \nabla \psi \, dx \, dt \longrightarrow -\langle \Gamma, \operatorname{div} (\rho_0 \psi) \rangle_{\mathcal{M} \times \mathcal{C}} = \langle \rho_0 \nabla \Gamma, \psi \rangle_{\mathcal{D}' \times \mathcal{D}}$$

when $\varepsilon \rightarrow 0$, for all test functions $\psi \in \mathcal{D}([0, T[\times \Omega)$ such that $\operatorname{div} \psi = 0$. In the previous formula, the notation $\langle \cdot, \cdot \rangle_{\mathcal{M} \times \mathcal{C}}$ denotes the duality product between Radon measures and continuous functions.

Therefore, starting from (43), where we have set $\psi = \nabla^\perp \phi$, in the limit for $\varepsilon \rightarrow 0$ we find

$$\begin{aligned}- \int_0^T \int_\Omega \rho_0 u \cdot \partial_t \nabla^\perp \phi - \int_0^T \int_\Omega \sigma \partial_t \phi - \\ - \int_0^T \int_\Omega \Gamma \operatorname{div} (\rho_0 \nabla^\perp \phi) + \nu \int_0^T \int_\Omega \nabla u : \nabla \nabla^\perp \phi = \int_\Omega m_0 \cdot \nabla^\perp \phi(0) + \int_\Omega r_0 \phi(0),\end{aligned}$$

where $\phi \in \mathcal{D}([0, T[\times \Omega)$ is any test function, with no additional restrictions, and where with a little abuse of notation, we have identified

$$- \int_0^T \int_{\Omega} \Gamma \operatorname{div} (\rho_0 \nabla^\perp \phi) := - \langle \Gamma, \operatorname{div} (\rho_0 \nabla^\perp \phi) \rangle_{\mathcal{M} \times \mathcal{C}}.$$

A further integration by parts allows us to arrive (after multiplying by -1) at the equation

$$\begin{aligned} & - \int_0^T \int_{\Omega} (\operatorname{curl} (\rho_0 u) - \sigma) \cdot \partial_t \phi \, dx \, dt + \\ & + \int_0^T \int_{\Omega} \Gamma \operatorname{div} (\rho_0 \nabla^\perp \phi) \, dx \, dt + \nu \int_0^T \int_{\Omega} \nabla \omega : \nabla \phi \, dx \, dt = \int_{\Omega} (\operatorname{curl} m_0 - r_0) \phi(0) \, dx, \end{aligned}$$

which coincides with the weak formulation of equation (13), in view of (59).

Theorem 2.7 is thus completely proved. \square

5.4 A conditional convergence result

In this subsection, we state and prove a convergence result for the fully non-homogeneous case, where we are able to pass to the limit to the full system, in which the dynamics of the density fluctuation function and the velocity field are decoupled.

This is just a conditional result, because very strong assumptions are required on the family of weak solutions: in particular, we need to assume uniform bounds in higher norms for the family of velocity fields (see in particular conditions (ii) and (iii) in Theorem 5.8 below), which cannot be deduced from classical energy estimates.

The statement is the following.

Theorem 5.8. *With the notation and under the assumptions of Theorem 2.7, assume moreover that $\rho_0 \in W^{3,\infty}(\Omega)$ and that the following conditions hold true:*

- (i) $(r_{0,\varepsilon})_\varepsilon \subset H^{1+\beta}(\Omega)$, for some $\beta \in]0, 1[$;
- (ii) $(u_\varepsilon)_\varepsilon \subset L_{\text{loc}}^\infty(\mathbb{R}_+; H^1(\Omega)) \cap L_{\text{loc}}^2(\mathbb{R}_+; H^2(\Omega))$;
- (iii) $(u_\varepsilon)_\varepsilon \subset \mathcal{C}_{\text{loc}}^{0,\gamma}(\mathbb{R}_+; L^2(\Omega))$, for some $\gamma \in]0, 1[$.

Then there exist distributions Π , Γ_0 and Γ_1 over $\mathbb{R}_+ \times \Omega$ such that the limit points σ and u satisfy the system

$$(60) \quad \begin{cases} \partial_t \sigma + u \cdot \nabla \sigma = \operatorname{curl} (\rho_0 \nabla \Gamma_1) \\ \rho_0 \partial_t u + \nabla \Pi + \rho_0 \nabla \Gamma_0 + \sigma u^\perp - \nu \Delta u = 0 \\ \operatorname{div} u = \operatorname{div} (\rho_0 u) = 0, \end{cases}$$

with initial data respectively $\sigma|_{t=0} = r_0$ and $(\rho_0 u)|_{t=0} = m_0$, where r_0 and m_0 have been defined in (6).

It goes without saying that, under conditions (i)-(ii)-(iii), the convergence properties for $(u_\varepsilon)_\varepsilon$ and $(\sigma_\varepsilon)_\varepsilon$ stated in Theorem 5.8 can be improved (see also Proposition 5.13 below). However, our focus here is on obtaining convergence to the full system rather than (13). In this respect, let us make some comments.

Remark 5.9. In order to make sense of the equation on σ_ε and to pass to the limit, we need enough regularity on σ_ε in order to define the product $\sigma_\varepsilon u_\varepsilon$, and to prove uniform bounds. Now, the only way to get information for σ_ε is to exploit equation (29): assumptions (i) and (ii) of Theorem 5.8 are designed so as to get the desired smoothness for all terms appearing in that equation, while assumption (iii) is needed in order to prove compactness properties for $(\sigma_\varepsilon)_\varepsilon$.

Remark 5.10. It is well-known that, for 2-D density-dependent Navier-Stokes equations *in the absence of vacuum*, if the initial velocity field is in H^1 , then there exists a weak solution $u \in \mathcal{C}_T(H^1) \cap L_T^2(H^2)$. We refer e.g. to pages 31-32 of [29] and to comments to Theorem 3.41 in [4] for precise statements and further details. Moreover, an $H^{1+\beta}$ assumption on the initial density is required for recovering uniqueness; see e.g. Section 5 of [11].

Nonetheless, it seems to us not possible to get *uniform bounds* for $(u_\varepsilon)_\varepsilon$ in $L_T^\infty(H^1) \cap L_T^2(H^2)$, even in absence of vacuum, using the strategy of proof adopted in the above mentioned results. The problem is that the Coriolis term, which does not affect basic energy estimates, does affect these higher order type estimates. This is why we have to assume the *a priori* bounds (ii).

Remark 5.11. As it will be apparent from the proof, see in particular Paragraph 5.4.2 below, the regularities of the distributions Γ_0 and Γ_1 with respect to the space variable will be enough to make sense of the product of their gradients by ρ_0 in \mathcal{D}' .

The proof of Theorem 5.8 consists in two main steps. In the first one, in Paragraph 5.4.1, we study the density variations: the new assumptions allow to improve the regularity of the σ_ε 's. This turns out to be fundamental in passing to the limit $\varepsilon \rightarrow 0$, as shown in Paragraph 5.4.2

5.4.1 Study of the density variations

First of all, let us consider the functions $s_\varepsilon := \rho_\varepsilon - \rho_0$, which satisfy equation (23). We can establish the following result.

Lemma 5.12. *Under assumptions (i) and (ii) of Theorem 5.8, for all $0 < \alpha < \beta$, one has the uniform bound*

$$(s_\varepsilon)_\varepsilon \subset L_{\text{loc}}^\infty(\mathbb{R}_+; H^{1+\alpha}(\Omega)) \cap \mathcal{C}_{\text{loc}}^{0,1/2}(\mathbb{R}_+; H^\alpha(\Omega)).$$

In particular, this family is compact e.g. in the space $L_T^\infty(H_{\text{loc}}^\alpha)$ for all fixed times $T > 0$.

Proof. The starting point is the transport equation (23). By assumption, the family of initial data $(\varepsilon r_{0,\varepsilon})_\varepsilon$ is uniformly bounded in $H^{1+\beta}$, and the family of external forces $(u_\varepsilon \cdot \nabla \rho_0)_\varepsilon$ is uniformly bounded in $L_T^2(H^2) \hookrightarrow L_T^2(H^{1+\beta})$. Moreover, the family of divergence-free vector fields $(u_\varepsilon)_\varepsilon$ is uniformly bounded in $L_T^2(H^2)$. Therefore, for all $0 < \alpha < \beta$, by an application of Proposition 5.2 of [11] we deduce that $(s_\varepsilon)_\varepsilon \subset L_T^\infty(H^{1+\alpha})$ for all fixed $T > 0$.

On the other hand, we can write $\partial_t s_\varepsilon = -\text{div}(s_\varepsilon u_\varepsilon) - u_\varepsilon \cdot \nabla \rho_0$: because $H^{1+\alpha}$ is a Banach algebra, we infer the uniform bound $(\partial_t s_\varepsilon)_\varepsilon \subset L_T^2(H^\alpha)$ for all times $T > 0$, which implies that $(s_\varepsilon)_\varepsilon \subset \mathcal{C}_T^{0,1/2}(H^\alpha)$.

Finally, the last assertion of the statement follows by the Ascoli-Arzelà theorem and interpolation of the previous two properties. The lemma is proved. \square

Thanks to the previous result, we can improve the regularity for the functions σ_ε .

Proposition 5.13. *Under assumptions (i) and (ii) of Theorem 5.8, one has the uniform bound*

$$(\sigma_\varepsilon)_\varepsilon \subset L_{\text{loc}}^\infty(\mathbb{R}_+; H^{-1}(\Omega)).$$

In particular, there holds $(\sigma_\varepsilon u_\varepsilon)_\varepsilon \subset L_{\text{loc}}^2(\mathbb{R}_+; H^{-1}(\Omega))$.

Moreover, under assumption (iii) $(\sigma_\varepsilon)_\varepsilon$ is compact in $L_T^\infty(H_{\text{loc}}^{-1-k-\delta})$, for any $0 < \delta < 1$ and for all times $T > 0$, where k is the index fixed in Proposition 3.9. In particular, one gathers also the convergence property

$$\sigma_\varepsilon u_\varepsilon \rightharpoonup \sigma u \quad \text{in} \quad L_T^2(H_{\text{loc}}^{-1-k-\delta}).$$

Proof. Integrating equation (29) in time, for all $\varepsilon \in]0, 1[$ we can write

$$\sigma_\varepsilon(t, x) = \varepsilon r_{0,\varepsilon}(x) + \eta_\varepsilon(t, x) - \operatorname{curl}(m_{0,\varepsilon}(x)) + \int_0^t \operatorname{curl} f_\varepsilon(\tau, x) d\tau,$$

where we recall that $\eta_\varepsilon := \operatorname{curl}(\rho_\varepsilon u_\varepsilon)$ and f_ε is defined in (26).

By assumption on the initial data, we have that both $(r_{0,\varepsilon})_\varepsilon$ and $(\operatorname{curl} m_{0,\varepsilon})_\varepsilon$ are uniformly bounded in H^{-1} . Furthermore, writing $\rho_\varepsilon = \rho_0 + s_\varepsilon$, by Lemma 5.12 and product rules we deduce that $(\eta_\varepsilon)_\varepsilon \subset L_T^2(L^2)$.

For convenience, let us choose $\alpha = \beta/2$ in Lemma 5.12, and keep it fixed throughout this proof. We now consider the convection term coming into play in the definition of f_ε . We remark, by assumption (ii) of Theorem 5.8 and interpolation, that $(u_\varepsilon)_\varepsilon$ is uniformly bounded in $L_T^4(H^{3/2})$. Therefore, the family $\rho_0 u_\varepsilon \otimes u_\varepsilon$ is uniformly bounded in the space $L_T^1(H^2) \cap L_T^2(H^{3/2})$, while the family of $s_\varepsilon u_\varepsilon \otimes u_\varepsilon$ is uniformly bounded in $L_T^2(H^{1+\alpha})$, since we have chosen $\alpha < 1/2$. On the other hand, $(\Delta\omega_\varepsilon)_\varepsilon$ is uniformly bounded in $L_T^2(H^{-1})$, so we finally deduce the uniform bound $(\operatorname{curl} f_\varepsilon)_\varepsilon \subset L_T^2(H^{-1})$.

Putting all these properties together, we deduce that the family $(\sigma_\varepsilon)_\varepsilon$ is uniformly bounded in $L_T^\infty(H^{-1})$, for all $T > 0$. By product rules of Corollary A.6 (iv), the uniform bounds for $(\sigma_\varepsilon u_\varepsilon)_\varepsilon$ follow immediately.

Remark that, by (29) again and the above study on $(f_\varepsilon)_\varepsilon$, it follows also that $(\eta_\varepsilon - \sigma_\varepsilon)_\varepsilon$ is compact in the space $\mathcal{C}_T^{0,1/2-\delta}(H_{\text{loc}}^{-1-\delta})$, for all $0 < \delta < 1$. On the other hand, thanks to Proposition 3.9, we can write

$$\eta_\varepsilon = \operatorname{curl}(\rho_0 u_\varepsilon) + \varepsilon^\theta \operatorname{curl}(\varepsilon^{-\theta} s_\varepsilon u_\varepsilon).$$

Observe that $(u_\varepsilon)_\varepsilon \subset L_T^\infty(H^1)$, so we deduce from Proposition 3.9 that $(\varepsilon^{-\theta} s_\varepsilon u_\varepsilon)_\varepsilon$ is uniformly bounded in $L_T^\infty(H^{-k-\delta})$, for any $\delta > 0$ small (we could actually improve this property, in view of Lemma 5.12 above, but it is enough for our purposes). Therefore, the latter term in the right-hand side converges strongly, as $\varepsilon \rightarrow 0$, in the space $L_T^\infty(H^{-k-\delta-1})$ for all $\delta > 0$ arbitrarily small. Moreover, by assumption (iii) we gather that the former term is uniformly bounded in $\mathcal{C}_T^{0,\gamma}(H^{-1})$, and so it is compact (by the Ascoli-Arzelà theorem) in $\mathcal{C}_T^{0,\gamma-\delta}(H_{\text{loc}}^{-1-\delta})$, where again $\delta > 0$ is arbitrarily small. From the previous properties, we immediately deduce that $(\sigma_\varepsilon)_\varepsilon$ is compact in the space $L_T^\infty(H_{\text{loc}}^{-1-k-\delta})$, for any $T > 0$ fixed and any $\delta > 0$ small.

In particular, we can assume, with no loss of generality, that $1 + k + \delta < 2$. Hence, combining this property with product rules of Corollary A.6 (iv) and the previous uniform bounds, we discover that $(\sigma_\varepsilon u_\varepsilon)_\varepsilon$ is weakly convergent to the product σu in $L_T^2(H_{\text{loc}}^{-1-k-\delta})$. \square

5.4.2 Passing to the limit

In view of the previous results, it is possible to pass to the limit in system (2).

First of all, we notice that we are now allowed to write (still in the weak sense) the equation

$$(61) \quad \partial_t \sigma_\varepsilon + \operatorname{div}(\sigma_\varepsilon u_\varepsilon) = -\varepsilon^{-1} u_\varepsilon \cdot \nabla \rho_0.$$

As a consequence of this relation and Proposition 5.13, arguing exactly in the same way as in establishing (32), we find the uniform embedding property

$$(\varepsilon^{-1} u_\varepsilon \cdot \nabla \rho_0)_\varepsilon \subset W_T^{-1,\infty}(H^{-1}) + L_T^2(H^{-1}) \hookrightarrow W^{-1,\infty}([0, T]; H^{-1}(\Omega)),$$

and then this term has to weakly converge, in the previous space, to some $g \in W_T^{-1,\infty}(H^{-1})$. Observe that, to be consistent with Theorem 2.7, g has to be equal to $\operatorname{curl}(\rho_0 \nabla \Gamma_1)$, for some scalar distribution Γ_1 . Indeed, since u_ε is divergence-free, we can write $u_\varepsilon = \nabla^\perp \Psi_\varepsilon$, hence

$$u_\varepsilon \cdot \nabla \rho_0 = \nabla^\perp \Psi_\varepsilon \cdot \nabla \rho_0 = -\nabla \Psi_\varepsilon \cdot \nabla^\perp \rho_0 = -\operatorname{curl}(\rho_0 \nabla \Psi_\varepsilon).$$

Therefore, thanks to these computations and Proposition 5.13, we can pass to the limit in equation (61) and find

$$\partial_t \sigma + \operatorname{div}(\sigma u) = \operatorname{curl}(\rho_0 \nabla \Gamma_1),$$

for a suitable distribution Γ_1 such that $\operatorname{curl}(\rho_0 \nabla \Gamma_1) \in W_T^{-1, \infty}(H^{-1})$ for all $T > 0$ fixed.

Let us consider now the weak formulation of the momentum equation:

$$\int_0^T \int_{\Omega} \left(-\rho_{\varepsilon} u_{\varepsilon} \cdot \partial_t \psi - \rho_{\varepsilon} u_{\varepsilon} \otimes u_{\varepsilon} : \nabla \psi + \frac{1}{\varepsilon} \rho_{\varepsilon} u_{\varepsilon}^{\perp} \cdot \psi + \nu \nabla u_{\varepsilon} : \nabla \psi \right) dx dt = \int_{\Omega} m_{0, \varepsilon} \cdot \psi(0) dx,$$

where ψ is a smooth divergence-free test function, compactly supported in $[0, T[\times \Omega$. Obviously, the convergence of the ∂_t term, the viscosity term and the initial datum present no difficulty, and it can be performed as in the previous paragraphs. Furthermore, the computations made in Subsection 5.2 allow us to pass to the limit also in the convective term, as done in (59), for a suitable Radon measure $\Gamma \in \mathcal{M}([0, T] \times \Omega)$. So the only changes concern the convergence in the Coriolis term. Using the decomposition $\rho_{\varepsilon} = \rho_0 + \varepsilon \sigma_{\varepsilon}$ and the fact that $u_{\varepsilon} = \nabla^{\perp} \Psi_{\varepsilon}$, we can write

$$\frac{1}{\varepsilon} \int_0^T \int_{\Omega} \rho_{\varepsilon} u_{\varepsilon}^{\perp} \cdot \psi = \frac{1}{\varepsilon} \int_0^T \int_{\Omega} \rho_0 u_{\varepsilon}^{\perp} \cdot \psi + \int_0^T \int_{\Omega} \sigma_{\varepsilon} u_{\varepsilon}^{\perp} \cdot \psi = -\frac{1}{\varepsilon} \int_0^T \int_{\Omega} \rho_0 \nabla \Psi_{\varepsilon} \cdot \psi + \int_0^T \int_{\Omega} \sigma_{\varepsilon} u_{\varepsilon}^{\perp} \cdot \psi.$$

Passing to the limit in the latter term in the right-hand side can be done as in the mass equation above (using Proposition 5.13); moreover the former term is exactly the same term (using the fact that $\psi = \nabla^{\perp} \phi$, for some smooth ϕ) we have encountered in the right-hand side of the mass equation. In the end, we deduce that

$$\frac{1}{\varepsilon} \int_0^T \int_{\Omega} \rho_{\varepsilon} u_{\varepsilon}^{\perp} \cdot \psi dx dt \longrightarrow \langle \rho_0 \nabla \Gamma_1, \psi \rangle_{\mathcal{D}' \times \mathcal{D}} + \int_0^T \int_{\Omega} \sigma u^{\perp} \cdot \psi dx dt.$$

Therefore, after setting $\Gamma_0 = \Gamma + \Gamma_1$, we have proved that the momentum equation converges, in the weak sense, to the equation

$$\rho_0 \partial_t u + \nabla \Pi + \rho_0 \nabla \Gamma_0 + \sigma u^{\perp} - \nu \Delta u = 0,$$

supplemented with the constraints $\operatorname{div} u = \operatorname{div}(\rho_0 u) = 0$.

Theorem 5.8 is completely proved. \square

A Appendix – Fourier and harmonic analysis toolbox

We recall here the main ideas of Littlewood-Paley theory, which we exploited in the previous analysis. We refer e.g. to Chapter 2 of [4] for details. For simplicity of exposition, let us deal with the \mathbb{R}^d case; however, the whole construction can be adapted also to the d -dimensional torus \mathbb{T}^d .

First of all, let us introduce the so called ‘‘Littlewood-Paley decomposition’’, based on a non-homogeneous dyadic partition of unity with respect to the Fourier variable. We fix a smooth radial function χ supported in the ball $B(0, 2)$, equal to 1 in a neighborhood of $B(0, 1)$ and such that $r \mapsto \chi(re)$ is nonincreasing over \mathbb{R}_+ for all unitary vectors $e \in \mathbb{R}^d$. Set $\varphi(\xi) = \chi(\xi) - \chi(2\xi)$ and $\varphi_j(\xi) := \varphi(2^{-j}\xi)$ for all $j \geq 0$.

The dyadic blocks $(\Delta_j)_{j \in \mathbb{Z}}$ are defined by¹

$$\Delta_j := 0 \quad \text{if } j \leq -2, \quad \Delta_{-1} := \chi(D) \quad \text{and} \quad \Delta_j := \varphi(2^{-j}D) \quad \text{if } j \geq 0.$$

¹Throughout we agree that $f(D)$ stands for the pseudo-differential operator $u \mapsto \mathcal{F}^{-1}(f \mathcal{F}u)$.

We also introduce the following low frequency cut-off operator:

$$(62) \quad S_j u := \chi(2^{-j} D) = \sum_{k \leq j-1} \Delta_k \quad \text{for} \quad j \geq 0.$$

The following classical property holds true: for any $u \in \mathcal{S}'$, then one has the equality $u = \sum_j \Delta_j u$ in the sense of \mathcal{S}' . Let us also mention the so-called *Bernstein inequalities*, which explain the way derivatives act on spectrally localized functions.

Lemma A.1. *Let $0 < r < R$. A constant C exists so that, for any nonnegative integer k , any couple (p, q) in $[1, +\infty]^2$, with $p \leq q$, and any function $u \in L^p$, we have, for all $\lambda > 0$,*

$$\begin{aligned} \text{supp } \widehat{u} \subset B(0, \lambda R) &\implies \|\nabla^k u\|_{L^q} \leq C^{k+1} \lambda^{k+d(\frac{1}{p}-\frac{1}{q})} \|u\|_{L^p}; \\ \text{supp } \widehat{u} \subset \{\xi \in \mathbb{R}^d \mid r\lambda \leq |\xi| \leq R\lambda\} &\implies C^{-k-1} \lambda^k \|u\|_{L^p} \leq \|\nabla^k u\|_{L^p} \leq C^{k+1} \lambda^k \|u\|_{L^p}. \end{aligned}$$

By use of Littlewood-Paley decomposition, we can define the class of Besov spaces.

Definition A.2. Let $s \in \mathbb{R}$ and $1 \leq p, r \leq +\infty$. The *non-homogeneous Besov space* $B_{p,r}^s$ is defined as the subset of tempered distributions u for which

$$\|u\|_{B_{p,r}^s} := \left\| \left(2^{js} \|\Delta_j u\|_{L^p} \right)_{j \geq -1} \right\|_{\ell^r} < +\infty.$$

Besov spaces are interpolation spaces between Sobolev spaces. In fact, for any $k \in \mathbb{N}$ and $p \in [1, +\infty]$ we have the following chain of continuous embeddings:

$$B_{p,1}^k \hookrightarrow W^{k,p} \hookrightarrow B_{p,\infty}^k,$$

where $W^{k,p}$ denotes the classical Sobolev space of L^p functions with all the derivatives up to the order k in L^p . Moreover, for all $s \in \mathbb{R}$ we have the equivalence $B_{2,2}^s \equiv H^s$, with

$$(63) \quad \|f\|_{H^s} \sim \left(\sum_{j \geq -1} 2^{2js} \|\Delta_j f\|_{L^2}^2 \right)^{1/2}.$$

As an immediate consequence of the first Bernstein inequality, one gets the following embedding result.

Proposition A.3. *The space $B_{p_1,r_1}^{s_1}$ is continuously embedded in the space $B_{p_2,r_2}^{s_2}$ for all indices satisfying $p_1 \leq p_2$ and*

$$s_2 < s_1 - d \left(\frac{1}{p_1} - \frac{1}{p_2} \right) \quad \text{or} \quad s_2 = s_1 - d \left(\frac{1}{p_1} - \frac{1}{p_2} \right) \quad \text{and} \quad r_1 \leq r_2.$$

We recall also Lemma 2.73 of [4].

Lemma A.4. *If $1 \leq r < +\infty$, for any $f \in B_{p,r}^s$ one has*

$$\lim_{j \rightarrow +\infty} \|f - S_j f\|_{B_{p,r}^s} = 0.$$

Let us now introduce the paraproduct operator (after J.-M. Bony, see [5]). Constructing the paraproduct operator relies on the observation that, formally, any product of two tempered distributions u and v , may be decomposed into

$$(64) \quad uv = T_u v + T_v u + R(u, v),$$

where we have defined

$$T_u v := \sum_j S_{j-1} u \Delta_j v, \quad \text{and} \quad R(u, v) := \sum_j \sum_{|j'-j| \leq 1} \Delta_j u \Delta_{j'} v.$$

The above operator T is called ‘‘paraproduct’’ whereas R is called ‘‘remainder’’. The paraproduct and remainder operators have many nice continuity properties. The following ones have been of constant use in this paper (see the proof in e.g. Chapter 2 of [4]).

Proposition A.5. *For any $(s, p, r) \in \mathbb{R} \times [1, \infty]^2$ and $t > 0$, the paraproduct operator T maps continuously $L^\infty \times B_{p,r}^s$ in $B_{p,r}^s$ and $B_{\infty,\infty}^{-t} \times B_{p,r}^s$ in $B_{p,r}^{s-t}$. Moreover, the following estimates hold:*

$$\|T_u v\|_{B_{p,r}^s} \leq C \|u\|_{L^\infty} \|\nabla v\|_{B_{p,r}^{s-1}} \quad \text{and} \quad \|T_u v\|_{B_{p,r}^{s-t}} \leq C \|u\|_{B_{\infty,\infty}^{-t}} \|\nabla v\|_{B_{p,r}^{s-1}}.$$

For any (s_1, p_1, r_1) and (s_2, p_2, r_2) in $\mathbb{R} \times [1, \infty]^2$ such that $s_1 + s_2 > 0$, $1/p := 1/p_1 + 1/p_2 \leq 1$ and $1/r := 1/r_1 + 1/r_2 \leq 1$, the remainder operator R maps continuously $B_{p_1,r_1}^{s_1} \times B_{p_2,r_2}^{s_2}$ into $B_{p,r}^{s_1+s_2}$. In the case $s_1 + s_2 = 0$, provided $r = 1$, operator R is continuous from $B_{p_1,r_1}^{s_1} \times B_{p_2,r_2}^{s_2}$ with values in $B_{p,\infty}^0$.

As a corollary of the previous proposition, we deduce the following continuity properties of the product in Sobolev spaces, which have been used in the course of the analysis. In the statement, we limit ourselves to the case of space dimension $d = 2$, the only relevant one for this study.

Corollary A.6. *Let $d = 2$.*

- (i) *For all η and all δ in $]0, 1[$, such that $1 - \eta - \delta > 0$, the product is a continuous map from $H^{-\eta} \times H^{1-\delta}$ into $H^{-\eta-\delta}$.*
- (ii) *For all $0 \leq \eta < 1$, the product is a continuous map from $H^{-\eta} \times H^1$ into $H^{-\eta-\delta}$ for all $\delta > 0$ arbitrarily small.*
- (iii) *For all $0 < \eta \leq 1$, the product is a continuous map from $H^\eta \times H^1$ into L^2 .*
- (iv) *For all $0 < \eta < 2$, the product is a continuous map from $H^{-\eta} \times H^2$ into $H^{-\eta}$.*
- (v) *The product is a continuous map from $H^1 \times H^1$ into $H^{1-\delta}$ for all $\delta > 0$ arbitrarily small.*

Proof. To begin with, we show that the product of two tempered distributions maps continuously $H^{-\eta} \times H^{1-\delta}$ into $H^{-\eta-\delta}$. Indeed, let us take two tempered distributions $a \in H^{-\eta}$ and $w \in H^{1-\delta}$, and let us write

$$(65) \quad a w = T_a w + T_w a + R(a, w).$$

By a systematic use of Proposition A.5 and of embeddings $H^s \hookrightarrow B_{\infty,\infty}^{s-1}$, we deduce that

$$\|T_a w + T_w a\|_{H^{-\eta-\delta}} + \|R(a, w)\|_{B_{1,1}^{1-\eta-\delta}} \leq C \|a\|_{H^{-\eta}} \|w\|_{H^{1-\delta}}.$$

At this point, the continuous embedding $B_{1,1}^{1-\eta-\delta} \hookrightarrow H^{-\eta-\delta}$ completes the proof of our claim.

For the second point, we apply Bony’s paraproduct decomposition to get once again (65), with $a \in H^{-\eta}$ and $w \in H^1$. Keeping in mind the embedding $H^s \hookrightarrow B_{\infty,\infty}^{s-1}$, by application of Proposition A.5 we get, for all $\delta > 0$,

$$\|T_a w\|_{H^{-\eta}} + \|T_w a\|_{H^{-\eta-\delta}} \leq C \|a\|_{H^{-\eta}} \|w\|_{H^1}.$$

On the other hand, the remainder operator can be controlled again by Proposition A.5 in the following way:

$$\|R(a, w)\|_{B_{1,1}^{1-\eta}} \leq C \|a\|_{H^{-\eta}} \|w\|_{H^1},$$

and the continuous embedding $B_{1,1}^{1-\eta} \hookrightarrow H^{-\eta}$ completes the proof.

As for point (iii) of the statement, for $a \in H^\eta$ and $w \in H^1$, we write for $\kappa > 0$

$$\|T_a w\|_{L^2} + \|T_w a\|_{L^2} \leq \|a\|_{B_{\infty,\infty}^{-\kappa}} \|w\|_{H^\kappa} + \|a\|_{H^1} \|w\|_{B_{\infty,\infty}^{-1}},$$

so in particular taking $\eta = 1 - \kappa$

$$\|T_a w\|_{L^2} + \|T_w a\|_{L^2} \leq \|a\|_{H^\eta} \|w\|_{H^1}.$$

On the other hand

$$\|R(a, w)\|_{L^2} \leq \|a\|_{H^\eta} \|w\|_{H^1}.$$

The fourth item of the claim easily follows from analogous arguments, keeping in mind the embedding $H^s \hookrightarrow L^\infty$ for all $s > 1$ (in dimension $d = 2$). The last item is also a straightforward consequence of Proposition A.5, its proof is omitted. The corollary is now proved. \square

We recall also a classical commutator estimate (see e.g. Lemma 2.97 of [4]), which is needed in our analysis.

Lemma A.7. *Let $h \in C^1(\mathbb{R}^d)$ such that $(1 + |\cdot|) \widehat{h} \in L^1$. There exists a constant C such that, for any Lipschitz function $\ell \in W^{1,\infty}(\mathbb{R}^d)$ and any $f \in L^p(\mathbb{R}^d)$ and for all $\lambda > 0$, one has*

$$\| [h(\lambda^{-1}D), \ell] f \|_{L^p} \leq C \lambda^{-1} \|\nabla \ell\|_{L^\infty} \|f\|_{L^p}.$$

Going along the lines of the proof, it is easy to see that the constant C depends just on the L^1 norm of the function $|x| k(x)$, where $k = \mathcal{F}_\xi^{-1} h$ is the inverse Fourier transform of h .

To conclude, let us recall Gagliardo-Nirenberg inequalities, which we have repeatedly used in Subsection 4.4. We refer e.g. to Corollary 1.2 of [9] for their proof.

Proposition A.8. *Let $p \in [2, +\infty[$ such that $1/p > 1/2 - 1/d$. There exists a constant $C > 0$ such that, for any domain $\Omega \subset \mathbb{R}^d$ and for all $u \in H_0^1(\Omega)$, the following inequality holds true:*

$$\|u\|_{L^p(\Omega)} \leq C \|u\|_{L^2(\Omega)}^{1-\lambda} \|\nabla u\|_{L^2(\Omega)}^\lambda, \quad \text{with} \quad \lambda = \frac{d(p-2)}{2p}.$$

References

- [1] A. Babin, A. Mahalov, B. Nicolaenko: *Global splitting, integrability and regularity of 3D Euler and Navier-Stokes equations for uniformly rotating fluids*. European J. Mech. B Fluids **15**, 291–300 (1996)
- [2] A. Babin, A. Mahalov, B. Nicolaenko: *Regularity and integrability of 3D Euler and Navier-Stokes equations for rotating fluids*. Asymptot. Anal. **15**, 103–150 (1997)
- [3] A. Babin, A. Mahalov, B. Nicolaenko: *Global regularity of 3D rotating Navier-Stokes equations for resonant domains*. Indiana Univ. Math. J. **48**, 1133–1176 (1999)
- [4] H. Bahouri, J.-Y. Chemin and R. Danchin: *“Fourier Analysis and Nonlinear Partial Differential Equations”*. Grundlehren der Mathematischen Wissenschaften (Fundamental Principles of Mathematical Sciences), Springer, Heidelberg (2011)
- [5] J.-M. Bony: *Calcul symbolique et propagation des singularités pour les équations aux dérivées partielles non linéaires*. Ann. Sci. École Norm. Sup. **14**, 209–246 (1981)
- [6] D. Bresch, B. Desjardins: *Existence of global weak solution for a 2D viscous shallow water equation and convergence to the quasi-geostrophic model*. Comm. Math. Phys. **238**, 211–223 (2003)
- [7] H. Brezis: *“Functional analysis, Sobolev spaces and partial differential equations”*. Universitext, Springer, New York (2011)

- [8] J.-Y. Chemin, B. Desjardins, I. Gallagher, E. Grenier: *Anisotropy and dispersion in rotating fluids*. In “Nonlinear partial differential equations and their applications”, Collège de France Seminar, Stud. Math. Appl. **31**, North-Holland, Amsterdam (2002)
- [9] J.-Y. Chemin, B. Desjardins, I. Gallagher, E. Grenier: “*Mathematical geophysics. An introduction to rotating fluids and the Navier-Stokes equations*”. Oxford Lecture Series in Mathematics and its Applications, Oxford University Press, Oxford (2006)
- [10] B. Cushman-Roisin: “*Introduction to Geophysical Fluid Dynamics*”. Prentice-Hall, Englewood Cliffs (1994)
- [11] R. Danchin: *Local and global well-posedness results for flows of inhomogeneous viscous fluids*. Adv. Differential Equations **9**, 353–386 (2004)
- [12] B. Desjardins, E. Grenier: *On the homogeneous model of wind-driven ocean circulation*. SIAM J. Appl. Math., **60**, 43–60 (2000)
- [13] R. DiPerna, P.-L. Lions: *Ordinary differential equations, transport theory and Sobolev spaces*. Invent. Math., **98**, 511–547 (1989)
- [14] A. Dutrifoy: *Fast averaging for long- and short-wave scaled equatorial shallow water equations with Coriolis parameter deviating from linearity*. Arch. Ration. Mech. Anal., **216**, 261–312 (2015)
- [15] A. Dutrifoy, A. Majda: *The dynamics of equatorial long waves: a singular limit with fast variable coefficients*. Commun. Math. Sci., **4**, 375–397 (2006)
- [16] A. Dutrifoy, A. Majda, S. Schochet: *A simple justification of the singular limit for equatorial shallow-water dynamics*. Comm. Pure Appl. Math., **62**, 322–333 (2009)
- [17] F. Fanelli: *Highly rotating viscous compressible fluids in presence of capillarity effects*. Math. Ann., **366**, 981–1033 (2016)
- [18] F. Fanelli: *A singular limit problem for rotating capillary fluids with variable rotation axis*. J. Math. Fluid Mech., **18**, 625–658 (2016)
- [19] E. Feireisl, I. Gallagher, D. Gérard-Varet, A. Novotný: *Multi-scale analysis of compressible viscous and rotating fluids*. Comm. Math. Phys. **314**, 641–670 (2012)
- [20] E. Feireisl, I. Gallagher, A. Novotný: *A singular limit for compressible rotating fluids*. SIAM J. Math. Anal. **44**, 192–205 (2012)
- [21] E. Feireisl, A. Novotný: “*Singular limits in thermodynamics of viscous fluids*”. Advances in Mathematical Fluid Mechanics, Birkhäuser Verlag, Basel (2009)
- [22] I. Gallagher: *Applications of Schochet’s methods to parabolic equations*. J. Math. Pures Appl. **77**, 989–1054 (1998)
- [23] I. Gallagher: *A mathematical review of the analysis of the betaplane model and equatorial waves*. Discrete Contin. Dyn. Syst. Ser. S **1**, 461–480 (2008)
- [24] I. Gallagher, L. Saint-Raymond: *Weak convergence results for inhomogeneous rotating fluid equations*. J. Anal. Math. **99**, 1–34 (2006)
- [25] I. Gallagher, L. Saint-Raymond: *Mathematical study of the betaplane model: equatorial waves and convergence results*. Mém. Soc. Math. Fr. **107** (2006)
- [26] A. E. Gill: “*Atmosphere-ocean dynamics*”. Academic Press, Orlando (1982)
- [27] M. Lesieur: “*Turbulence in fluids*”. Fluid Mechanics and its applications, Springer, Dordrecht, The Netherlands (2008)
- [28] J. Leray: *Sur le mouvement d’un liquid visqueux emplissant l’espace*. Acta Math. **63**, 193–248 (1934)
- [29] P.-L. Lions: “*Mathematical topics in Fluid Mechanics. Vol. 1. Incompressible models*”. Oxford Lecture Series in Mathematics, Oxford University Press, New York (1996)
- [30] P.-L. Lions, N. Masmoudi: *Incompressible limit for a viscous compressible fluid*. J. Math. Pures Appl. **77**, 585–627 (1998)

- [31] A. Majda: *“Introduction to PDEs and waves for the atmosphere and ocean”*. Courant Lecture Notes in Mathematics, **9**, New York University, Courant Institute of Mathematical Sciences, New York; American Mathematical Society, Providence, RI (2003)
- [32] J. Pedlosky: *“Geophysical fluid dynamics”*. Springer-Verlag, New-York (1987)